



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 07:51 AM EDT

PDB ID : 7BBD  
Title : Crystal structure of monoubiquitinated TRIM21 RING (Ub-RING) In complex with ubiquitin charged Ube2N (Ube2N Ub) and Ube2V2  
Authors : Kiss, L.; Neuhaus, D.; James, L.C.  
Deposited on : 2020-12-17  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	164	 12% 87% 10% ..
2	D	76	 5% 92% 8%
3	A	150	 41% 91% 8% ..
4	C	153	 96% .

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 4434 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyubiquitin-B,E3 ubiquitin-protein ligase TRIM21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	160	1241	776	222	231	12	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP J3QS39
B	-1	SER	-	expression tag	UNP J3QS39
B	0	HIS	-	expression tag	UNP J3QS39
B	75	ALA	-	linker	UNP J3QS39
B	76	ALA	-	linker	UNP J3QS39

- Molecule 2 is a protein called Polyubiquitin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	76	597	376	104	116	1	0	0	0

- Molecule 3 is a protein called Ubiquitin-conjugating enzyme E2 variant 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	149	1099	690	187	214	8	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q15819
A	-3	SER	-	expression tag	UNP Q15819
A	-2	GLN	-	expression tag	UNP Q15819
A	-1	GLU	-	expression tag	UNP Q15819
A	0	PHE	-	expression tag	UNP Q15819

- Molecule 4 is a protein called Ubiquitin-conjugating enzyme E2 N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	C	153	1210	777	209	220	4	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	GLY	-	expression tag	UNP P61088
C	87	LYS	CYS	engineered mutation	UNP P61088
C	92	ALA	LYS	engineered mutation	UNP P61088

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
5	B	2	2	2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	87	Total 87	O 87	0	0
6	D	22	Total 22	O 22	0	0
6	A	48	Total 48	O 48	0	0
6	C	128	Total 128	O 128	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Polyubiquitin-B,E3 ubiquitin-protein ligase TRIM21




- Molecule 2: Polyubiquitin-C



- Molecule 3: Ubiquitin-conjugating enzyme E2 variant 2



- Molecule 4: Ubiquitin-conjugating enzyme E2 N



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.15Å 108.36Å 75.14Å 90.00° 104.99° 90.00°	Depositor
Resolution (Å)	19.99 – 2.20 19.99 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (19.99-2.20) 99.6 (19.99-2.20)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.19Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874, PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.222 , 0.252 0.222 , 0.252	Depositor DCC
$R_{free}$ test set	2004 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.0	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 64.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4434	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.24	0/1259	0.44	0/1699
2	D	0.24	0/603	0.44	0/811
3	A	0.24	0/1123	0.39	0/1529
4	C	0.25	0/1240	0.43	0/1688
All	All	0.24	0/4225	0.42	0/5727

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1241	0	1260	9	0
2	D	597	0	623	4	0
3	A	1099	0	1018	10	0
4	C	1210	0	1226	4	0
5	B	2	0	0	0	0
6	A	48	0	0	1	0
6	B	87	0	0	1	0
6	C	128	0	0	1	0
6	D	22	0	0	0	0
All	All	4434	0	4127	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:14:ARG:NH2	3:A:111:ASN:O	2.24	0.70
1:B:0:HIS:HB3	1:B:16:GLU:HG2	1.76	0.68
3:A:126:MET:O	6:A:201:HOH:O	2.16	0.62
1:B:23:ILE:HB	1:B:52:ASP:HA	1.81	0.61
1:B:46:ALA:O	6:B:301:HOH:O	2.16	0.59
1:B:70:VAL:HG11	1:B:73:LEU:HG	1.84	0.58
4:C:7:ARG:NH2	4:C:99:LEU:O	2.37	0.56
3:A:28:ASP:N	3:A:28:ASP:OD1	2.37	0.53
3:A:53:PRO:O	3:A:56:THR:OG1	2.28	0.52
4:C:93:ASP:OD2	6:C:201:HOH:O	2.19	0.50
1:B:24:GLU:HB3	1:B:52:ASP:HB3	1.93	0.49
2:D:45:PHE:O	2:D:48:LYS:HG2	2.13	0.48
3:A:63:TYR:HD1	3:A:84:THR:HG1	1.61	0.47
3:A:52:GLY:HA3	3:A:58:TYR:O	2.15	0.46
3:A:57:ASN:HB2	3:A:132:MET:O	2.16	0.46
2:D:23:ILE:HB	2:D:52:ASP:HA	1.97	0.46
1:B:44:ILE:HB	1:B:68:HIS:HB2	2.00	0.44
1:B:14:THR:O	1:B:33:LYS:NZ	2.44	0.44
3:A:128:SER:O	3:A:132:MET:HB2	2.19	0.43
2:D:73:LEU:HD22	4:C:107:SER:HB3	2.00	0.43
1:B:37:PRO:HB2	1:B:39:ASP:OD1	2.18	0.42
1:B:99:PHE:HB3	1:B:102:PRO:HB3	2.00	0.42
3:A:52:GLY:HA2	3:A:53:PRO:HD3	1.88	0.41
2:D:11:LYS:HB3	2:D:11:LYS:HE2	1.91	0.41
3:A:56:THR:HA	3:A:132:MET:HG3	2.03	0.40
4:C:41:GLY:HA3	4:C:47:PHE:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	158/164 (96%)	151 (96%)	7 (4%)	0	100	100
2	D	74/76 (97%)	73 (99%)	1 (1%)	0	100	100
3	A	147/150 (98%)	142 (97%)	5 (3%)	0	100	100
4	C	151/153 (99%)	148 (98%)	3 (2%)	0	100	100
All	All	530/543 (98%)	514 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	140/145 (97%)	137 (98%)	3 (2%)	53	67
2	D	67/68 (98%)	67 (100%)	0	100	100
3	A	111/133 (84%)	110 (99%)	1 (1%)	78	88
4	C	128/128 (100%)	128 (100%)	0	100	100
All	All	446/474 (94%)	442 (99%)	4 (1%)	78	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	71	LEU
1	B	73	LEU
1	B	113	GLN
3	A	14	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	160/164 (97%)	0.73	20 (12%) <b>3</b>   <b>3</b>	17, 48, 95, 115	0
2	D	76/76 (100%)	0.64	4 (5%) 26   25	23, 56, 88, 96	0
3	A	149/150 (99%)	1.81	61 (40%) <b>0</b>   <b>0</b>	23, 84, 126, 140	0
4	C	153/153 (100%)	-0.28	0 <b>100</b>   <b>100</b>	7, 20, 52, 66	0
All	All	538/543 (99%)	0.73	85 (15%) <b>2</b>   <b>1</b>	7, 48, 116, 140	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	138	PRO	8.0
3	A	129	LYS	6.8
3	A	90	GLY	6.2
1	B	73	LEU	5.8
3	A	132	MET	5.6
3	A	94	SER	5.6
3	A	128	SER	5.2
3	A	140	GLY	5.2
1	B	10	GLY	5.0
3	A	96	GLY	4.9
3	A	95	SER	4.7
3	A	139	GLU	4.6
3	A	97	MET	4.5
3	A	60	ASN	4.5
1	B	75	ALA	4.5
3	A	91	ILE	4.5
3	A	89	ASN	4.3
3	A	93	ASN	4.2
3	A	135	PRO	4.1
3	A	131	ASN	4.1
3	A	144	ASN	4.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	76	ALA	4.1
1	B	77	MET	4.1
3	A	105	VAL	4.0
3	A	86	ILE	4.0
1	B	78	ALA	4.0
2	D	57	SER	4.0
3	A	102	SER	4.0
3	A	115	ILE	3.9
3	A	124	ARG	3.9
3	A	143	TYR	3.8
3	A	142	THR	3.8
1	B	71	LEU	3.8
3	A	99	ASP	3.7
3	A	127	MET	3.6
3	A	136	GLN	3.6
3	A	54	PRO	3.5
1	B	158	GLU	3.4
3	A	61	ARG	3.3
3	A	122	LEU	3.3
3	A	55	ARG	3.2
1	B	156	SER	3.2
3	A	141	GLN	3.1
3	A	26	VAL	3.0
3	A	88	MET	3.0
3	A	104	PRO	3.0
3	A	-4	GLY	2.9
3	A	109	TRP	2.9
3	A	126	MET	2.9
3	A	113	TYR	2.9
3	A	59	GLU	2.9
1	B	9	THR	2.8
2	D	1	MET	2.8
3	A	27	GLY	2.7
3	A	51	ILE	2.6
3	A	100	ALA	2.6
3	A	101	ARG	2.6
1	B	122	GLY	2.6
3	A	87	ASN	2.6
3	A	92	ASN	2.6
3	A	130	GLU	2.6
1	B	157	GLN	2.6
3	A	56	THR	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	A	107	ALA	2.5
1	B	34	GLU	2.5
1	B	80	ALA	2.5
3	A	57	ASN	2.5
3	A	28	ASP	2.5
3	A	133	LYS	2.4
2	D	51	GLU	2.4
3	A	98	VAL	2.4
3	A	31	VAL	2.4
1	B	81	ALA	2.3
1	B	74	ARG	2.3
1	B	16	GLU	2.3
1	B	13	ILE	2.3
1	B	38	PRO	2.2
1	B	49	GLN	2.2
3	A	110	GLN	2.2
3	A	121	GLU	2.2
3	A	85	LYS	2.2
3	A	29	GLY	2.1
2	D	2	GLN	2.1
3	A	45	ARG	2.0
3	A	82	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	ZN	B	201	1/1	0.99	0.04	26,26,26,26	0


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ZN	B	202	1/1	0.99	0.04	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers ⓘ

There are no such residues in this entry.