



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 3, 2024 – 09:32 PM EDT

PDB ID : 9BBG
Title : Co-crystal structure of human DDB1 bound to fragment UB028671
Authors : Zeng, H.; Dong, A.; Frommlet, A.; Seitova, A.; Loppnau, P.; Ackloo, S.; Arrow-smith, C.H.; Edwards, A.M.; Halabelian, L.; Structural Genomics Consortium (SGC)
Deposited on : 2024-04-05
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1142	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	UNX	A	1210	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DNA damage-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1111	8673	5513	1456	1655	49	0	18	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q16531
A	0	SER	-	expression tag	UNP Q16531

- Molecule 2 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	4	6	0	0
2	A	1	10	4	6	0	0

- Molecule 3 is 1H-indol-6-amine (three-letter code: A1ALA) (formula: $C_8H_8N_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
3	A	1	20	16	4	0	1

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

- Molecule 5 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total X 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	586	Total O 590 590	0	4

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA damage-binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.14Å 125.75Å 169.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.01 – 1.70 46.97 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.01-1.70) 99.9 (46.97-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.182 , 0.218 0.191 , 0.224	Depositor DCC
R_{free} test set	7350 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9329	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, TLA, EDO, A1ALA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	1/8829 (0.0%)	0.74	6/11980 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	958	GLU	CD-OE1	5.50	1.31	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	18	CYS	CB-CA-C	6.56	123.52	110.40
1	A	57	MET	CG-SD-CE	6.52	110.63	100.20
1	A	477	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	931	LEU	N-CA-CB	-5.19	100.02	110.40
1	A	889	ARG	NE-CZ-NH1	-5.03	117.79	120.30
1	A	477	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1080	ARG	Sidechain
1	A	198[A]	ARG	Sidechain
1	A	327	ARG	Sidechain
1	A	626	ARG	Sidechain
1	A	928	ARG	Sidechain
1	A	947	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8673	0	8481	51	1
2	A	20	0	8	0	0
3	A	20	0	0	0	0
4	A	24	0	35	2	0
5	A	2	0	0	0	0
6	A	590	0	0	3	0
All	All	9329	0	8524	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198[A]:ARG:HG2	1:A:198[A]:ARG:HH11	1.04	1.14
1:A:275:ASP:OD2	1:A:279[B]:ARG:HD3	1.66	0.95
1:A:127:GLU:OE2	6:A:1301:HOH:O	1.93	0.86
1:A:198[A]:ARG:HG2	1:A:198[A]:ARG:NH1	1.80	0.85
1:A:413:LEU:HD21	1:A:461:GLY:HA2	1.60	0.82
1:A:413:LEU:CD2	1:A:461:GLY:HA2	2.19	0.71
1:A:118:THR:O	1:A:118:THR:HG22	1.90	0.71
1:A:413:LEU:HD21	1:A:461:GLY:CA	2.20	0.71
1:A:1105:MET:O	1:A:1109:VAL:HG13	1.94	0.67
1:A:742:VAL:HG13	1:A:752:LEU:HD21	1.78	0.65
1:A:118:THR:O	1:A:118:THR:CG2	2.49	0.60
1:A:275:ASP:OD2	1:A:279[B]:ARG:CD	2.45	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:277:GLU:OE1	1:A:279[A]:ARG:NH1	2.35	0.59
1:A:390:ILE:HG12	1:A:712:ILE:HD12	1.85	0.58
1:A:554:PRO:HB2	1:A:570:LYS:HE3	1.88	0.56
1:A:596:PHE:HB3	1:A:661:SER:HB2	1.88	0.55
1:A:277:GLU:OE1	1:A:279[A]:ARG:NH2	2.41	0.53
4:A:1209:EDO:H12	6:A:1428:HOH:O	2.07	0.53
1:A:502[B]:SER:OG	6:A:1302:HOH:O	2.19	0.52
1:A:434:ARG:NH1	1:A:445:GLU:OE2	2.42	0.52
1:A:146[B]:ASP:OD2	1:A:148:ASP:HB2	2.11	0.51
1:A:1102:ARG:N	1:A:1103:PRO:CD	2.74	0.51
1:A:198[A]:ARG:NH1	1:A:198[A]:ARG:CG	2.62	0.51
1:A:18:CYS:SG	1:A:315:THR:HG23	2.50	0.51
1:A:740:ILE:HG21	1:A:784:GLU:HA	1.93	0.51
1:A:1080:ARG:HB3	1:A:1080:ARG:NH2	2.28	0.47
1:A:277:GLU:OE1	1:A:279[A]:ARG:CZ	2.63	0.47
1:A:516:LEU:HD11	1:A:558:ILE:HD12	1.96	0.47
1:A:642:ARG:HG2	1:A:647[B]:THR:OG1	2.15	0.47
1:A:184:ASP:HB2	1:A:185:PRO:CD	2.45	0.46
1:A:749:THR:HG21	1:A:786:VAL:HG11	1.97	0.46
1:A:288:GLU:HB2	1:A:298:LYS:HB2	1.98	0.45
1:A:384:GLU:O	4:A:1209:EDO:H21	2.17	0.45
1:A:690:SER:C	1:A:691:LEU:HD12	2.36	0.45
1:A:1136:LEU:O	1:A:1139:ILE:HG12	2.17	0.45
1:A:1030:PHE:CZ	1:A:1038:GLY:HA3	2.52	0.44
1:A:18:CYS:SG	1:A:315:THR:CG2	3.06	0.44
1:A:141:LYS:HB2	1:A:141:LYS:HE2	1.77	0.42
1:A:1057:ARG:HD3	1:A:1108:VAL:O	2.19	0.42
1:A:218:MET:HE1	1:A:258:ILE:HG22	2.01	0.42
1:A:1109:VAL:O	1:A:1110:ALA:C	2.58	0.42
1:A:1109:VAL:HG11	1:A:1126:ALA:HA	2.01	0.42
1:A:630:THR:HG23	1:A:1134:GLU:OE1	2.19	0.41
1:A:1104:LYS:O	1:A:1108:VAL:HG23	2.21	0.41
1:A:207:TRP:HB2	1:A:242:GLY:HA2	2.03	0.41
1:A:317:LEU:O	1:A:318:ASP:HB2	2.21	0.41
1:A:219:VAL:HG22	1:A:231:ILE:CD1	2.51	0.41
1:A:1109:VAL:HG12	1:A:1129:LEU:HD12	2.02	0.40
1:A:390:ILE:HG12	1:A:712:ILE:CD1	2.49	0.40
1:A:385:GLY:HA3	1:A:719:GLU:O	2.22	0.40
1:A:255:GLN:OE1	1:A:279[A]:ARG:NH2	2.54	0.40
1:A:242:GLY:C	1:A:244:LYS:H	2.24	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:840:GLU:OE2	1:A:1080:ARG:NH2[4_445]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1117/1142 (98%)	1087 (97%)	30 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	934/1000 (93%)	926 (99%)	8 (1%)	78 70

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	183	GLN
1	A	186	GLN
1	A	198[A]	ARG
1	A	198[B]	ARG
1	A	244	LYS
1	A	245	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	969	GLU
1	A	1080	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	GLN
1	A	648	ASN
1	A	990	GLN
1	A	991	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are unknown - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	1207	-	3,3,3	0.18	0	2,2,2	0.19	0
3	A1ALA	A	1203[A]	-	9,11,11	0.77	0	12,15,15	0.68	0
4	EDO	A	1209	-	3,3,3	0.45	0	2,2,2	0.63	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A1ALA	A	1203[B]	-	9,11,11	0.79	0	12,15,15	0.69	0
4	EDO	A	1204	-	3,3,3	0.80	0	2,2,2	0.41	0
4	EDO	A	1208	-	3,3,3	0.50	0	2,2,2	0.17	0
2	TLA	A	1202	-	9,9,9	1.12	0	12,12,12	1.01	0
2	TLA	A	1201	-	9,9,9	1.38	2 (22%)	12,12,12	0.89	0
4	EDO	A	1206	-	3,3,3	0.63	0	2,2,2	0.24	0
4	EDO	A	1205	-	3,3,3	0.31	0	2,2,2	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1207	-	-	0/1/1/1	-
3	A1ALA	A	1203[A]	-	-	-	0/2/2/2
4	EDO	A	1209	-	-	1/1/1/1	-
3	A1ALA	A	1203[B]	-	-	-	0/2/2/2
4	EDO	A	1204	-	-	0/1/1/1	-
4	EDO	A	1208	-	-	1/1/1/1	-
2	TLA	A	1202	-	-	0/12/12/12	-
4	EDO	A	1206	-	-	1/1/1/1	-
2	TLA	A	1201	-	-	0/12/12/12	-
4	EDO	A	1205	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	TLA	O1-C1	2.58	1.30	1.22
2	A	1201	TLA	O11-C1	-2.05	1.23	1.30

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1208	EDO	O1-C1-C2-O2
4	A	1209	EDO	O1-C1-C2-O2
4	A	1205	EDO	O1-C1-C2-O2
4	A	1206	EDO	O1-C1-C2-O2


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1209	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1111/1142 (97%)	0.13	50 (4%) 33 37	22, 35, 65, 115	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	46	ALA	6.4
1	A	1114	TYR	5.5
1	A	1109	VAL	4.4
1	A	750	THR	4.4
1	A	338	VAL	4.1
1	A	1120	MET	3.8
1	A	344	GLY	3.8
1	A	1119	GLY	3.8
1	A	768	SER	3.7
1	A	342	GLU	3.6
1	A	1121	LYS	3.5
1	A	1115	ASP	3.5
1	A	1112	LEU	3.4
1	A	94	SER	3.2
1	A	784	GLU	3.1
1	A	1123	GLU	3.1
1	A	359	ILE	3.0
1	A	687	TYR	3.0
1	A	403	ASP	2.9
1	A	700[A]	THR	2.9
1	A	841	ALA	2.9
1	A	749	THR	2.9
1	A	148	ASP	2.8
1	A	981	SER	2.8
1	A	339	ASP	2.7
1	A	1037	ILE	2.7
1	A	767	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1111	ASN	2.5
1	A	343	GLN	2.5
1	A	769	LYS	2.4
1	A	987	GLU	2.4
1	A	318	ASP	2.4
1	A	7	VAL	2.3
1	A	71	GLY	2.3
1	A	1122	ARG	2.3
1	A	586	ILE	2.2
1	A	1022	THR	2.2
1	A	8	THR	2.2
1	A	837	TYR	2.2
1	A	980	ASP	2.2
1	A	1006	VAL	2.2
1	A	548	ASP	2.2
1	A	1030	PHE	2.1
1	A	418	ASN	2.1
1	A	346	TYR	2.1
1	A	340	SER	2.1
1	A	1124	ALA	2.1
1	A	356	LEU	2.1
1	A	686	GLY	2.1
1	A	697	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	UNX	A	1210	1/1	0.66	0.44	41,41,41,41	0
4	EDO	A	1207	4/4	0.71	0.12	67,68,71,73	0
4	EDO	A	1209	4/4	0.73	0.24	37,47,50,51	0
2	TLA	A	1201	10/10	0.75	0.33	36,58,63,65	0
2	TLA	A	1202	10/10	0.79	0.28	45,62,65,67	0
3	A1ALA	A	1203[B]	10/10	0.83	0.33	43,63,70,71	10
3	A1ALA	A	1203[A]	10/10	0.83	0.33	35,39,40,41	10
4	EDO	A	1206	4/4	0.89	0.14	32,41,44,53	0
4	EDO	A	1204	4/4	0.91	0.22	22,38,39,52	0
5	UNX	A	1211	1/1	0.91	0.29	37,37,37,37	0
4	EDO	A	1205	4/4	0.94	0.10	52,53,54,54	0
4	EDO	A	1208	4/4	0.95	0.07	34,40,42,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1ALA A 1203 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.