



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 09:24 pm GMT

PDB ID : 4BF2  
Title : Crystal Structures of Ask1-inhibitor Complexes  
Authors : Singh, O.; Shillings, A.; Craggs, P.; Wall, I.; Rowland, P.; Skarzynski, T.;  
Hobbs, C.I.; Hardwick, P.; Tanner, R.; Blunt, M.; Witty, D.R.; Smith, K.J.  
Deposited on : 2013-03-13  
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtrriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	5449 (2.14-2.10)
Clashscore	122126	5972 (2.14-2.10)
Ramachandran outliers	120053	5914 (2.14-2.10)
Sidechain outliers	120020	5915 (2.14-2.10)
RSRZ outliers	108989	5321 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	
1	B	334	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	253	1973	1270	325	369	9	0	0	0
1	B	252	1961	1262	325	365	9	0	0	0

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	644	MET	-	expression tag	UNP Q99683
A	645	ASP	-	expression tag	UNP Q99683
A	646	TYR	-	expression tag	UNP Q99683
A	647	LYS	-	expression tag	UNP Q99683
A	648	ASP	-	expression tag	UNP Q99683
A	649	ASP	-	expression tag	UNP Q99683
A	650	ASP	-	expression tag	UNP Q99683
A	651	ASP	-	expression tag	UNP Q99683
A	652	LYS	-	expression tag	UNP Q99683
A	653	GLU	-	expression tag	UNP Q99683
A	654	ASN	-	expression tag	UNP Q99683
A	655	LEU	-	expression tag	UNP Q99683
A	656	TYR	-	expression tag	UNP Q99683
A	657	PHE	-	expression tag	UNP Q99683
A	658	GLN	-	expression tag	UNP Q99683
A	659	GLY	-	expression tag	UNP Q99683
A	838	GLU	THR	engineered mutation	UNP Q99683
B	644	MET	-	expression tag	UNP Q99683
B	645	ASP	-	expression tag	UNP Q99683
B	646	TYR	-	expression tag	UNP Q99683
B	647	LYS	-	expression tag	UNP Q99683
B	648	ASP	-	expression tag	UNP Q99683
B	649	ASP	-	expression tag	UNP Q99683
B	650	ASP	-	expression tag	UNP Q99683

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	651	ASP	-	expression tag	UNP Q99683
B	652	LYS	-	expression tag	UNP Q99683
B	653	GLU	-	expression tag	UNP Q99683
B	654	ASN	-	expression tag	UNP Q99683
B	655	LEU	-	expression tag	UNP Q99683
B	656	TYR	-	expression tag	UNP Q99683
B	657	PHE	-	expression tag	UNP Q99683
B	658	GLN	-	expression tag	UNP Q99683
B	659	GLY	-	expression tag	UNP Q99683
B	838	GLU	THR	engineered mutation	UNP Q99683

- Molecule 2 is STAUROSPORINE (three-letter code: STU) (formula:  $C_{28}H_{26}N_4O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			35	28	4	3		
2	B	1	Total	C	N	O	0	0
			35	28	4	3		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2^-$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	131	Total 131	O 131	0	0
5	B	89	Total 89	O 89	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5



- Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.25Å 78.25Å 421.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	13.36 – 2.11 33.45 – 2.11	Depositor EDS
% Data completeness (in resolution range)	93.4 (13.36-2.11) 93.0 (33.45-2.11)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 2.12Å)	Xtrriage
Refinement program	BUSTER 2.11.1	Depositor
R, $R_{free}$	0.206 , 0.247 0.219 , 0.264	Depositor DCC
$R_{free}$ test set	1728 reflections (4.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.4	Xtrriage
Anisotropy	0.152	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 61.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4234	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, STU, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/2015	0.63	0/2717
1	B	0.46	0/2002	0.63	0/2699
All	All	0.49	0/4017	0.63	0/5416

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1941	6	0
1	B	1961	0	1945	6	0
2	A	35	0	26	2	0
2	B	35	0	26	1	0
3	A	6	0	8	0	0
4	B	4	0	3	0	0
5	A	131	0	0	0	0
5	B	89	0	0	1	0
All	All	4234	0	3949	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1941:STU:H261	2:B:1941:STU:H16	1.71	0.73
1:B:869:CYS:HB3	1:B:880:PRO:HG3	1.77	0.66
1:B:679:GLU:H	1:B:679:GLU:CD	2.06	0.57
1:A:821:SER:OG	1:A:822:ASP:N	2.36	0.56
1:B:846:MET:HG2	1:B:850:ILE:HB	1.87	0.56
1:A:869:CYS:HB3	1:A:880:PRO:HG3	1.91	0.52
2:A:1940:STU:H261	2:A:1940:STU:H16	1.92	0.52
1:A:786:GLN:NE2	1:A:818:LEU:H	2.09	0.51
1:A:786:GLN:HE22	1:A:818:LEU:H	1.60	0.49
1:B:872:ILE:HD11	1:B:915:ILE:HG21	1.99	0.45
1:B:676:GLU:HG2	5:B:2004:HOH:O	2.17	0.43
1:A:675:TYR:OH	1:A:710:GLU:OE2	2.30	0.41
2:A:1940:STU:C17	2:A:1940:STU:H273	2.50	0.41
1:A:786:GLN:HE22	1:A:817:VAL:HA	1.86	0.40
1:B:706:ILE:HD12	1:B:753:PHE:HD1	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	247/334 (74%)	237 (96%)	9 (4%)	1 (0%)	36 34
1	B	246/334 (74%)	237 (96%)	9 (4%)	0	100 100
All	All	493/668 (74%)	474 (96%)	18 (4%)	1 (0%)	49 50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	822	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/289 (71%)	202 (98%)	4 (2%)	60	65
1	B	206/289 (71%)	200 (97%)	6 (3%)	45	47
All	All	412/578 (71%)	402 (98%)	10 (2%)	52	55

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	679	GLU
1	A	731	HIS
1	A	844	GLN
1	A	936	GLU
1	B	728	LEU
1	B	733	LYS
1	B	737	ILE
1	B	905	GLU
1	B	910	GLU
1	B	936	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	786	GLN
1	B	756	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	STU	A	1940	-	30,42,42	2.01	11 (36%)	30,68,68	1.75	9 (30%)
3	GOL	A	1941	-	5,5,5	0.54	0	5,5,5	1.40	1 (20%)
2	STU	B	1941	-	30,42,42	2.13	12 (40%)	30,68,68	1.71	5 (16%)
4	ACT	B	1942	-	1,3,3	3.51	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	STU	A	1940	-	-	0/4/42/42	0/0/8/8
3	GOL	A	1941	-	-	0/4/4/4	0/0/0/0
2	STU	B	1941	-	-	0/4/42/42	0/0/8/8
4	ACT	B	1942	-	-	0/0/0/0	0/0/0/0

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1941	STU	C6-C19	-4.00	1.37	1.42
2	A	1940	STU	C11-C18	-3.29	1.38	1.42
2	B	1941	STU	C7-C8	-2.24	1.45	1.49
2	B	1941	STU	C11-C18	-2.04	1.40	1.42
2	A	1940	STU	C3-C2	2.01	1.42	1.38
2	A	1940	STU	C3-C4	2.05	1.41	1.36

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1941	STU	C8-N1	2.12	1.37	1.35
2	A	1940	STU	C28-N4	2.15	1.52	1.46
2	B	1941	STU	O6-C22	2.31	1.46	1.42
2	A	1940	STU	C9-N1	2.40	1.48	1.45
2	B	1941	STU	C1-C20	2.41	1.46	1.41
2	B	1941	STU	C2-C1	2.65	1.42	1.36
2	B	1941	STU	C22-C23	2.69	1.55	1.52
2	A	1940	STU	C14-C13	2.86	1.43	1.36
2	B	1941	STU	C14-C13	3.09	1.43	1.36
2	A	1940	STU	C4-C5	3.09	1.47	1.41
2	B	1941	STU	C26-C21	3.12	1.55	1.51
2	A	1940	STU	C2-C1	3.14	1.44	1.36
2	B	1941	STU	C24-C25	3.41	1.56	1.51
4	B	1942	ACT	CH3-C	3.51	1.53	1.48
2	A	1940	STU	C24-C25	3.78	1.57	1.51
2	B	1941	STU	C24-C23	3.91	1.59	1.53
2	A	1940	STU	C26-C21	3.94	1.56	1.51
2	A	1940	STU	C24-C23	4.16	1.60	1.53

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1941	STU	O5-C8-C7	-5.19	123.16	129.32
2	B	1941	STU	C26-C21-C22	-4.47	103.80	112.66
2	A	1940	STU	C9-N1-C8	-4.07	109.39	113.78
2	A	1940	STU	O5-C8-C7	-3.94	124.64	129.32
2	B	1941	STU	C14-C13-C12	-3.26	115.23	120.83
2	A	1940	STU	C26-C21-C22	-2.65	107.41	112.66
3	A	1941	GOL	C3-C2-C1	-2.45	102.18	111.63
2	A	1940	STU	C3-C4-C5	-2.35	116.79	120.83
2	A	1940	STU	C9-C10-C7	-2.30	107.56	109.61
2	A	1940	STU	C27-O6-C22	-2.08	110.55	114.31
2	B	1941	STU	C9-N1-C8	-2.02	111.60	113.78
2	A	1940	STU	C11-C12-C17	2.14	108.70	106.37
2	B	1941	STU	C13-C12-C17	2.64	122.73	119.39
2	A	1940	STU	C10-C9-N1	2.66	104.92	101.86
2	A	1940	STU	O4-C25-C24	3.36	117.17	112.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1940	STU	2	0
2	B	1941	STU	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	253/334 (75%)	0.11	12 (4%) 31 37	31, 45, 76, 114	0
1	B	252/334 (75%)	0.59	31 (12%) 4 5	35, 57, 87, 104	0
All	All	505/668 (75%)	0.35	43 (8%) 11 14	31, 50, 82, 114	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	854	GLY	6.0
1	B	855	PRO	5.7
1	B	714	ARG	5.5
1	B	851	ILE	5.4
1	B	892	PHE	5.3
1	B	940	VAL	5.2
1	A	837	GLU	5.2
1	A	830	ALA	5.1
1	B	852	ASP	4.2
1	B	856	ARG	4.1
1	B	867	LEU	3.9
1	B	853	LYS	3.9
1	A	731	HIS	3.9
1	A	939	LYS	3.5
1	B	830	ALA	3.5
1	B	804	ILE	3.5
1	B	840	THR	3.3
1	B	750	ILE	3.1
1	B	787	ILE	3.0
1	A	727	ALA	2.9
1	B	748	GLY	2.9
1	B	713	GLU	2.9
1	B	871	ILE	2.8
1	A	838	GLU	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	724	GLU	2.8
1	B	712	PRO	2.7
1	B	939	LYS	2.7
1	B	691	TYR	2.7
1	B	809	VAL	2.7
1	A	839	PHE	2.7
1	B	889	ALA	2.6
1	A	828	ARG	2.4
1	B	886	GLU	2.4
1	B	857	GLY	2.4
1	B	724	GLU	2.3
1	B	747	ASN	2.2
1	B	870	THR	2.2
1	A	829	LEU	2.2
1	B	885	GLY	2.2
1	B	820	ILE	2.1
1	A	690	THR	2.1
1	B	711	ILE	2.1
1	A	934	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ACT	B	1942	4/4	0.71	0.24	64,64,65,66	0
3	GOL	A	1941	6/6	0.96	0.14	50,51,51,51	0
2	STU	B	1941	35/35	0.97	0.09	37,38,39,41	0
2	STU	A	1940	35/35	0.98	0.13	32,33,34,35	0



## 6.5 Other polymers [i](#)

There are no such residues in this entry.