



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 2, 2024 – 12:09 PM EDT

PDB ID : 9BFG  
Title : Structure of the crosslinked PCP-E didomain of tyrocidine synthetase A  
Authors : Heberlig, G.W.; Burkart, M.D.  
Deposited on : 2024-04-17  
Resolution : 2.18 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	8336 (2.20-2.16)
Clashscore	180529	9404 (2.20-2.16)
Ramachandran outliers	177936	9297 (2.20-2.16)
Sidechain outliers	177891	9297 (2.20-2.16)
RSRZ outliers	164620	8337 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	 4% 90% 6%
1	B	555	 2% 88% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	612	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 17521 atoms, of which 8525 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrocidine synthase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	533	8460	2706	4204	733	809	8	129	0	0
1	B	532	8449	2706	4198	730	807	8	131	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP P09095
A	2	VAL	-	expression tag	UNP P09095
A	370	ALA	GLU	engineered mutation	UNP P09095
A	546	GLY	-	expression tag	UNP P09095
A	547	SER	-	expression tag	UNP P09095
A	548	ARG	-	expression tag	UNP P09095
A	549	SER	-	expression tag	UNP P09095
A	550	HIS	-	expression tag	UNP P09095
A	551	HIS	-	expression tag	UNP P09095
A	552	HIS	-	expression tag	UNP P09095
A	553	HIS	-	expression tag	UNP P09095
A	554	HIS	-	expression tag	UNP P09095
A	555	HIS	-	expression tag	UNP P09095
B	1	MET	-	expression tag	UNP P09095
B	2	VAL	-	expression tag	UNP P09095
B	370	ALA	GLU	engineered mutation	UNP P09095
B	546	GLY	-	expression tag	UNP P09095
B	547	SER	-	expression tag	UNP P09095
B	548	ARG	-	expression tag	UNP P09095
B	549	SER	-	expression tag	UNP P09095
B	550	HIS	-	expression tag	UNP P09095
B	551	HIS	-	expression tag	UNP P09095
B	552	HIS	-	expression tag	UNP P09095
B	553	HIS	-	expression tag	UNP P09095
B	554	HIS	-	expression tag	UNP P09095

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	555	HIS	-	expression tag	UNP P09095

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	B	1	5	4	1	0	0

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
3	A	1	25	6	13	1	4	1	0	0
3	B	1	25	6	13	1	4	1	0	0
3	B	1	25	6	13	1	4	1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
5	A	1	49	14	26	2	6	1	0
5	B	1	49	14	26	2	6	1	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	167	Total	O	0	0
			167	167		
6	B	156	Total	O	0	0
			156	156		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrocidine synthase 1



- Molecule 1: Tyrocidine synthase 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	216.38Å 216.38Å 59.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.85 – 2.18 46.85 – 2.18	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.85-2.18) 100.0 (46.85-2.18)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.18Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.188 , 0.228 0.188 , 0.229	Depositor DCC
$R_{free}$ test set	4242 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.3	Xtrriage
Anisotropy	0.243	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 34.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	17521	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2GH, GOL, MES, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	0/4352	0.81	0/5910
1	B	0.70	0/4348	0.83	0/5905
All	All	0.70	0/8700	0.82	0/11815

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4256	4204	4180	22	2
1	B	4251	4198	4172	24	2
2	A	40	0	0	0	0
2	B	20	0	0	1	0
3	A	12	13	13	1	0
3	B	24	26	26	2	0
4	A	12	16	16	6	0
4	B	12	16	16	1	0
5	A	23	26	27	3	0
5	B	23	26	28	5	0
6	A	167	0	0	5	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	156	0	0	5	0
All	All	8996	8525	8478	48	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 48 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:TYR:CB	6:A:855:HOH:O	2.01	1.08
4:A:605:GOL:O1	5:A:610:2GH:H26	1.82	0.78
3:A:603:MES:H52	6:A:773:HOH:O	1.92	0.70
1:B:196:GLU:HG2	6:B:807:HOH:O	1.91	0.69
1:B:196:GLU:CG	6:B:807:HOH:O	2.42	0.67

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:SER:HG	1:B:364:GLN:HE22[5_554]	1.11	0.49
1:A:75:ASN:OD1	1:B:441:SER:OG[5_554]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	531/555 (96%)	517 (97%)	14 (3%)	0	100	100
1	B	530/555 (96%)	515 (97%)	15 (3%)	0	100	100
All	All	1061/1110 (96%)	1032 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/484 (96%)	453 (98%)	9 (2%)	52	64
1	B	461/484 (95%)	450 (98%)	11 (2%)	44	54
All	All	923/968 (95%)	903 (98%)	20 (2%)	47	58

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	168	ASN
1	B	401	LYS
1	B	487	SER
1	B	418	ARG
1	A	367	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	HIS
1	B	531	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	611	-	4,4,4	0.26	0	6,6,6	0.23	0
4	GOL	B	606	-	5,5,5	0.13	0	5,5,5	0.40	0
3	MES	B	605	-	12,12,12	0.71	0	15,16,16	0.53	0
2	SO4	A	609	-	4,4,4	0.43	0	6,6,6	0.24	0
2	SO4	B	607	-	4,4,4	0.31	0	6,6,6	0.12	0
4	GOL	A	612	-	5,5,5	0.18	0	5,5,5	0.49	0
2	SO4	B	602	-	4,4,4	0.30	0	6,6,6	0.11	0
2	SO4	B	604	-	4,4,4	0.31	0	6,6,6	0.20	0
2	SO4	A	607	-	4,4,4	0.28	0	6,6,6	0.10	0
5	2GH	B	608	1	18,22,23	0.63	0	21,28,31	1.62	5 (23%)
2	SO4	B	601	-	4,4,4	0.34	0	6,6,6	0.11	0
4	GOL	B	609	-	5,5,5	0.11	0	5,5,5	0.36	0
2	SO4	A	604	-	4,4,4	0.26	0	6,6,6	0.19	0
2	SO4	A	608	-	4,4,4	0.29	0	6,6,6	0.15	0
5	2GH	A	610	1	18,22,23	0.37	0	21,28,31	0.94	2 (9%)
2	SO4	A	601	-	4,4,4	0.34	0	6,6,6	0.10	0
3	MES	B	603	-	12,12,12	0.67	0	15,16,16	0.74	1 (6%)
2	SO4	A	602	-	4,4,4	0.30	0	6,6,6	0.15	0
3	MES	A	603	-	12,12,12	0.76	0	15,16,16	0.96	1 (6%)
4	GOL	A	605	-	5,5,5	0.22	0	5,5,5	0.41	0
2	SO4	A	606	-	4,4,4	0.30	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	B	603	-	-	0/6/14/14	0/1/1/1
4	GOL	B	609	-	-	2/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	606	-	-	2/4/4/4	-
3	MES	B	605	-	-	3/6/14/14	0/1/1/1
5	2GH	B	608	1	-	12/26/28/29	-
5	2GH	A	610	1	-	4/26/28/29	-
3	MES	A	603	-	-	6/6/14/14	0/1/1/1
4	GOL	A	612	-	-	4/4/4/4	-
4	GOL	A	605	-	-	2/4/4/4	-

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	608	2GH	CAN-CAO-CAT	3.52	118.26	112.39
5	B	608	2GH	OAE-CAU-CAV	-3.21	111.94	120.89
5	B	608	2GH	CAB-CAW-CAP	3.08	113.31	108.22
5	B	608	2GH	CAM-NAQ-CAT	2.57	127.61	122.82
3	B	603	MES	O2S-S-C8	-2.29	103.27	106.73

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	603	MES	C8-C7-N4-C5
3	A	603	MES	N4-C7-C8-S
3	A	603	MES	C7-C8-S-O1S
3	A	603	MES	C7-C8-S-O3S
3	B	605	MES	C7-C8-S-O1S

There are no ring outliers.

8 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	605	MES	2	0
4	A	612	GOL	4	0
5	B	608	2GH	5	0
2	B	601	SO4	1	0
4	B	609	GOL	1	0
5	A	610	2GH	3	0
3	A	603	MES	1	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	605	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	533/555 (96%)	-0.03	23 (4%) 40 39	24, 41, 76, 115	0
1	B	532/555 (95%)	-0.03	13 (2%) 59 58	26, 41, 72, 111	0
All	All	1065/1110 (95%)	-0.03	36 (3%) 48 47	24, 41, 74, 115	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	GLY	5.3
1	A	301	VAL	4.8
1	B	310	THR	4.3
1	B	162	GLY	3.9
1	B	209	GLY	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	607	5/5	0.65	0.16	106,111,119,127	0
3	MES	B	605	12/12	0.72	0.20	71,82,131,152	0
2	SO4	A	609	5/5	0.75	0.15	58,63,79,81	0
4	GOL	A	612	6/6	0.79	0.19	60,63,70,71	2
4	GOL	B	609	6/6	0.80	0.18	56,65,79,80	2
2	SO4	A	607	5/5	0.82	0.10	97,109,121,122	0
2	SO4	A	608	5/5	0.85	0.11	70,72,76,90	0
4	GOL	A	605	6/6	0.85	0.21	56,59,67,67	2
2	SO4	A	611	5/5	0.87	0.11	78,85,89,95	0
2	SO4	A	604	5/5	0.87	0.11	64,67,74,80	0
2	SO4	B	604	5/5	0.91	0.09	57,64,75,83	0
2	SO4	A	602	5/5	0.91	0.07	67,75,78,81	0
4	GOL	B	606	6/6	0.91	0.19	46,59,77,78	2
2	SO4	B	601	5/5	0.91	0.07	67,70,75,78	0
5	2GH	B	608	23/24	0.92	0.12	37,47,58,64	1
2	SO4	A	606	5/5	0.93	0.11	110,113,123,126	0
2	SO4	A	601	5/5	0.94	0.06	78,79,83,88	0
2	SO4	B	602	5/5	0.94	0.07	62,63,77,77	0
5	2GH	A	610	23/24	0.96	0.09	33,40,60,64	1
3	MES	A	603	12/12	0.96	0.09	34,39,42,44	0
3	MES	B	603	12/12	0.98	0.05	30,33,34,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 2GH B 608:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 2GH A 610:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.