



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 03:11 am BST

PDB ID : 3BGT
Title : Structural Studies of Acetoacetate Decarboxylase
Authors : Ho, M.; Allen, K.N.
Deposited on : 2007-11-27
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	
1	B	246	
1	C	246	
1	D	246	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8082 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable acetoacetate decarboxylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	244	Total 1924	C 1241	N 327	O 346	S 2	Se 8	0	2	0
1	B	244	Total 1913	C 1235	N 323	O 345	S 2	Se 8	0	1	0
1	C	244	Total 1913	C 1235	N 323	O 345	S 2	Se 8	0	1	0
1	D	245	Total 1924	C 1241	N 327	O 346	S 2	Se 8	0	1	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	114	Total 114	O 114	0	0
2	B	122	Total 122	O 122	0	0
2	C	86	Total 86	O 86	0	0
2	D	86	Total 86	O 86	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Probable acetoacetate decarboxylase




- Molecule 1: Probable acetoacetate decarboxylase




- Molecule 1: Probable acetoacetate decarboxylase



- Molecule 1: Probable acetoacetate decarboxylase





4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	105.45Å 105.45Å 252.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.30 – 2.10 30.28 – 1.99	Depositor EDS
% Data completeness (in resolution range)	94.3 (30.30-2.10) 93.1 (30.28-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005, CNS	Depositor
R, R_{free}	0.201 , 0.245 0.196 , 0.237	Depositor DCC
R_{free} test set	11784 reflections (8.81%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 58.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for -h,1/3*h-1/3*k-1/3*l,-4/3*h-8/3*k +1/3*l 0.009 for -1/3*h+1/3*k+1/3*l,-k,8/3*h+4/ 3*k+1/3*l 0.004 for -2/3*h-1/3*k-1/3*l,-1/3*h-2/3*k+ 1/3*l,-4/3*h+4/3*k+1/3*l 0.002 for 1/3*h+2/3*k-1/3*l,-k,-8/3*h-4/3* k-1/3*l 0.005 for -1/3*h-2/3*k+1/3*l,-2/3*h-1/3*k- 1/3*l,4/3*h-4/3*k-1/3*l 0.003 for -h,2/3*h+1/3*k+1/3*l,4/3*h+8/3 *k-1/3*l 0.020 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8082	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.59	46/1972 (2.3%)	1.05	11/2677 (0.4%)
1	B	1.01	12/1961 (0.6%)	0.89	7/2663 (0.3%)
1	C	1.10	23/1961 (1.2%)	0.88	8/2663 (0.3%)
1	D	1.31	38/1972 (1.9%)	1.00	12/2677 (0.4%)
All	All	1.27	119/7866 (1.5%)	0.96	38/10680 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

The worst 5 of 119 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	90	ARG	CB-CG	-16.57	1.07	1.52
1	B	89	GLU	CB-CG	-13.94	1.25	1.52
1	B	31	GLU	CD-OE1	-13.66	1.10	1.25
1	A	205	GLU	CB-CG	-12.84	1.27	1.52
1	B	31	GLU	CD-OE2	-12.84	1.11	1.25

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	155	ARG	NE-CZ-NH1	17.62	129.11	120.30
1	D	155	ARG	NE-CZ-NH2	-11.43	114.59	120.30
1	C	191	ASP	CB-CG-OD1	10.64	127.87	118.30
1	A	90	ARG	NE-CZ-NH1	-9.59	115.51	120.30
1	B	31	GLU	OE1-CD-OE2	-8.30	113.34	123.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	90	ARG	CA

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1924	0	1923	57	3
1	B	1913	0	1911	54	0
1	C	1913	0	1911	42	0
1	D	1924	0	1924	39	0
2	A	114	0	0	3	0
2	B	122	0	0	3	0
2	C	86	0	0	2	3
2	D	86	0	0	5	0
All	All	8082	0	7669	189	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 189 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:GLU:HG2	1:B:97[B]:MSE:CE	1.51	1.38
1:D:54:MSE:CG	1:D:54:MSE:SE	2.24	1.36
1:A:77:GLU:HG2	1:A:97[B]:MSE:CE	1.56	1.35
1:D:54:MSE:CE	1:D:54:MSE:SE	2.24	1.35
1:A:77:GLU:CG	1:A:97[B]:MSE:HE3	1.66	1.24

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:TYR:OH	2:C:301:HOH:O[2_765]	0.18	2.02
1:A:150:TYR:CZ	2:C:301:HOH:O[2_765]	1.17	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:TYR:CE2	2:C:301:HOH:O[2_765]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/246 (99%)	236 (97%)	8 (3%)	0	100	100
1	B	243/246 (99%)	233 (96%)	10 (4%)	0	100	100
1	C	243/246 (99%)	233 (96%)	10 (4%)	0	100	100
1	D	244/246 (99%)	233 (96%)	10 (4%)	1 (0%)	34	32
All	All	974/984 (99%)	935 (96%)	38 (4%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	159	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/198 (104%)	187 (91%)	18 (9%)	10	6
1	B	204/198 (103%)	192 (94%)	12 (6%)	19	17
1	C	204/198 (103%)	188 (92%)	16 (8%)	12	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	205/198 (104%)	193 (94%)	12 (6%)	19	17
All	All	818/792 (103%)	760 (93%)	58 (7%)	15	11

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	99	LEU
1	C	52	LEU
1	D	155	ARG
1	B	161	SER
1	B	234	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	GLN
1	B	102	GLN
1	D	166	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/246 (96%)	-0.24	3 (1%) 77 80	19, 31, 48, 71	3 (1%)
1	B	237/246 (96%)	-0.32	1 (0%) 92 93	18, 29, 54, 68	3 (1%)
1	C	237/246 (96%)	-0.07	4 (1%) 70 74	25, 37, 57, 79	2 (0%)
1	D	238/246 (96%)	-0.05	6 (2%) 57 62	25, 39, 63, 80	2 (0%)
All	All	949/984 (96%)	-0.17	14 (1%) 73 77	18, 34, 57, 80	10 (1%)

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	62	GLU	2.9
1	C	8	GLN	2.8
1	D	4	GLN	2.6
1	D	62	GLU	2.5
1	D	245	ARG	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.