

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 25, 2023 – 07:24 AM EDT

PDB ID : 3BK9

Title: H55A mutant of tryptophan 2,3-dioxygenase from Xanthomonas campestris

Authors: Bruckmann, C.; Mowat, C.G.

Deposited on : 2007-12-06

Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\# \text{Entries, resolution range}(\mathring{A}))$			
$R_{free}$	130704	1479 (2.16-2.16)			
Clashscore	141614	1585 (2.16-2.16)			
Ramachandran outliers	138981	1560 (2.16-2.16)			
Sidechain outliers	138945	1559 (2.16-2.16)			
RSRZ outliers	127900	1456 (2.16-2.16)			

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain							
1	A	306	64%	20% 6% • 9%						
1	В	306	60%	25% 6% 8%						
1	С	306	58%	25% 5% 11%						
1	D	306	59%	23% • 14%						
1	Е	306	53%	26% 5% 15%						



Continued from previous page...

Mol	Chain	Length	Quality of chain							
1	F	306	55%	23%		18%				
1	G	306	5%	27%	5% •	14%				
1	Н	306	59%	21%	•	16%				



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 19073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Tryptophan 2,3-dioxygenase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	277	Total	С	N	О	S	0	0	0
1	A	211	2285	1462	404	412	7	0	0	
1	В	280	Total	С	N	О	S	0	0	0
1	1 B	200	2308	1475	408	418	7	0	0	
1	С	272	Total	С	N	О	S	0	0	0
1		212	2244	1437	395	405	7	0		
1	D	262	Total	С	N	О	S	0	0	0
1	D	202	2165	1389	381	388	7	U	O	0
1	E	260	Total	С	N	Ο	S	0	0	0
1	ш	200	2153	1381	379	386	7	0	U	
1	F	250	Total	С	N	О	S	0	0	0
1	I.	250	2071	1330	362	372	7	0	0	
1	G	262	Total	С	N	О	S	0	0	0
1	1 G	202	2172	1393	384	388	7	0	0	
1	Н	257	Total	С	N	О	S	0	0	0
1	11	201	2120	1362	372	379	7			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	ALA	HIS	engineered mutation	UNP Q8PDA8
A	299	LEU	-	expression tag	UNP Q8PDA8
A	300	GLU	ı	expression tag	UNP Q8PDA8
A	301	HIS	-	expression tag	UNP Q8PDA8
A	302	HIS	-	expression tag	UNP Q8PDA8
A	303	HIS	-	expression tag	UNP Q8PDA8
A	304	HIS	-	expression tag	UNP Q8PDA8
A	305	HIS	-	expression tag	UNP Q8PDA8
A	306	HIS	-	expression tag	UNP Q8PDA8
В	55	ALA	HIS	engineered mutation	UNP Q8PDA8
В	299	LEU	-	expression tag	UNP Q8PDA8
В	300	GLU	-	expression tag	UNP Q8PDA8
В	301	HIS	-	expression tag	UNP Q8PDA8



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	302	HIS	-	expression tag	UNP Q8PDA8
В	303	HIS	-	expression tag	UNP Q8PDA8
В	304	HIS	-	expression tag	UNP Q8PDA8
В	305	HIS	_	expression tag	UNP Q8PDA8
В	306	HIS	-	expression tag	UNP Q8PDA8
С	55	ALA	HIS	engineered mutation	UNP Q8PDA8
С	299	LEU	-	expression tag	UNP Q8PDA8
С	300	GLU	-	expression tag	UNP Q8PDA8
С	301	HIS	-	expression tag	UNP Q8PDA8
С	302	HIS	-	expression tag	UNP Q8PDA8
С	303	HIS	-	expression tag	UNP Q8PDA8
С	304	HIS	-	expression tag	UNP Q8PDA8
С	305	HIS	-	expression tag	UNP Q8PDA8
С	306	HIS	-	expression tag	UNP Q8PDA8
D	55	ALA	HIS	engineered mutation	UNP Q8PDA8
D	299	LEU	-	expression tag	UNP Q8PDA8
D	300	GLU	-	expression tag	UNP Q8PDA8
D	301	HIS	-	expression tag	UNP Q8PDA8
D	302	HIS	-	expression tag	UNP Q8PDA8
D	303	HIS	-	expression tag	UNP Q8PDA8
D	304	HIS	-	expression tag	UNP Q8PDA8
D	305	HIS	-	expression tag	UNP Q8PDA8
D	306	HIS	-	expression tag	UNP Q8PDA8
Е	55	ALA	HIS	engineered mutation	UNP Q8PDA8
Е	299	LEU	-	expression tag	UNP Q8PDA8
Е	300	GLU	-	expression tag	UNP Q8PDA8
Е	301	HIS	-	expression tag	UNP Q8PDA8
Е	302	HIS	-	expression tag	UNP Q8PDA8
Е	303	HIS	-	expression tag	UNP Q8PDA8
Е	304	HIS	-	expression tag	UNP Q8PDA8
Е	305	HIS	-	expression tag	UNP Q8PDA8
Е	306	HIS	_	expression tag	UNP Q8PDA8
F	55	ALA	HIS	engineered mutation	UNP Q8PDA8
F	299	LEU	-	expression tag	UNP Q8PDA8
F	300	GLU	-	expression tag	UNP Q8PDA8
F	301	HIS	-	expression tag	UNP Q8PDA8
F	302	HIS	-	expression tag	UNP Q8PDA8
F	303	HIS	-	expression tag	UNP Q8PDA8
F	304	HIS	-	expression tag	UNP Q8PDA8
F	305	HIS	-	expression tag	UNP Q8PDA8
F	306	HIS	-	expression tag	UNP Q8PDA8
G	55	ALA	HIS	engineered mutation	UNP Q8PDA8



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
G	299	LEU	-	expression tag	UNP Q8PDA8
G	300	GLU	-	expression tag	UNP Q8PDA8
G	301	HIS	-	expression tag	UNP Q8PDA8
G	302	HIS	ı	expression tag	UNP Q8PDA8
G	303	HIS	-	expression tag	UNP Q8PDA8
G	304	HIS	ı	expression tag	UNP Q8PDA8
G	305	HIS	-	expression tag	UNP Q8PDA8
G	306	HIS	-	expression tag	UNP Q8PDA8
Н	55	ALA	HIS	engineered mutation	UNP Q8PDA8
Н	299	LEU	-	expression tag	UNP Q8PDA8
Н	300	GLU	-	expression tag	UNP Q8PDA8
Н	301	HIS	-	expression tag	UNP Q8PDA8
Н	302	HIS	-	expression tag	UNP Q8PDA8
Н	303	HIS	-	expression tag	UNP Q8PDA8
Н	304	HIS		expression tag	UNP Q8PDA8
Н	305	HIS	=	expression tag	UNP Q8PDA8
Н	306	HIS	-	expression tag	UNP Q8PDA8

 $\bullet$  Molecule 2 is TRYPTOPHAN (three-letter code: TRP) (formula:  $C_{11}H_{12}N_2O_2).$ 




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total				0	0	
	2 11	1	15	11	2	2			
9	Λ	1	Total	$\mathbf{C}$	Ν	O	0	0	
	2 A	1	15	11	2	2	0		
9	D	1	Total	С	N	О	0	0	
	$2 \mid B$	1	15	11	2	2	U	0	



Continued from previous page...

Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf	
2	В	1	Total	С	N	О	0	0	
	Б	1	15	11	2	2	U	0	
2	С	1	Total	С	N	Ο	0	0	
		1	15	11	2	2	U	U	
2	С	1	Total	С	N	Ο	0	0	
		1	15	11	2	2	0	U	
2	D	1	Total	С	N	Ο	0	0	
	D	1	15	11	2	2	0	U	
2	D	1	Total	С	N	Ο	0	0	
	D	D I	15	11	2	2	O	U	
$\frac{1}{2}$	E	1	Total	$\mathbf{C}$	N	Ο	0	0	
	Ш	1	15	11	2	2	O	U	
$\frac{1}{2}$	E	1	Total	$\mathbf{C}$	N	Ο	0	0	
	Ш	1	15	11	2	2	O	U	
$\frac{1}{2}$	F	1	Total	$\mathbf{C}$	N	Ο	0	0	
	I.	1	15	11	2	2	O	U	
2	G	1	Total	С	N	O	0	0	
	2 G	1	15	11	2	2	U	U	
2	2 H	П	1	Total	С	N	Ο	0	0
	11	1	15	11	2	2	U		

 $\bullet$  Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $\rm C_{34}H_{32}FeN_4O_4).$ 



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Λ	1	Total	С	Fe	N	О	0	0
3	A	1	43	34	1	4	4	0	



Continued from previous page...

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
3	В	1	Total	С	Fe	N	О	0	0	
3	9 В	1	43	34	1	4	4	0	0	
3	С	1	Total	С	Fe	N	О	0	0	
3		1	43	34	1	4	4	0	U	
3	D	1	Total	С	Fe	N	О	0	0	
3	<i>D</i>	D	1	43	34	1	4	4		U
3	F	E	1	Total	С	Fe	N	О	0	0
3	ت ا	1	43	34	1	4	4		U	
3	F	1	Total	С	Fe	N	О	0	0	
3	I'	1	43	34	1	4	4		0	
3	G	1	Total	С	Fe	N	Ο	0	0	
	3   G	1	43	34	1	4	4		U	
3	Н	Ц 1		С	Fe	N	О	0	0	
3	3   H	1	43	34	1	4	4		U	

#### • Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	170	Total O 170 170	0	0
4	В	152	Total O 152 152	0	0
4	С	129	Total O 129 129	0	0
4	D	124	Total O 124 124	0	0
4	E	133	Total O 133 133	0	0
4	F	96	Total O 96 96	0	0
4	G	113	Total O 113 113	0	0
4	Н	99	Total O 99 99	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tryptophan 2,3-dioxygenase










• Molecule 1: Tryptophan 2,3-dioxygenase








 $\bullet$  Molecule 1: Tryptophan 2,3-dioxygenase









# 4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	78.22Å 117.61Å 139.28Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 95.73° 90.00°	Depositor	
Resolution (Å)	54.13 - 2.15	Depositor	
rtesolution (A)	54.13 - 2.15	EDS	
% Data completeness	96.7 (54.13-2.15)	Depositor	
(in resolution range)	96.7 (54.13-2.15)	EDS	
$R_{merge}$	0.10	Depositor	
$R_{sym}$	(Not available)	Depositor	
$< I/\sigma(I) > 1$	3.20 (at 2.14Å)	Xtriage	
Refinement program	REFMAC 5.2.0019	Depositor	
P. P.	0.200 , 0.285	Depositor	
$R, R_{free}$	0.201 , 0.281	DCC	
$R_{free}$ test set	6624 reflections (5.03%)	wwPDB-VP	
Wilson B-factor (Å <sup>2</sup> )	22.6	Xtriage	
Anisotropy	0.136	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, 46.9	EDS	
L-test for twinning <sup>2</sup>	$ < L >=0.46, < L^2>=0.29$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
$F_o, F_c$ correlation	0.93	EDS	
Total number of atoms	19073	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.40% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain D		ond lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5
1	A	1.46	15/2342~(0.6%)	1.35	26/3174 (0.8%)
1	В	1.52	27/2365 (1.1%)	1.50	24/3206 (0.7%)
1	С	1.35	$13/2299 \ (0.6\%)$	1.31	19/3115 (0.6%)
1	D	1.41	$12/2220 \ (0.5\%)$	1.17	9/3009 (0.3%)
1	Е	1.36	$14/2208 \; (0.6\%)$	1.24	$14/2993 \; (0.5\%)$
1	F	1.28	5/2122~(0.2%)	1.16	13/2872 (0.5%)
1	G	1.27	5/2227~(0.2%)	1.21	14/3018 (0.5%)
1	Н	1.18	2/2173~(0.1%)	1.05	5/2944 (0.2%)
All	All	1.36	93/17956 (0.5%)	1.26	124/24331 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0
1	В	0	1
1	G	0	1
All	All	1	2

The worst 5 of 93 bond length outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
1	В	161	GLU	CG-CD	10.31	1.67	1.51
1	С	106	GLU	CB-CG	8.86	1.69	1.52
1	A	17	ASP	CB-CG	7.96	1.68	1.51
1	D	106	GLU	CB-CG	7.73	1.66	1.52
1	D	106	GLU	CD-OE2	7.31	1.33	1.25

The worst 5 of 124 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	252	ARG	NE-CZ-NH2	-23.35	108.62	120.30
1	В	252	ARG	NE-CZ-NH1	19.05	129.83	120.30
1	В	8	ARG	NE-CZ-NH2	-15.85	112.38	120.30
1	В	8	ARG	NE-CZ-NH1	13.25	126.92	120.30
1	С	279	ARG	NE-CZ-NH2	12.71	126.65	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	17	ASP	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	283	GLY	Peptide
1	G	251	LYS	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2285	0	2240	70	0
1	В	2308	0	2259	58	0
1	С	2244	0	2201	66	0
1	D	2165	0	2123	64	0
1	E	2153	0	2109	69	0
1	F	2071	0	2033	58	0
1	G	2172	0	2133	77	0
1	Н	2120	0	2083	60	0
2	A	30	0	18	3	0
2	В	30	0	18	0	0
2	С	30	0	18	3	0
2	D	30	0	18	0	0
2	Е	30	0	18	1	0
2	F	15	0	9	0	0
2	G	15	0	9	1	0
2	Н	15	0	9	2	0
3	A	43	0	30	1	0



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	43	0	30	4	0
3	С	43	0	30	2	0
3	D	43	0	30	4	0
3	Е	43	0	30	7	0
3	F	43	0	30	5	0
3	G	43	0	30	3	0
3	Н	43	0	30	5	0
4	A	170	0	0	4	0
4	В	152	0	0	6	0
4	С	129	0	0	7	0
4	D	124	0	0	5	0
4	Е	133	0	0	6	0
4	F	96	0	0	2	0
4	G	113	0	0	11	0
4	Н	99	0	0	1	0
All	All	19073	0	17538	470	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 470 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:E:114:MET:SD	1:H:283:GLY:HA2	1.77	1.24
1:E:38:GLN:NE2	1:F:29:ARG:HH11	1.37	1.23
1:E:38:GLN:HE22	1:F:29:ARG:NH1	1.38	1.20
1:A:241:MET:CE	1:A:245:MET:HE2	1.75	1.16
1:C:241:MET:SD	1:C:245:MET:HE3	1.86	1.15

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	275/306~(90%)	269 (98%)	5 (2%)	1 (0%)	34	29
1	В	278/306 (91%)	268 (96%)	9 (3%)	1 (0%)	34	29
1	$\mathbf{C}$	268/306~(88%)	258 (96%)	9 (3%)	1 (0%)	34	29
1	D	260/306~(85%)	250 (96%)	9 (4%)	1 (0%)	34	29
1	E	258/306 (84%)	245 (95%)	13 (5%)	0	100	100
1	F	246/306 (80%)	235 (96%)	9 (4%)	2 (1%)	19	12
1	G	260/306~(85%)	247 (95%)	12 (5%)	1 (0%)	34	29
1	Н	253/306 (83%)	241 (95%)	12 (5%)	0	100	100
All	All	2098/2448 (86%)	2013 (96%)	78 (4%)	7 (0%)	41	37

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	284	VAL
1	A	118	ASP
1	G	114	MET
1	С	198	ALA
1	D	184	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	241/266~(91%)	225 (93%)	16 (7%)	16 11
1	В	244/266~(92%)	229 (94%)	15 (6%)	18 14
1	С	238/266~(90%)	220 (92%)	18 (8%)	13 8
1	D	228/266~(86%)	214 (94%)	14 (6%)	18 14
1	E	227/266~(85%)	204 (90%)	23 (10%)	7 4
1	F	220/266~(83%)	204 (93%)	16 (7%)	14 9
1	G	229/266 (86%)	211 (92%)	18 (8%)	12 7
1	Н	224/266 (84%)	205 (92%)	19 (8%)	10 6



Continued from previous page...

Mol	Chain	Analysed	Analysed Rotameric Out		Percentiles
All	All	$1851/2128 \; (87\%)$	1712 (92%)	139 (8%)	13 8

5 of 139 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	235	LEU
1	G	270	THR
1	Н	151	TYR
1	С	264	GLN
1	С	257	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 74 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	185	GLN
1	Н	130	GLN
1	G	37	GLN
1	G	188	GLN
1	С	83	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TRP	Н	402	-	14,16,16	1.30	2 (14%)	16,22,22	1.09	2 (12%)
2	TRP	С	403	-	14,16,16	1.15	1 (7%)	16,22,22	1.32	2 (12%)
2	TRP	D	403	-	14,16,16	1.32	2 (14%)	16,22,22	1.28	3 (18%)
2	TRP	Е	403	-	14,16,16	1.35	3 (21%)	16,22,22	0.87	0
2	TRP	В	403	-	14,16,16	0.91	1 (7%)	16,22,22	1.49	5 (31%)
2	TRP	Е	402	-	14,16,16	1.58	3 (21%)	16,22,22	1.43	2 (12%)
3	HEM	F	401	1	41,50,50	2.51	13 (31%)	45,82,82	2.69	20 (44%)
3	HEM	G	401	1	41,50,50	1.92	11 (26%)	45,82,82	2.62	15 (33%)
3	HEM	В	401	4,1	41,50,50	2.33	11 (26%)	45,82,82	2.03	12 (26%)
2	TRP	A	403	_	14,16,16	1.30	3 (21%)	16,22,22	1.07	1 (6%)
2	TRP	F	402	-	14,16,16	1.51	4 (28%)	16,22,22	1.39	2 (12%)
3	HEM	Е	401	4,1	41,50,50	2.22	9 (21%)	45,82,82	2.58	16 (35%)
2	TRP	G	402	-	14,16,16	1.37	2 (14%)	16,22,22	1.06	2 (12%)
2	TRP	A	402	-	14,16,16	1.78	2 (14%)	16,22,22	1.28	2 (12%)
3	HEM	С	401	1	41,50,50	2.16	12 (29%)	45,82,82	2.27	16 (35%)
3	HEM	D	401	1	41,50,50	2.12	11 (26%)	45,82,82	2.28	19 (42%)
3	HEM	Н	401	1	41,50,50	2.23	11 (26%)	45,82,82	2.40	18 (40%)
2	TRP	D	402	-	14,16,16	1.85	4 (28%)	16,22,22	1.19	2 (12%)
2	TRP	С	402	-	14,16,16	1.64	2 (14%)	16,22,22	0.96	2 (12%)
2	TRP	В	402	_	14,16,16	1.57	3 (21%)	16,22,22	1.05	2 (12%)
3	HEM	A	401	1	41,50,50	2.09	11 (26%)	45,82,82	2.51	19 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
2	TRP	Н	402	-	-	5/7/8/8	0/2/2/2
2	TRP	С	403	-	-	0/7/8/8	0/2/2/2
2	TRP	D	403	-	-	0/7/8/8	0/2/2/2
2	TRP	Ε	403	-	-	1/7/8/8	0/2/2/2
2	TRP	В	403	-	-	0/7/8/8	0/2/2/2



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRP	E	402	-	-	3/7/8/8	0/2/2/2
3	HEM	F	401	1	-	3/12/54/54	-
3	HEM	G	401	1	-	4/12/54/54	-
3	HEM	В	401	4,1	-	6/12/54/54	-
2	TRP	A	403	-	-	0/7/8/8	0/2/2/2
2	TRP	F	402	-	-	5/7/8/8	0/2/2/2
3	HEM	Е	401	4,1	-	4/12/54/54	-
2	TRP	G	402	-	-	4/7/8/8	0/2/2/2
2	TRP	A	402	-	-	5/7/8/8	0/2/2/2
3	HEM	С	401	1	-	2/12/54/54	-
3	HEM	D	401	1	-	3/12/54/54	-
3	HEM	Н	401	1	-	3/12/54/54	-
2	TRP	D	402	-	-	5/7/8/8	0/2/2/2
2	TRP	С	402	-	-	6/7/8/8	0/2/2/2
2	TRP	В	402	-	-	5/7/8/8	0/2/2/2
3	HEM	A	401	1	-	4/12/54/54	-

The worst 5 of 121 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	Ideal(A)
3	Н	401	HEM	C3D-C2D	8.89	1.55	1.36
3	F	401	HEM	C3D-C2D	8.79	1.55	1.36
3	В	401	HEM	C3C-C2C	-8.72	1.28	1.40
3	Е	401	HEM	C3D-C2D	8.42	1.54	1.36
3	A	401	HEM	C3D-C2D	6.98	1.51	1.36

The worst 5 of 162 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
3	F	401	HEM	C4D-ND-C1D	9.93	115.33	105.07
3	Ε	401	HEM	C4D-ND-C1D	8.96	114.33	105.07
3	G	401	HEM	C4D-ND-C1D	8.46	113.81	105.07
3	Н	401	HEM	C4D-ND-C1D	8.20	113.54	105.07
3	G	401	HEM	C4C-CHD-C1D	7.43	132.36	122.56

There are no chirality outliers.

5 of 68 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	В	402	TRP	C-CA-CB-CG
2	D	402	TRP	C-CA-CB-CG
2	Е	402	TRP	C-CA-CB-CG
2	F	402	TRP	C-CA-CB-CG
2	G	402	TRP	C-CA-CB-CG


There are no ring outliers.

14 monomers are involved in 40 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Н	402	TRP	2	0
2	Е	403	TRP	1	0
3	F	401	HEM	5	0
3	G	401	HEM	3	0
3	В	401	HEM	4	0
2	A	403	TRP	1	0
3	Е	401	HEM	7	0
2	G	402	TRP	1	0
2	A	402	TRP	2	0
3	С	401	HEM	2	0
3	D	401	HEM	4	0
3	Н	401	HEM	5	0
2	С	402	TRP	3	0
3	A	401	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































# 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	277/306 (90%)	-0.07	4 (1%) 75 80	5, 15, 27, 45	0
1	В	280/306 (91%)	-0.03	5 (1%) 68 75	4, 15, 31, 42	0
1	С	272/306 (88%)	0.06	6 (2%) 62 69	10, 19, 41, 54	0
1	D	262/306~(85%)	0.01	0 100 100	9, 19, 38, 44	0
1	E	260/306 (84%)	0.12	7 (2%) 54 63	9, 24, 37, 43	0
1	F	250/306 (81%)	0.32	11 (4%) 34 43	15, 26, 45, 55	0
1	G	262/306~(85%)	0.21	15 (5%) 23 32	9, 25, 42, 47	0
1	Н	257/306 (83%)	0.63	27 (10%) 6 9	16, 31, 52, 63	0
All	All	2120/2448 (86%)	0.15	75 (3%) 44 52	4, 22, 41, 63	0

The worst 5 of 75 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	185	GLN	4.6
1	Н	119	VAL	4.4
1	С	19	GLU	4.3
1	Н	150	ALA	4.3
1	Н	192	TRP	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ extbf{B-factors}( extbf{A}^2)$	Q<0.9
2	TRP	Н	402	15/15	0.82	0.24	32,36,40,41	0
2	TRP	Е	402	15/15	0.84	0.20	29,31,33,36	0
2	TRP	G	402	15/15	0.85	0.24	30,35,40,43	0
2	TRP	Е	403	15/15	0.86	0.19	24,27,38,39	0
2	TRP	С	402	15/15	0.92	0.15	13,21,28,31	0
2	TRP	D	402	15/15	0.92	0.16	14,18,25,26	0
2	TRP	С	403	15/15	0.94	0.13	14,18,27,28	0
2	TRP	F	402	15/15	0.94	0.18	25,27,32,35	0
2	TRP	В	403	15/15	0.94	0.11	12,20,26,26	0
2	TRP	A	402	15/15	0.94	0.12	14,16,23,25	0
2	TRP	A	403	15/15	0.95	0.13	14,21,28,28	0
3	HEM	G	401	43/43	0.95	0.11	20,25,31,38	0
2	TRP	В	402	15/15	0.96	0.11	7,9,18,18	0
3	HEM	Е	401	43/43	0.96	0.11	11,23,27,34	0
3	HEM	F	401	43/43	0.96	0.12	13,20,27,34	0
2	TRP	D	403	15/15	0.96	0.10	11,15,27,30	0
3	HEM	Н	401	43/43	0.96	0.12	17,23,27,28	0
3	HEM	С	401	43/43	0.97	0.11	4,10,16,19	0
3	HEM	D	401	43/43	0.97	0.12	2,9,17,21	0
3	HEM	A	401	43/43	0.98	0.12	2,8,13,17	0
3	HEM	В	401	43/43	0.98	0.12	2,5,10,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







































# 6.5 Other polymers (i)

There are no such residues in this entry.

