

Full wwPDB X-ray Structure Validation Report (i)

Jul 1, 2024 - 08:04 PM EDT

PDB ID	:	9BK2
Title	:	Crystal structure of Lactate dehydrogenase in complex with 4-((4-(1-methyl-
		1H-imidazole-2-carbonyl)phenyl)amino)-4-oxo-2-(4-(trifluoromethyl)phenyl)b
		utanoic acid (S-enantiomer, monoclinic P form)
Authors	:	Lovell, S.; Cooper, A.; Battaile, K.P.; Sharma, H.; Seattle Structural Genomics
		Center for Infectious Disease (SSGCID)
Deposited on	:	2024-04-26
Resolution	:	1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : $4.02b-467$	
Mogul : $1.8.5 (274361), \text{CSD as}541\text{be} (2020)$	
Xtriage (Phenix) : 1.13	
EDS : 2.37.1	
buster-report : $1.1.7$ (2018)	
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 201	19)
Refmac : 5.8.0158	
CCP4 : $7.0.044$ (Gargrove)	
Ideal geometry (proteins) : Engh & Huber (2001)	
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)	
Validation Pipeline (wwPDB-VP) : 2.37.1	




1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	2469(1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	352	88%	6%	7%
1	В	352	% 91%	•	7%
1	С	352	89%	•	7%
1	D	352	90%	•	6%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10713 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	328	Total	С	Ν	0	\mathbf{S}	0	1	0
	А	320	2510	1606	427	464	13	0	1	0
1	В	329	Total	С	Ν	0	S	0	2	0
	D	329	2505	1600	424	468	13	0	2	0
1	C	329	Total	С	Ν	0	S	0	2	0
	C	329	2497	1596	424	464	13	0		0
1	Л	330	Total	С	Ν	0	S	0	2	0
		550	2523	1614	429	466	14	0	2	

• Molecule 1 is a protein called L-lactate dehydrogenase A chain.

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-20	MET	-	initiating methionine	UNP P00338
А	-19	HIS	-	expression tag	UNP P00338
А	-18	HIS	-	expression tag	UNP P00338
А	-17	HIS	-	expression tag	UNP P00338
А	-16	HIS	-	expression tag	UNP P00338
А	-15	HIS	-	expression tag	UNP P00338
А	-14	HIS	-	expression tag	UNP P00338
А	-13	SER	-	expression tag	UNP P00338
А	-12	THR	-	expression tag	UNP P00338
А	-11	SER	-	expression tag	UNP P00338
А	-10	VAL	-	expression tag	UNP P00338
А	-9	ASP	-	expression tag	UNP P00338
A	-8	LEU	-	expression tag	UNP P00338
А	-7	GLY	-	expression tag	UNP P00338
A	-6	THR	-	expression tag	UNP P00338
А	-5	GLU	-	expression tag	UNP P00338
А	-4	ASN	-	expression tag	UNP P00338
А	-3	LEU	-	expression tag	UNP P00338
А	-2	TYR	-	expression tag	UNP P00338
А	-1	PHE	-	expression tag	UNP P00338
А	0	GLN	-	expression tag	UNP P00338



Chain	Residue	Modelled	Actual	Comment	Reference
В	-20	MET	-	initiating methionine	UNP P00338
В	-19	HIS	_	expression tag	UNP P00338
В	-18	HIS	-	expression tag	UNP P00338
В	-17	HIS	_	expression tag	UNP P00338
В	-16	HIS	_	expression tag	UNP P00338
В	-15	HIS	-	expression tag	UNP P00338
В	-14	HIS	-	expression tag	UNP P00338
В	-13	SER	-	expression tag	UNP P00338
В	-12	THR	-	expression tag	UNP P00338
В	-11	SER	-	expression tag	UNP P00338
В	-10	VAL	-	expression tag	UNP P00338
В	-9	ASP	-	expression tag	UNP P00338
В	-8	LEU	-	expression tag	UNP P00338
В	-7	GLY	-	expression tag	UNP P00338
В	-6	THR	-	expression tag	UNP P00338
В	-5	GLU	-	expression tag	UNP P00338
В	-4	ASN	-	expression tag	UNP P00338
В	-3	LEU	-	expression tag	UNP P00338
В	-2	TYR	-	expression tag	UNP P00338
В	-1	PHE	-	expression tag	UNP P00338
В	0	GLN	-	expression tag	UNP P00338
С	-20	MET	-	initiating methionine	UNP P00338
С	-19	HIS	-	expression tag	UNP P00338
С	-18	HIS	-	expression tag	UNP P00338
С	-17	HIS	-	expression tag	UNP P00338
С	-16	HIS	-	expression tag	UNP P00338
С	-15	HIS	-	expression tag	UNP P00338
С	-14	HIS	-	expression tag	UNP P00338
С	-13	SER	-	expression tag	UNP P00338
С	-12	THR	-	expression tag	UNP P00338
С	-11	SER	-	expression tag	UNP P00338
С	-10	VAL	-	expression tag	UNP P00338
С	-9	ASP	-	expression tag	UNP P00338
С	-8	LEU	_	expression tag	UNP P00338
С	-7	GLY	_	expression tag	UNP P00338
С	-6	THR	-	expression tag	UNP P00338
С	-5	GLU		expression tag	UNP P00338
С	-4	ASN	-	expression tag	UNP P00338
С	-3	LEU	-	expression tag	UNP P00338
С	-2	TYR	-	expression tag	UNP P00338
С	-1	PHE	-	expression tag	UNP P00338
С	0	GLN	_	expression tag	UNP P00338


Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
D	-20	MET	-	initiating methionine	UNP P00338
D	-19	HIS	-	expression tag	UNP P00338
D	-18	HIS	-	expression tag	UNP P00338
D	-17	HIS	-	expression tag	UNP P00338
D	-16	HIS	-	expression tag	UNP P00338
D	-15	HIS	-	expression tag	UNP P00338
D	-14	HIS	-	expression tag	UNP P00338
D	-13	SER	-	expression tag	UNP P00338
D	-12	THR	-	expression tag	UNP P00338
D	-11	SER	-	expression tag	UNP P00338
D	-10	VAL	-	expression tag	UNP P00338
D	-9	ASP	-	expression tag	UNP P00338
D	-8	LEU	-	expression tag	UNP P00338
D	-7	GLY	-	expression tag	UNP P00338
D	-6	THR	-	expression tag	UNP P00338
D	-5	GLU	-	expression tag	UNP P00338
D	-4	ASN	-	expression tag	UNP P00338
D	-3	LEU	-	expression tag	UNP P00338
D	-2	TYR	-	expression tag	UNP P00338
D	-1	PHE	-	expression tag	UNP P00338
D	0	GLN	-	expression tag	UNP P00338

Continued from previous page...

• Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues					ZeroOcc	AltConf
2	А	1	Total 4	$\begin{array}{c} \mathrm{C} \\ \mathrm{2} \end{array}$	0 1	S 1	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
2	С	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0


• Molecule 3 is (2S)-4-[4-(1-methyl-1H-imidazole-2-carbonyl)anilino]-4-oxo-2-[4-(trifluoromethyl)phenyl]butanoic acid (three-letter code: A1AQ0) (formula: $C_{22}H_{18}F_3N_3O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	\mathbf{ms}			ZeroOcc	AltConf
3	Δ	1	Total	С	F	Ν	0	0	Ο
0	11	I	32	22	3	3	4	0	0

• Molecule 4 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 7 & 3 & 4 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 7 & 3 & 4 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 3 4 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	138	Total O 138 138	0	0
5	В	145	Total O 145 145	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	С	122	Total O 122 122	0	0
5	D	155	Total O 155 155	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: L-lactate dehydrogenase A chain





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	74.28Å 99.24Å 82.66Å	Depositor
a, b, c, α , β , γ	90.00° 98.69° 90.00°	Depositor
Resolution (Å)	40.85 - 1.85	Depositor
Resolution (A)	49.62 - 1.85	EDS
% Data completeness	99.9 (40.85-1.85)	Depositor
(in resolution range)	99.9 (49.62-1.85)	EDS
R _{merge}	0.12	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.80 (at 1.86 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
D D.	0.154 , 0.192	Depositor
R, R_{free}	0.163 , 0.196	DCC
R_{free} test set	5022 reflections $(4.97%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	24.1	Xtriage
Anisotropy	0.431	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.38 , 47.3	EDS
L-test for twinning ²	$ \langle L \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10713	wwPDB-VP
Average B, all atoms $(Å^2)$	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, DMS, $\rm A1AQ0$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.38	0/2556	0.55	0/3464	
1	В	0.40	0/2554	0.56	0/3467	
1	С	0.40	0/2546	0.55	0/3458	
1	D	0.39	0/2573	0.54	0/3489	
All	All	0.39	0/10229	0.55	0/13878	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2510	0	2560	12	0
1	В	2505	0	2525	9	0
1	С	2497	0	2526	13	0
1	D	2523	0	2564	14	0
2	А	4	0	6	0	0
2	В	4	0	6	0	0
2	С	4	0	6	0	0
2	D	4	0	6	0	0
3	А	32	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes					
4	А	14	0	4	0	0					
4	В	21	0	6	0	0					
4	С	21	0	6	1	0					
4	D	14	0	4	0	0					
5	А	138	0	0	1	0					
5	В	145	0	0	0	0					
5	С	122	0	0	0	0					
5	D	155	0	0	0	0					
All	All	10713	0	10219	37	0					

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 1	A +	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:C:302:LEU:HD13	1:D:11:LEU:HD11	1.64	0.77
1:C:276:ILE:HD11	1:C:286:VAL:HG13	1.73	0.70
1:A:11:LEU:HD11	1:B:302:LEU:HD13	1.74	0.69
1:C:11:LEU:CD1	1:D:302:LEU:HD13	2.35	0.57
1:C:11:LEU:HD11	1:D:302:LEU:HD13	1.87	0.55
1:A:50:VAL:HG21	1:A:85:THR:CG2	2.38	0.54
1:D:276:ILE:HD11	1:D:286:VAL:HG13	1.90	0.52
1:C:260:GLU:OE2	1:D:72:ARG:NH1	2.42	0.52
1:D:276:ILE:HD11	1:D:286:VAL:CG1	2.41	0.50
1:A:56:LYS:NZ	1:A:60:GLU:OE2	2.47	0.48
1:A:124:VAL:HG21	1:A:150:SER:HB2	1.95	0.47
1:C:276:ILE:CD1	1:C:282:ILE:HG21	2.45	0.46
1:D:141:ILE:HD13	1:D:325:ILE:HG21	1.96	0.46
1:C:113:VAL:HG22	1:C:145:VAL:HG21	1.98	0.46
1:C:302:LEU:HD13	1:D:11:LEU:CD1	2.40	0.46
1:C:52:VAL:HG13	4:C:404:MLI:H11	1.99	0.45
1:D:50:VAL:HG21	1:D:85:THR:CG2	2.46	0.45
1:A:106:LEU:O	1:A:109:VAL:HG23	2.17	0.44
1:D:276:ILE:HD12	1:D:276:ILE:C	2.38	0.44
1:B:292:CYS:HB3	1:B:299:ILE:HG23	1.98	0.44
1:C:10:ASN:ND2	1:D:301:ASP:OD2	2.34	0.44
1:A:11:LEU:CD1	1:B:302:LEU:HD13	2.46	0.43
1:A:145:VAL:HG12	1:A:149:ILE:HD12	2.00	0.43
1:C:276:ILE:CD1	1:C:286:VAL:HG13	2.44	0.43
1:B:276:ILE:CD1	1:B:282:ILE:HD13	2.48	0.43



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:TYR:HA	1:A:181:PRO:HG3	2.01	0.43
1:B:276:ILE:HD12	1:B:282:ILE:CD1	2.49	0.43
1:B:180:HIS:HB2	1:C:266:LEU:O	2.19	0.42
1:B:276:ILE:HD12	1:B:282:ILE:HD13	2.01	0.42
5:A:528:HOH:O	1:B:264:LYS:HE3	2.20	0.42
1:A:180:HIS:HB2	1:D:266:LEU:O	2.20	0.42
1:D:165:ASP:OD1	1:D:192:HIS:ND1	2.46	0.41
1:A:30:VAL:CG2	1:A:98:ARG:CZ	2.99	0.41
1:D:12:LEU:N	1:D:12:LEU:HD23	2.36	0.41
1:B:266:LEU:O	1:C:180:HIS:HB2	2.20	0.41
1:A:293:ILE:HD12	1:A:301:ASP:HB2	2.03	0.41
1:A:200:TRP:CE3	1:A:203:MET:SD	3.15	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	325/352~(92%)	318~(98%)	7 (2%)	0	100	100
1	В	327/352~(93%)	320~(98%)	7 (2%)	0	100	100
1	С	329/352~(94%)	322~(98%)	7(2%)	0	100	100
1	D	330/352~(94%)	323~(98%)	7(2%)	0	100	100
All	All	1311/1408~(93%)	1283~(98%)	28 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	274/307~(89%)	274 (100%)	0	100	100
1	В	272/307~(89%)	271 (100%)	1 (0%)	91	89
1	С	270/307~(88%)	268 (99%)	2 (1%)	84	79
1	D	275/307~(90%)	275 (100%)	0	100	100
All	All	1091/1228~(89%)	1088 (100%)	3 (0%)	92	91

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	247	THR
1	С	89	LYS
1	С	111	ARG

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	gles
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
4	MLI	В	402	-	$6,\!6,\!6$	1.56	1 (16%)	7,7,7	1.14	0
4	MLI	С	402	-	$6,\!6,\!6$	1.36	0	7,7,7	1.25	0
4	MLI	D	403	-	$6,\!6,\!6$	1.55	1 (16%)	7,7,7	1.32	0
4	MLI	В	404	-	$6,\!6,\!6$	1.49	1 (16%)	7,7,7	1.28	1 (14%)
4	MLI	А	403	-	$6,\!6,\!6$	1.62	1 (16%)	7,7,7	1.52	1 (14%)
2	DMS	A	401	-	3,3,3	0.88	0	3,3,3	0.49	0
2	DMS	В	401	-	3,3,3	0.73	0	3,3,3	0.75	0
4	MLI	А	404	-	$6,\!6,\!6$	1.58	1 (16%)	7,7,7	1.43	1 (14%)
2	DMS	D	401	-	3,3,3	0.86	0	3,3,3	0.69	0
4	MLI	C	404	-	$6,\!6,\!6$	1.62	1 (16%)	7,7,7	0.90	0
4	MLI	D	402	-	$6,\!6,\!6$	1.57	1 (16%)	7,7,7	1.36	1 (14%)
2	DMS	С	401	-	3,3,3	0.84	0	3,3,3	0.63	0
4	MLI	В	403	-	6,6,6	1.60	1 (16%)	7,7,7	1.22	0
4	MLI	С	403	-	6,6,6	1.59	1 (16%)	7,7,7	1.32	1 (14%)
3	A1AQ0	А	402	-	32,34,34	0.56	0	46,49,49	0.90	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLI	В	402	-	-	3/4/4/4	-
4	MLI	С	402	-	-	4/4/4/4	-
4	MLI	D	403	-	-	0/4/4/4	-
4	MLI	В	404	-	-	0/4/4/4	-
4	MLI	А	403	-	-	4/4/4/4	-
4	MLI	D	402	-	-	2/4/4/4	-
4	MLI	А	404	-	-	1/4/4/4	-
4	MLI	С	404	-	-	0/4/4/4	-
4	MLI	В	403	-	-	2/4/4/4	-
4	MLI	С	403	-	-	2/4/4/4	-
3	A1AQ0	А	402	-	-	0/26/30/30	0/3/3/3



Mol	Chain	Res	Type	Atoms	Z	$Observed(\text{\AA})$	Ideal(Å)
4	С	403	MLI	C1-C3	2.84	1.55	1.51
4	В	403	MLI	C1-C3	2.80	1.55	1.51
4	А	403	MLI	C1-C3	2.76	1.55	1.51
4	D	403	MLI	C1-C3	2.69	1.55	1.51
4	D	402	MLI	C1-C3	2.58	1.55	1.51
4	С	404	MLI	C1-C3	2.53	1.55	1.51
4	А	404	MLI	C1-C3	2.48	1.54	1.51
4	В	402	MLI	C1-C3	2.38	1.54	1.51
4	В	404	MLI	C1-C3	2.23	1.54	1.51

All (9) bond length outliers are listed below:

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	403	MLI	O6-C2-C1	-2.44	114.96	122.08
3	А	402	A1AQ0	C13-C14-C16	-2.25	107.55	111.96
4	D	402	MLI	O6-C2-C1	-2.25	115.51	122.08
4	В	404	MLI	O6-C2-C1	-2.20	115.65	122.08
3	А	402	A1AQ0	C13-C14-C15	2.11	115.11	111.28
4	А	404	MLI	O6-C2-C1	-2.09	115.96	122.08
4	С	403	MLI	O6-C2-C1	-2.07	116.04	122.08

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	В	402	MLI	C3-C1-C2-O7
4	С	402	MLI	C3-C1-C2-O7
4	В	403	MLI	C3-C1-C2-O6
4	В	402	MLI	C3-C1-C2-O6
4	А	403	MLI	C2-C1-C3-O9
4	С	402	MLI	C2-C1-C3-O9
4	С	403	MLI	C3-C1-C2-O7
4	D	402	MLI	C2-C1-C3-O8
4	А	403	MLI	C2-C1-C3-O8
4	В	403	MLI	C3-C1-C2-O7
4	С	402	MLI	C3-C1-C2-O6
4	С	402	MLI	C2-C1-C3-O8
4	С	403	MLI	C3-C1-C2-O6
4	D	402	MLI	C2-C1-C3-O9
4	А	403	MLI	C3-C1-C2-O7
4	В	402	MLI	C2-C1-C3-O9



Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	А	403	MLI	C3-C1-C2-O6
4	А	404	MLI	C3-C1-C2-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	С	404	MLI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2		$OWAB(Å^2)$	Q<0.9
1	А	328/352~(93%)	-0.30	2(0%) 89	89	17, 26, 43, 63	0
1	В	329/352~(93%)	-0.45	3 (0%) 84	84	17, 24, 41, 60	0
1	С	329/352~(93%)	-0.36	1 (0%) 94	93	18, 25, 40, 55	0
1	D	330/352~(93%)	-0.32	1 (0%) 94	93	18, 24, 42, 68	0
All	All	1316/1408~(93%)	-0.36	7 (0%) 91	91	17, 25, 42, 68	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	331	PHE	6.0
1	D	331	PHE	4.7
1	В	331	PHE	4.7
1	С	329	LEU	2.8
1	В	118	PHE	2.8
1	В	12	LEU	2.3
1	А	100	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	$B-factors(A^2)$	Q<0.9
4	MLI	С	404	7/7	0.75	0.22	$33,\!41,\!55,\!58$	0
4	MLI	В	404	7/7	0.84	0.29	30,41,47,49	0
2	DMS	С	401	4/4	0.88	0.13	25,27,29,47	0
4	MLI	С	403	7/7	0.90	0.10	31,32,37,40	0
4	MLI	В	403	7/7	0.92	0.12	27,34,37,43	0
3	A1AQ0	А	402	32/32	0.92	0.14	$19,\!33,\!47,\!58$	0
2	DMS	В	401	4/4	0.93	0.12	28,29,34,39	0
4	MLI	С	402	7/7	0.93	0.15	22,26,33,38	0
4	MLI	D	402	7/7	0.93	0.09	27,30,33,36	0
2	DMS	D	401	4/4	0.94	0.12	25,28,35,40	0
2	DMS	А	401	4/4	0.94	0.15	27,28,31,40	0
4	MLI	А	403	7/7	0.94	0.11	31,34,36,39	0
4	MLI	В	402	7/7	0.95	0.12	22,26,39,39	0
4	MLI	D	403	7/7	0.95	0.16	23,31,42,48	0
4	MLI	А	404	7/7	0.96	0.10	25,27,43,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

