



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 11, 2023 – 05:04 pm BST

PDB ID : 8BLM  
Title : Structure of RutB  
Authors : Rajendran, C.  
Deposited on : 2022-11-09  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	
1	B	230	
1	C	230	
1	D	230	
1	E	230	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	230	 6% 75% 20% .
1	G	230	 11% 75% 20% ..
1	H	230	 10% 78% 18% .
1	I	230	 4% 81% 15% .
1	J	230	 6% 72% 24% .
1	K	230	 7% 74% 21% .
1	L	230	 11% 77% 17% ..
1	M	230	 11% 73% 23% .
1	N	230	 11% 79% 17% ..
1	O	230	 13% 72% 24% .
1	P	230	 15% 77% 18% .

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 28516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ureidoacrylate amidohydrolase RutB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	222	Total 1709	C 1098	N 285	O 321	S 5	0	0	0
1	B	222	Total 1715	C 1104	N 285	O 321	S 5	0	0	0
1	C	222	Total 1715	C 1104	N 285	O 321	S 5	0	0	0
1	D	220	Total 1702	C 1096	N 283	O 318	S 5	0	0	0
1	E	222	Total 1703	C 1098	N 283	O 317	S 5	0	0	0
1	F	220	Total 1694	C 1092	N 282	O 315	S 5	0	0	0
1	G	222	Total 1711	C 1101	N 284	O 321	S 5	0	0	0
1	H	222	Total 1715	C 1104	N 285	O 321	S 5	0	0	0
1	I	222	Total 1708	C 1099	N 285	O 319	S 5	0	0	0
1	J	222	Total 1696	C 1091	N 281	O 319	S 5	0	0	0
1	K	220	Total 1692	C 1091	N 282	O 314	S 5	0	0	0
1	L	220	Total 1702	C 1096	N 283	O 318	S 5	0	0	0
1	M	222	Total 1713	C 1103	N 285	O 320	S 5	0	0	0
1	N	222	Total 1711	C 1102	N 284	O 320	S 5	0	0	0
1	O	222	Total 1715	C 1104	N 285	O 321	S 5	0	0	0
1	P	220	Total 1683	C 1085	N 279	O 314	S 5	0	0	0


- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	95	Total O 95 95	0	0
2	B	87	Total O 87 87	0	0
2	C	96	Total O 96 96	0	0
2	D	81	Total O 81 81	0	0
2	E	78	Total O 78 78	0	0
2	F	70	Total O 70 70	0	0
2	G	78	Total O 78 78	0	0
2	H	81	Total O 81 81	0	0
2	I	93	Total O 93 93	0	0
2	J	84	Total O 84 84	0	0
2	K	71	Total O 71 71	0	0
2	L	82	Total O 82 82	0	0
2	M	68	Total O 68 68	0	0
2	N	48	Total O 48 48	0	0
2	O	62	Total O 62 62	0	0
2	P	58	Total O 58 58	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ureidoacrylate amidohydrolase RutB

Chain A: 





- Molecule 1: Ureidoacrylate amidohydrolase RutB

Chain B: 





- Molecule 1: Ureidoacrylate amidohydrolase RutB

Chain C: 





- Molecule 1: Ureidoacrylate amidohydrolase RutB



Chain D: 





- Molecule 1: Ureidoacrylate amidohydrolase RutB





- Molecule 1: Ureidoacrylate amidohydrolase RutB




- Molecule 1: Ureidoacrylate amidohydrolase RutB





- Molecule 1: Ureidoacrylate amidohydrolase RutB








- Molecule 1: Ureidoacrylate amidohydrolase RutB





- Molecule 1: Ureidoacrylate amidohydrolase RutB



- Molecule 1: Ureidoacrylate amidohydrolase RutB




- Molecule 1: Ureidoacrylate amidohydrolase RutB







- Molecule 1: Ureidoacrylate amidohydrolase RutB




- Molecule 1: Ureidoacrylate amidohydrolase RutB



- Molecule 1: Ureidoacrylate amidohydrolase RutB



- Molecule 1: Ureidoacrylate amidohydrolase RutB



5212	SER
D213	PRO
	THR
	SER
	PHE
	ALA
	HIS
	ILE
	ALA

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.59Å 110.89Å 128.71Å 90.00° 99.13° 90.00°	Depositor
Resolution (Å)	47.67 – 1.90 47.67 – 1.93	Depositor EDS
% Data completeness (in resolution range)	81.3 (47.67-1.90) 85.7 (47.67-1.93)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.95 (at 1.94Å)	Xtrriage
Refinement program	PHENIX 1.13-2998	Depositor
R, $R_{free}$	0.263 , 0.318 0.265 , 0.318	Depositor DCC
$R_{free}$ test set	10818 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.7	Xtrriage
Anisotropy	0.561	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 33.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	28516	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.57 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6160e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/1753	0.61	0/2392
1	B	0.46	0/1760	0.62	0/2401
1	C	0.44	0/1760	0.64	0/2401
1	D	0.46	0/1746	0.59	0/2381
1	E	0.44	0/1748	0.60	0/2386
1	F	0.42	0/1738	0.56	0/2371
1	G	0.46	0/1756	0.61	0/2397
1	H	0.45	0/1760	0.58	0/2401
1	I	0.44	0/1753	0.60	0/2392
1	J	0.44	0/1741	0.60	0/2379
1	K	0.43	0/1736	0.61	0/2369
1	L	0.46	1/1746 (0.1%)	0.57	0/2381
1	M	0.41	0/1758	0.58	0/2398
1	N	0.41	0/1756	0.57	0/2396
1	O	0.41	0/1760	0.58	0/2401
1	P	0.41	0/1727	0.56	0/2359
All	All	0.44	1/27998 (0.0%)	0.59	0/38205

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	6	ALA	C-N	-5.04	1.22	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1709	0	1668	31	0
1	B	1715	0	1675	35	1
1	C	1715	0	1675	37	0
1	D	1702	0	1663	25	0
1	E	1703	0	1659	27	0
1	F	1694	0	1653	30	0
1	G	1711	0	1664	37	0
1	H	1715	0	1675	33	0
1	I	1708	0	1662	24	0
1	J	1696	0	1632	42	1
1	K	1692	0	1648	33	0
1	L	1702	0	1663	25	1
1	M	1713	0	1670	38	0
1	N	1711	0	1669	26	0
1	O	1715	0	1675	40	0
1	P	1683	0	1627	22	1
2	A	95	0	0	1	0
2	B	87	0	0	5	0
2	C	96	0	0	4	0
2	D	81	0	0	6	0
2	E	78	0	0	6	0
2	F	70	0	0	1	0
2	G	78	0	0	4	0
2	H	81	0	0	7	0
2	I	93	0	0	5	0
2	J	84	0	0	5	0
2	K	71	0	0	6	0
2	L	82	0	0	3	0
2	M	68	0	0	3	0
2	N	48	0	0	1	0
2	O	62	0	0	4	0
2	P	58	0	0	1	0
All	All	28516	0	26578	474	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 474 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:72:ASN:ND2	1:J:133:LYS:HZ2	1.53	1.06
1:J:72:ASN:HD22	1:J:133:LYS:NZ	1.57	1.03
1:M:163:THR:HG22	1:M:188:THR:HB	1.53	0.90
1:I:172:ARG:NH2	2:I:301:HOH:O	2.11	0.82
1:O:98:MET:HE2	1:O:105:GLN:HA	1.63	0.81

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:209:GLY:O	1:P:91:LYS:NZ[2_656]	2.04	0.16
1:B:105:GLN:O	1:L:107:LYS:NZ[2_556]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/230 (96%)	214 (97%)	6 (3%)	0	100	100
1	B	220/230 (96%)	211 (96%)	9 (4%)	0	100	100
1	C	220/230 (96%)	213 (97%)	6 (3%)	1 (0%)	29	18
1	D	218/230 (95%)	211 (97%)	7 (3%)	0	100	100
1	E	220/230 (96%)	213 (97%)	7 (3%)	0	100	100
1	F	218/230 (95%)	210 (96%)	8 (4%)	0	100	100
1	G	220/230 (96%)	212 (96%)	7 (3%)	1 (0%)	29	18
1	H	220/230 (96%)	210 (96%)	10 (4%)	0	100	100
1	I	220/230 (96%)	211 (96%)	9 (4%)	0	100	100
1	J	220/230 (96%)	213 (97%)	7 (3%)	0	100	100
1	K	218/230 (95%)	205 (94%)	12 (6%)	1 (0%)	29	18
1	L	218/230 (95%)	210 (96%)	8 (4%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	M	220/230 (96%)	212 (96%)	7 (3%)	1 (0%)	29	18
1	N	220/230 (96%)	211 (96%)	9 (4%)	0	100	100
1	O	220/230 (96%)	211 (96%)	9 (4%)	0	100	100
1	P	218/230 (95%)	210 (96%)	8 (4%)	0	100	100
All	All	3510/3680 (95%)	3377 (96%)	129 (4%)	4 (0%)	51	42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	35	TYR
1	K	165	VAL
1	G	165	VAL
1	M	165	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/188 (96%)	177 (98%)	3 (2%)	60	57
1	B	181/188 (96%)	178 (98%)	3 (2%)	60	57
1	C	181/188 (96%)	178 (98%)	3 (2%)	60	57
1	D	179/188 (95%)	175 (98%)	4 (2%)	52	47
1	E	178/188 (95%)	174 (98%)	4 (2%)	52	47
1	F	177/188 (94%)	174 (98%)	3 (2%)	60	57
1	G	180/188 (96%)	178 (99%)	2 (1%)	73	73
1	H	181/188 (96%)	178 (98%)	3 (2%)	60	57
1	I	179/188 (95%)	175 (98%)	4 (2%)	52	47
1	J	176/188 (94%)	175 (99%)	1 (1%)	86	87
1	K	176/188 (94%)	173 (98%)	3 (2%)	60	57
1	L	179/188 (95%)	175 (98%)	4 (2%)	52	47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	180/188 (96%)	178 (99%)	2 (1%)	73	73
1	N	180/188 (96%)	174 (97%)	6 (3%)	38	29
1	O	181/188 (96%)	179 (99%)	2 (1%)	73	73
1	P	174/188 (93%)	169 (97%)	5 (3%)	42	35
All	All	2862/3008 (95%)	2810 (98%)	52 (2%)	59	55

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	141	ASN
1	L	58	THR
1	P	149	ARG
1	J	35	TYR
1	K	141	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	16	GLN
1	M	72	ASN
1	O	103	GLN
1	N	141	ASN
1	O	72	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	222/230 (96%)	0.79	12 (5%) 25 29	21, 27, 36, 44	0
1	B	222/230 (96%)	0.82	17 (7%) 13 15	18, 27, 36, 44	0
1	C	222/230 (96%)	0.84	17 (7%) 13 15	19, 28, 37, 53	0
1	D	220/230 (95%)	0.92	19 (8%) 10 12	19, 28, 37, 48	0
1	E	222/230 (96%)	0.99	25 (11%) 5 5	18, 31, 44, 57	0
1	F	220/230 (95%)	0.87	13 (5%) 22 25	21, 30, 40, 46	0
1	G	222/230 (96%)	0.98	26 (11%) 4 5	20, 30, 39, 48	0
1	H	222/230 (96%)	0.97	23 (10%) 6 7	19, 29, 41, 51	0
1	I	222/230 (96%)	0.77	10 (4%) 33 36	19, 27, 37, 47	0
1	J	222/230 (96%)	0.83	14 (6%) 20 22	19, 28, 36, 42	0
1	K	220/230 (95%)	0.96	17 (7%) 13 15	21, 30, 41, 46	0
1	L	220/230 (95%)	0.98	26 (11%) 4 5	21, 31, 40, 48	0
1	M	222/230 (96%)	1.05	26 (11%) 4 5	21, 34, 46, 55	0
1	N	222/230 (96%)	1.09	26 (11%) 4 5	21, 34, 49, 62	0
1	O	222/230 (96%)	1.13	31 (13%) 2 2	20, 32, 48, 61	0
1	P	220/230 (95%)	1.16	34 (15%) 2 2	21, 33, 48, 62	0
All	All	3542/3680 (96%)	0.95	336 (9%) 8 9	18, 29, 43, 62	0

The worst 5 of 336 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	N	223	PRO	5.0
1	K	167	VAL	4.7
1	M	62	ALA	4.4
1	B	62	ALA	4.2
1	O	50	ILE	4.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.