



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 07:53 pm GMT

PDB ID : 2BNU  
Title : Structural and kinetic basis for heightened immunogenicity of T cell vaccines  
Authors : Chen, J.-L.; Stewart-Jones, G.; Bossi, G.; Lissin, N.M.; Wooldridge, L.; Choi, E.M.L.; Held, G.; Dunbar, P.R.; Esnouf, R.M.; Sami, M.; Boultier, J.M.; Rizkallah, P.J.; Renner, C.; Sewell, A.; Van Der Merwe, P.A.; Jackobsen, B.K.; Griffiths, G.; Jones, E.Y.; Cerundolo, V.  
Deposited on : 2005-04-04  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	1541 (1.40-1.40)
Ramachandran outliers	120053	1500 (1.40-1.40)
Sidechain outliers	120020	1499 (1.40-1.40)
RSRZ outliers	108989	1412 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	203	
2	B	241	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-CELL RECEPTOR ALPHA CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	203	1560	970	265	318	7	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	115	TYR	ASN	conflict	UNP P01848
A	128	ARG	ASP	conflict	UNP P01848
A	162	CYS	THR	conflict	UNP P01848

- Molecule 2 is a protein called T-CELL RECEPTOR BETA CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	241	1902	1196	327	370	9	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	116	LYS	ASN	conflict	UNP P01848
B	117	ASN	LYS	conflict	UNP P01848
B	149	TYR	PHE	conflict	UNP P01848
B	169	CYS	SER	conflict	UNP P01848
B	187	ALA	CYS	conflict	UNP P01848
B	201	ASP	ASN	conflict	UNP P01848

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	104	104	104	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	139	Total 139	O 139	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-CELL RECEPTOR ALPHA CHAIN C REGION



- Molecule 2: T-CELL RECEPTOR BETA CHAIN C REGION



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.99Å 60.30Å 82.64Å 90.00° 89.54° 90.00°	Depositor
Resolution (Å)	20.00 – 1.40 19.70 – 1.39	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.40) 89.7 (19.70-1.39)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.75 (at 1.39Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.245 , (Not available) 0.247 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtrriage
Anisotropy	0.078	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3705	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/1592	0.63	1/2162 (0.0%)
2	B	0.31	0/1953	0.62	1/2659 (0.0%)
All	All	0.31	0/3545	0.62	2/4821 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	CYS	CA-CB-SG	5.12	123.22	114.00
2	B	122	GLU	N-CA-C	-5.01	97.47	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1560	0	1498	69	0
2	B	1902	0	1796	49	0
3	A	104	0	0	0	0
3	B	139	0	0	2	0
All	All	3705	0	3294	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:ASN:HD22	1:A:72:ARG:HH22	1.03	1.02
1:A:130:LYS:HA	1:A:130:LYS:HE3	1.44	1.00
2:B:114:ASP:OD1	2:B:116:LYS:HG2	1.77	0.85
1:A:152:LYS:HD2	1:A:193:ASN:HD21	1.44	0.82
1:A:127:ARG:H	1:A:127:ARG:HD3	1.48	0.78
2:B:95:VAL:HG13	2:B:96:GLY:H	1.50	0.74
1:A:99:SER:HB3	1:A:101:ILE:HG12	1.70	0.73
1:A:22:ASN:HD22	1:A:72:ARG:NH2	1.83	0.71
1:A:22:ASN:ND2	1:A:72:ARG:HH22	1.84	0.71
1:A:18:ASN:HD22	1:A:79:ALA:H	1.40	0.70
2:B:116:LYS:HD2	2:B:223:GLN:CG	2.21	0.70
1:A:101:ILE:HG23	2:B:44:LEU:HD22	1.72	0.70
1:A:13:VAL:HG21	1:A:19:LEU:HD21	1.72	0.69
1:A:129:SER:O	1:A:130:LYS:HB2	1.93	0.66
1:A:22:ASN:HB3	1:A:72:ARG:NH2	2.11	0.66
1:A:18:ASN:ND2	1:A:79:ALA:H	1.95	0.65
1:A:128:ARG:HG2	1:A:130:LYS:N	2.11	0.65
1:A:101:ILE:HD12	2:B:44:LEU:HD21	1.81	0.63
2:B:160:ASN:HD21	2:B:204:ASN:HD22	1.45	0.63
1:A:128:ARG:HG2	1:A:130:LYS:H	1.64	0.61
1:A:150:GLN:H	1:A:150:GLN:CD	2.04	0.60
2:B:46:HIS:ND1	2:B:56:GLN:HA	2.17	0.60
1:A:49:LEU:HD22	2:B:100:GLU:HB2	1.84	0.59
2:B:116:LYS:HE3	2:B:223:GLN:HE21	1.68	0.59
2:B:95:VAL:HG13	2:B:96:GLY:N	2.18	0.58
1:A:195:ILE:O	1:A:195:ILE:HG13	2.02	0.58
1:A:63:ASN:OD1	1:A:76:TYR:HB2	2.03	0.58
1:A:160:ASP:OD1	2:B:172:PRO:HD2	2.04	0.58
1:A:129:SER:OG	2:B:124:ALA:HB1	2.03	0.58
1:A:101:ILE:HD12	2:B:44:LEU:CD2	2.33	0.58
1:A:99:SER:CB	1:A:101:ILE:HG12	2.34	0.57
1:A:13:VAL:HG21	1:A:19:LEU:CD2	2.35	0.57
2:B:45:ILE:HG12	2:B:59:VAL:HG23	1.87	0.57
1:A:127:ARG:N	1:A:127:ARG:HD3	2.18	0.56
2:B:53:ILE:N	2:B:53:ILE:HD12	2.21	0.56
1:A:25:PHE:CE1	1:A:71:GLY:HA2	2.40	0.56
1:A:130:LYS:HA	1:A:130:LYS:CE	2.26	0.56
2:B:69:THR:HG22	2:B:70:THR:N	2.21	0.55
1:A:101:ILE:HD11	2:B:47:TYR:CE1	2.41	0.55
2:B:54:THR:HG21	2:B:65:VAL:HG23	1.89	0.54
2:B:24:GLN:NE2	2:B:26:MET:H	2.05	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:SER:H	1:A:194:SER:HB3	1.72	0.54
1:A:99:SER:C	1:A:101:ILE:H	2.11	0.53
1:A:93:ARG:HG2	1:A:102:PRO:HG3	1.91	0.53
1:A:58:THR:HG22	1:A:59:SER:N	2.24	0.52
2:B:20:LEU:HD12	2:B:75:LEU:HD23	1.90	0.52
1:A:198:GLU:N	1:A:198:GLU:OE1	2.42	0.52
1:A:152:LYS:HD3	1:A:152:LYS:H	1.74	0.52
1:A:114:PRO:HG3	1:A:163:VAL:HG11	1.92	0.51
2:B:35:ARG:HB3	2:B:45:ILE:HD11	1.93	0.51
1:A:130:LYS:O	1:A:131:SER:HB3	2.11	0.51
2:B:116:LYS:HD2	2:B:223:GLN:HG2	1.93	0.50
1:A:128:ARG:HG2	1:A:129:SER:N	2.27	0.50
1:A:153:ASP:OD2	1:A:156:VAL:HG23	2.12	0.50
2:B:119:PHE:CE2	2:B:225:ARG:NH1	2.80	0.50
1:A:106:ARG:HH11	1:A:106:ARG:HG2	1.77	0.49
2:B:160:ASN:HD21	2:B:204:ASN:ND2	2.06	0.49
2:B:227:LYS:HE3	2:B:229:VAL:CG1	2.43	0.49
1:A:129:SER:O	1:A:130:LYS:CB	2.61	0.49
2:B:19:THR:HG22	2:B:21:GLN:HE21	1.79	0.48
1:A:20:VAL:C	1:A:21:LEU:HD12	2.35	0.47
1:A:166:MET:HA	3:B:2092:HOH:O	2.15	0.47
1:A:101:ILE:HG23	2:B:44:LEU:CD2	2.42	0.46
1:A:181:ASN:O	1:A:182:LYS:C	2.53	0.46
2:B:85:THR:HG23	2:B:110:THR:HA	1.98	0.46
1:A:192:ASN:ND2	1:A:193:ASN:OD1	2.48	0.46
2:B:221:TRP:HZ2	2:B:225:ARG:HE	1.62	0.46
1:A:192:ASN:HD22	1:A:192:ASN:C	2.19	0.46
2:B:217:GLU:HA	2:B:227:LYS:NZ	2.31	0.46
1:A:101:ILE:HD11	2:B:47:TYR:CZ	2.51	0.45
1:A:163:VAL:HG22	1:A:174:ASN:OD1	2.17	0.45
1:A:157:TYR:O	1:A:178:ALA:HA	2.17	0.44
2:B:10:GLN:HG2	2:B:18:MET:SD	2.57	0.44
1:A:127:ARG:H	1:A:127:ARG:CD	2.26	0.44
2:B:56:GLN:HB2	2:B:60:PRO:CB	2.48	0.44
1:A:160:ASP:CG	2:B:172:PRO:HD2	2.38	0.43
1:A:99:SER:C	1:A:101:ILE:N	2.70	0.43
2:B:18:MET:HG2	2:B:19:THR:N	2.32	0.43
1:A:152:LYS:CD	1:A:152:LYS:H	2.31	0.43
1:A:55:ARG:C	1:A:55:ARG:HD3	2.39	0.43
1:A:71:GLY:O	1:A:72:ARG:HD2	2.17	0.43
1:A:72:ARG:CZ	1:A:72:ARG:HB3	2.49	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:51:ALA:HA	2:B:67:ARG:HG3	1.99	0.43
2:B:56:GLN:HB2	2:B:60:PRO:HB3	2.01	0.42
1:A:114:PRO:HG3	1:A:163:VAL:CG1	2.49	0.42
1:A:48:LEU:HD21	1:A:59:SER:OG	2.18	0.42
2:B:19:THR:CG2	2:B:21:GLN:HE21	2.32	0.42
1:A:33:LEU:HD13	1:A:73:SER:HB2	2.01	0.42
1:A:2:GLN:HG3	1:A:3:GLU:HG3	2.01	0.42
2:B:52:GLY:N	2:B:67:ARG:O	2.45	0.42
1:A:128:ARG:NE	2:B:127:GLU:OE2	2.49	0.42
2:B:221:TRP:NE1	2:B:223:GLN:HB2	2.34	0.41
1:A:127:ARG:HH22	2:B:240:ARG:NH1	2.18	0.41
1:A:191:PHE:C	1:A:193:ASN:H	2.23	0.41
1:A:49:LEU:CD2	2:B:100:GLU:HB2	2.50	0.41
2:B:38:PRO:C	2:B:40:MET:H	2.24	0.41
2:B:91:ALA:HA	2:B:102:PHE:O	2.20	0.41
2:B:74:PRO:HG3	3:B:2013:HOH:O	2.20	0.41
1:A:156:VAL:HG21	1:A:185:PHE:CE2	2.56	0.41
1:A:2:GLN:HA	1:A:2:GLN:HE21	1.86	0.41
1:A:99:SER:C	1:A:100:TYR:HD1	2.24	0.40
2:B:37:ASP:HB2	2:B:40:MET:HB3	2.03	0.40
1:A:3:GLU:O	1:A:25:PHE:HA	2.21	0.40
2:B:45:ILE:HG22	2:B:46:HIS:CD2	2.57	0.40
2:B:69:THR:HG22	2:B:70:THR:H	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	201/203 (99%)	181 (90%)	15 (8%)	5 (2%)	<b>6</b> <b>0</b>
2	B	239/241 (99%)	228 (95%)	10 (4%)	1 (0%)	<b>36</b> <b>12</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	440/444 (99%)	409 (93%)	25 (6%)	6 (1%)	12   1

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	128	ARG
1	A	130	LYS
1	A	182	LYS
1	A	192	ASN
1	A	185	PHE
2	B	95	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	179/179 (100%)	172 (96%)	7 (4%)	35   7
2	B	208/208 (100%)	205 (99%)	3 (1%)	69   42
All	All	387/387 (100%)	377 (97%)	10 (3%)	49   14

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	ARG
1	A	127	ARG
1	A	128	ARG
1	A	130	LYS
1	A	152	LYS
1	A	192	ASN
1	A	198	GLU
2	B	24	GLN
2	B	181	LEU
2	B	191	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	18	ASN
1	A	22	ASN
1	A	38	GLN
1	A	51	GLN
1	A	57	GLN
1	A	192	ASN
1	A	193	ASN
2	B	21	GLN
2	B	24	GLN
2	B	36	GLN
2	B	204	ASN
2	B	211	GLN
2	B	223	GLN
2	B	231	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	203/203 (100%)	1.70	56 (27%) <b>0</b>   <b>0</b>	11, 22, 51, 58	0
2	B	241/241 (100%)	1.01	32 (13%) <b>3</b>   <b>2</b>	10, 20, 43, 48	0
All	All	444/444 (100%)	1.33	88 (19%) <b>1</b>   <b>0</b>	10, 21, 47, 58	0

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	132	SER	17.6
1	A	184	ASP	11.1
1	A	100	TYR	10.8
1	A	129	SER	10.1
1	A	185	PHE	10.1
2	B	224	ASP	9.3
2	B	59	VAL	8.1
1	A	131	SER	8.1
1	A	130	LYS	8.0
2	B	242	ASP	7.7
1	A	183	SER	7.6
1	A	152	LYS	7.3
1	A	101	ILE	6.9
1	A	182	LYS	6.6
1	A	96	SER	6.3
2	B	60	PRO	6.3
1	A	99	SER	6.3
1	A	95	THR	5.9
1	A	128	ARG	5.9
1	A	195	ILE	5.7
2	B	222	THR	5.0
1	A	192	ASN	4.9
1	A	98	GLY	4.9
2	B	40	MET	4.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	133	ASP	4.8
1	A	181	ASN	4.6
1	A	193	ASN	4.4
2	B	220	GLU	4.2
1	A	150	GLN	4.2
2	B	95	VAL	4.1
2	B	218	ASN	4.0
1	A	155	ASP	4.0
2	B	217	GLU	4.0
1	A	40	PRO	3.9
1	A	72	ARG	3.9
1	A	127	ARG	3.8
1	A	55	ARG	3.8
2	B	55	ASP	3.8
1	A	199	ASP	3.7
1	A	97	GLY	3.7
2	B	58	GLU	3.6
1	A	53	SER	3.5
1	A	31	TYR	3.4
1	A	153	ASP	3.4
1	A	135	SER	3.3
1	A	68	LYS	3.1
2	B	53	ILE	3.1
2	B	76	ARG	3.0
2	B	182	ASN	3.0
2	B	78	LEU	3.0
1	A	170	ASP	2.9
2	B	43	ARG	2.9
1	A	136	VAL	2.9
1	A	90	CYS	2.8
2	B	56	GLN	2.8
2	B	241	ALA	2.8
1	A	2	GLN	2.7
2	B	68	SER	2.6
1	A	27	ASP	2.6
1	A	30	ILE	2.6
1	A	52	SER	2.5
2	B	238	TRP	2.5
1	A	198	GLU	2.5
1	A	19	LEU	2.4
1	A	94	PRO	2.4
1	A	67	ASP	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	61	ASN	2.3
2	B	49	VAL	2.3
2	B	69	THR	2.3
1	A	138	LEU	2.3
1	A	147	ASN	2.3
1	A	151	SER	2.2
2	B	4	THR	2.2
1	A	56	GLU	2.2
2	B	88	TYR	2.2
2	B	39	GLY	2.1
1	A	3	GLU	2.1
1	A	134	LYS	2.1
1	A	63	ASN	2.1
2	B	215	LEU	2.1
1	A	160	ASP	2.1
2	B	33	TRP	2.1
2	B	34	TYR	2.1
1	A	70	SER	2.0
2	B	71	GLU	2.0
2	B	116	LYS	2.0
1	A	178	ALA	2.0
1	A	75	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.