



# Full wwPDB X-ray Structure Validation Report ⓘ

May 19, 2018 – 10:57 AM EDT

PDB ID : 6BNC  
Title : Crystal structure of the intrinsic colistin resistance enzyme ICR(Mc) from *Moraxella catarrhalis*, catalytic domain, Thr315Ala mutant di-zinc and PEG complex  
Authors : Stogios, P.J.; Evdokimova, E.; Wawrzak, Z.; Savchenko, A.; Anderson, W.F.; Satchell, K.J.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)  
Deposited on : 2017-11-16  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031021  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031021


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2534 (1.50-1.50)
Clashscore	122126	2727 (1.50-1.50)
Ramachandran outliers	120053	2661 (1.50-1.50)
Sidechain outliers	120020	2659 (1.50-1.50)
RSRZ outliers	108989	2481 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	 5% 91% 6%
1	B	344	 6% 93% 5%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoethanolamine transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	335	2637	1659	446	515	17	0	9	0
1	B	335	2656	1675	449	514	18	0	12	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	315	ALA	THR	engineered mutation	UNP A0A1E9VP98
B	315	ALA	THR	engineered mutation	UNP A0A1E9VP98

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is POLYETHYLENE GLYCOL (N=34) (three-letter code: 15P) (formula: C<sub>69</sub>H<sub>140</sub>O<sub>35</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			19	12	7		
3	A	1	Total	C	O	0	0
			21	14	7		
3	A	1	Total	C	O	0	0
			11	7	4		
3	A	1	Total	C	O	0	0
			5	3	2		
3	A	1	Total	C	O	0	0
			9	6	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			10	6	4		
3	B	1	Total	C	O	0	0
			11	7	4		
3	B	1	Total	C	O	0	0
			9	6	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	520	Total 545	O 545	0	25
5	B	523	Total 548	O 548	0	25


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoethanolamine transferase



- Molecule 1: Phosphoethanolamine transferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.05Å 154.25Å 66.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.82 – 1.50 29.82 – 1.50	Depositor EDS
% Data completeness (in resolution range)	90.7 (29.82-1.50) 98.5 (29.82-1.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.82 (at 1.50Å)	Xtrriage
Refinement program	PHENIX (dev_2733: ???)	Depositor
R, $R_{free}$	0.132 , 0.153 0.140 , 0.160	Depositor DCC
$R_{free}$ test set	6110 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.1	Xtrriage
Anisotropy	0.012	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 58.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6493	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 15P, ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.77	1/2726 (0.0%)	0.88	5/3712 (0.1%)
1	B	0.79	0/2754	0.92	4/3748 (0.1%)
All	All	0.78	1/5480 (0.0%)	0.90	9/7460 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	456	CYS	CB-SG	5.36	1.91	1.82

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	409	ASP	CB-CG-OD2	-7.98	111.12	118.30
1	B	368	ARG	NE-CZ-NH2	-7.79	116.40	120.30
1	A	368	ARG	NE-CZ-NH2	-7.79	116.41	120.30
1	B	409	ASP	CB-CG-OD2	-7.63	111.43	118.30
1	B	368	ARG	NE-CZ-NH1	6.93	123.77	120.30
1	B	361	ASP	CB-CG-OD1	5.85	123.56	118.30
1	A	456	CYS	CA-CB-SG	5.74	124.33	114.00
1	A	368	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	276	ARG	NE-CZ-NH2	-5.17	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2637	0	2538	15	0
1	B	2656	0	2576	9	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	82	0	96	7	0
3	B	20	0	20	2	0
4	B	1	0	0	0	0
5	A	545	0	0	5	0
5	B	548	0	0	2	0
All	All	6493	0	5230	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:571:THR:H	3:B:604:15P:H532	1.47	0.78
1:A:412:LYS:HA	3:A:605:15P:H17	1.76	0.67
1:A:458:ARG:NH2	5:A:703:HOH:O	2.25	0.67
1:B:343[B]:ILE:HG23	1:B:353:ILE:HD13	1.77	0.65
1:B:513[B]:MET:HG3	1:B:522:GLN:HE22	1.61	0.65
1:A:256:LYS:HE3	1:A:563:SER:HB2	1.81	0.61
1:B:454:ALA:HA	5:B:1032:HOH:O	2.05	0.56
1:A:513:MET:HE2	1:A:518:ALA:HA	1.88	0.56
1:A:557:LEU:HD12	1:A:567:THR:HG21	1.89	0.55
1:A:394:PRO:O	3:A:609:15P:H27	2.08	0.53
1:B:571:THR:N	3:B:604:15P:H532	2.22	0.50
1:B:329:GLU:OE1	1:B:330:LYS:NZ	2.46	0.49
1:B:513[B]:MET:HG3	1:B:522:GLN:NE2	2.27	0.49
1:B:307:ASN:HB2	5:B:1104[B]:HOH:O	2.14	0.48
1:A:343[B]:ILE:HG23	1:A:353:ILE:HD13	1.96	0.47
3:A:603:15P:H582	5:A:1070:HOH:O	2.15	0.47
1:A:394:PRO:HG3	3:A:609:15P:H23	1.97	0.46
3:A:603:15P:H541	5:A:1070:HOH:O	2.15	0.46
1:A:343[A]:ILE:HG23	1:A:353:ILE:HD13	1.98	0.45
3:A:605:15P:H27	3:A:605:15P:H15	1.42	0.43
1:A:515:LYS:HE3	5:A:1108:HOH:O	2.18	0.43
1:A:338:TYR:HE2	1:B:429:HIS:CE1	2.38	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:567:THR:OG1	5:A:702:HOH:O	2.22	0.42
1:A:568:ALA:O	1:A:569:ASP:HB3	2.20	0.41
1:A:429:HIS:CG	3:A:603:15P:H602	2.55	0.41
1:A:553:THR:HB	1:A:554:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	342/344 (99%)	333 (97%)	8 (2%)	1 (0%)	43 19
1	B	345/344 (100%)	336 (97%)	8 (2%)	1 (0%)	43 19
All	All	687/688 (100%)	669 (97%)	16 (2%)	2 (0%)	43 19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	360	SER
1	B	360	SER

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/287 (100%)	287 (100%)	1 (0%)	93	86
1	B	291/287 (101%)	288 (99%)	3 (1%)	78	59
All	All	579/574 (101%)	575 (99%)	4 (1%)	90	71

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	458	ARG
1	B	286[A]	GLN
1	B	286[B]	GLN
1	B	561	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 5 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all  $Z$  scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	15P	A	603	-	18,18,103	0.59	0	17,17,102	0.68	0
3	15P	A	604	-	20,20,103	0.61	0	19,19,102	0.69	0
3	15P	A	605	-	10,10,103	0.45	0	9,9,102	1.16	1 (11%)
3	15P	A	606	-	4,4,103	0.56	0	2,3,102	0.38	0
3	15P	A	607	-	8,8,103	0.47	0	7,7,102	0.63	0
3	15P	A	608	-	6,6,103	0.36	0	5,5,102	1.50	1 (20%)
3	15P	A	609	-	9,9,103	0.57	0	8,8,102	0.62	0
3	15P	B	603	-	10,10,103	0.55	0	9,9,102	1.07	1 (11%)
3	15P	B	604	-	8,8,103	0.40	0	7,7,102	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	15P	A	603	-	-	0/16/16/101	0/0/0/0
3	15P	A	604	-	-	0/18/18/101	0/0/0/0
3	15P	A	605	-	-	0/8/8/101	0/0/0/0
3	15P	A	606	-	-	0/2/2/101	0/0/0/0
3	15P	A	607	-	-	0/6/6/101	0/0/0/0
3	15P	A	608	-	-	0/4/4/101	0/0/0/0
3	15P	A	609	-	-	0/7/7/101	0/0/0/0
3	15P	B	603	-	-	0/8/8/101	0/0/0/0
3	15P	B	604	-	-	0/6/6/101	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	605	15P	C5-O2-C4	-2.81	100.98	113.30
3	A	608	15P	C7-O3-C6	-2.77	101.13	113.30
3	B	603	15P	C5-O2-C4	-2.52	102.23	113.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	15P	3	0
3	A	605	15P	2	0
3	A	609	15P	2	0
3	B	604	15P	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/344 (97%)	0.13	17 (5%) 28 31	8, 15, 36, 71	0
1	B	335/344 (97%)	0.17	19 (5%) 24 26	8, 14, 37, 75	0
All	All	670/688 (97%)	0.15	36 (5%) 26 28	8, 14, 37, 75	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	517	PHE	6.8
1	A	578	ASP	5.8
1	B	517	PHE	5.6
1	A	244	THR	5.3
1	B	545	ALA	5.1
1	B	261	THR	4.9
1	B	244	THR	4.6
1	A	455	GLU	4.2
1	B	260	SER	4.2
1	A	261	THR	4.1
1	B	541	ASN	4.0
1	B	578	ASP	3.8
1	B	270	VAL	3.7
1	A	456	CYS	3.7
1	B	455	GLU	3.5
1	A	260	SER	3.3
1	A	568	ALA	3.1
1	B	267	VAL	2.9
1	A	270	VAL	2.7
1	B	268	VAL	2.7
1	B	259	ALA	2.7
1	A	454	ALA	2.6
1	B	540	ALA	2.6
1	B	542	SER	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	267	VAL	2.5
1	B	454	ALA	2.4
1	B	416	ASN	2.4
1	A	329	GLU	2.3
1	A	569	ASP	2.3
1	B	565	GLN	2.3
1	A	567	THR	2.2
1	A	452	GLU	2.2
1	B	421	ILE	2.2
1	A	416	ASN	2.0
1	B	544	THR	2.0
1	A	415	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	15P	A	608	7/104	0.47	0.27	39,40,42,43	7
3	15P	A	609	10/104	0.56	0.26	52,54,54,54	10
3	15P	A	605	11/104	0.56	0.20	35,36,39,40	11
3	15P	B	603	11/104	0.62	0.22	34,37,45,46	11
3	15P	B	604	9/104	0.63	0.22	57,58,58,59	0
3	15P	A	606	5/104	0.65	0.14	43,43,44,44	5
3	15P	A	604	21/104	0.65	0.24	57,60,62,62	0
3	15P	A	607	9/104	0.65	0.18	43,43,45,45	9
3	15P	A	603	19/104	0.73	0.17	48,53,56,56	0
4	CL	B	605	1/1	0.91	0.07	29,29,29,29	1
2	ZN	A	602	1/1	1.00	0.09	11,11,11,11	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	601	1/1	1.00	0.09	9,9,9,9	0
2	ZN	B	601	1/1	1.00	0.08	9,9,9,9	0
2	ZN	B	602	1/1	1.00	0.09	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.