



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 09:35 PM BST

PDB ID : 4BPZ
Title : Crystal structure of lamA_E269S from *Zobellia galactanivorans* in complex with a trisaccharide of 1,3-1,4-beta-D-glucan.
Authors : Labourel, A.; Jam, M.; Jeudy, A.; Czjzek, M.; Michel, G.
Deposited on : 2013-05-29
Resolution : 1.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1168 (1.14-1.10)
Clashscore	141614	1205 (1.14-1.10)
Ramachandran outliers	138981	1168 (1.14-1.10)
Sidechain outliers	138945	1165 (1.14-1.10)
RSRZ outliers	127900	1146 (1.14-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	 2% 80% 14% . .
1	B	256	 % 83% 11% . .
2	C	3	 100%
2	D	3	 33% 67%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ENDO-1,3-BETA-GLUCANASE, FAMILY GH16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	251	2133	1364	345	420	4	0	18	0
1	B	252	2068	1318	337	409	4	0	8	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	128	HIS	-	expression tag	UNP G0L5X4
A	129	HIS	-	expression tag	UNP G0L5X4
A	130	HIS	-	expression tag	UNP G0L5X4
A	131	HIS	-	expression tag	UNP G0L5X4
A	132	HIS	-	expression tag	UNP G0L5X4
A	133	HIS	-	expression tag	UNP G0L5X4
A	134	GLY	-	expression tag	UNP G0L5X4
A	135	SER	-	expression tag	UNP G0L5X4
A	269	SER	GLU	engineered mutation	UNP G0L5X4
B	128	HIS	-	expression tag	UNP G0L5X4
B	129	HIS	-	expression tag	UNP G0L5X4
B	130	HIS	-	expression tag	UNP G0L5X4
B	131	HIS	-	expression tag	UNP G0L5X4
B	132	HIS	-	expression tag	UNP G0L5X4
B	133	HIS	-	expression tag	UNP G0L5X4
B	134	GLY	-	expression tag	UNP G0L5X4
B	135	SER	-	expression tag	UNP G0L5X4
B	269	SER	GLU	engineered mutation	UNP G0L5X4

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-3)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	3	Total	C	O	0	0	0
			34	18	16			
2	D	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	333	Total	O	0	0
			333	333		
4	B	251	Total	O	0	0
			251	251		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.53Å 76.48Å 142.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.34 – 1.13 42.51 – 1.13	Depositor EDS
% Data completeness (in resolution range)	99.1 (71.34-1.13) 99.1 (42.51-1.13)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 1.13Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.142 , 0.175 0.144 , 0.176	Depositor DCC
R_{free} test set	9107 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	13.7	Xtrriage
Anisotropy	0.450	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4855	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, CA, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.29	12/2240 (0.5%)	1.23	17/3044 (0.6%)
1	B	1.27	6/2155 (0.3%)	1.24	16/2932 (0.5%)
All	All	1.28	18/4395 (0.4%)	1.24	33/5976 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	147	GLU	CD-OE2	11.32	1.38	1.25
1	A	271[A]	ASP	CG-OD1	7.55	1.42	1.25
1	A	271[B]	ASP	CG-OD1	7.55	1.42	1.25
1	A	149[A]	GLU	CD-OE2	-7.41	1.17	1.25
1	A	149[B]	GLU	CD-OE2	-7.41	1.17	1.25
1	A	367	GLU	CG-CD	6.67	1.61	1.51
1	B	297	TYR	CG-CD2	6.42	1.47	1.39
1	A	344	PRO	C-O	6.08	1.35	1.23
1	B	269[A]	SER	CB-OG	-5.99	1.34	1.42
1	B	269[B]	SER	CB-OG	-5.99	1.34	1.42
1	A	367	GLU	CD-OE2	5.60	1.31	1.25
1	A	367	GLU	CD-OE1	5.58	1.31	1.25
1	B	290	SER	CA-CB	5.53	1.61	1.52
1	A	259	TYR	CE2-CZ	-5.43	1.31	1.38
1	A	137	PHE	CG-CD2	5.42	1.46	1.38
1	B	336	GLU	CD-OE1	-5.29	1.19	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	TYR	CD1-CE1	5.07	1.47	1.39
1	B	157	TRP	CD2-CE2	5.03	1.47	1.41

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	291	ASP	CB-CG-OD2	-11.95	107.55	118.30
1	A	291	ASP	CB-CG-OD1	9.99	127.29	118.30
1	A	292[A]	LEU	CB-CG-CD1	-8.87	95.93	111.00
1	A	292[B]	LEU	CB-CG-CD1	-8.87	95.93	111.00
1	B	374	PHE	CB-CG-CD2	-7.62	115.47	120.80
1	A	271[A]	ASP	CB-CG-OD1	-7.23	111.79	118.30
1	A	271[B]	ASP	CB-CG-OD1	-7.23	111.79	118.30
1	B	279	ASP	CB-CG-OD2	-7.15	111.86	118.30
1	B	228	ARG	NE-CZ-NH1	-7.05	116.77	120.30
1	B	182	GLU	CG-CD-OE1	6.91	132.12	118.30
1	A	269[A]	SER	N-CA-CB	-6.80	100.30	110.50
1	A	269[B]	SER	N-CA-CB	-6.80	100.30	110.50
1	B	350	TYR	CZ-CE2-CD2	-6.73	113.74	119.80
1	A	380	ARG	NE-CZ-NH2	-6.52	117.04	120.30
1	B	350	TYR	CG-CD2-CE2	6.44	126.45	121.30
1	A	259	TYR	CB-CG-CD1	6.11	124.67	121.00
1	B	159	TYR	CB-CG-CD2	-6.07	117.36	121.00
1	A	213	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	B	364[A]	ASP	CB-CG-OD1	5.87	123.58	118.30
1	B	364[B]	ASP	CB-CG-OD1	5.87	123.58	118.30
1	B	299	ASN	N-CA-CB	-5.83	100.10	110.60
1	B	328	PHE	CB-CG-CD1	-5.81	116.73	120.80
1	A	271[A]	ASP	CB-CG-OD2	5.78	123.50	118.30
1	A	271[B]	ASP	CB-CG-OD2	5.78	123.50	118.30
1	B	182	GLU	OE1-CD-OE2	-5.62	116.56	123.30
1	B	213	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	159	TYR	CB-CG-CD1	5.45	124.27	121.00
1	A	159	TYR	CB-CG-CD2	-5.27	117.84	121.00
1	A	213	ARG	NE-CZ-NH1	5.27	122.93	120.30
1	B	335	TYR	CB-CG-CD2	-5.24	117.86	121.00
1	A	153	ASP	CB-CG-OD1	5.15	122.94	118.30
1	B	178	THR	CA-CB-CG2	-5.05	105.33	112.40
1	A	151	LYS	CD-CE-NZ	5.04	123.30	111.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	175	GLN	Peptide
1	A	271[A]	ASP	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2133	0	2001	37	1
1	B	2068	0	1924	25	1
2	C	34	0	30	0	0
2	D	34	0	30	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	333	0	0	14	1
4	B	251	0	0	3	0
All	All	4855	0	3985	63	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294[A]:SER:HB3	4:A:2244:HOH:O	1.32	1.26
1:A:360[B]:THR:HG22	4:A:2070:HOH:O	1.41	1.21
1:A:286[B]:HIS:CE1	1:A:299:ASN:OD1	2.17	0.98
1:A:286[A]:HIS:CE1	1:A:299:ASN:OD1	2.17	0.98
1:A:286[B]:HIS:HE1	1:A:299:ASN:OD1	1.46	0.98
1:A:286[A]:HIS:HE1	1:A:299:ASN:OD1	1.47	0.97
1:A:296:GLU:O	4:A:2241:HOH:O	1.88	0.90
1:A:294[B]:SER:HB3	4:A:2244:HOH:O	1.81	0.80
1:A:155:GLU:HG2	4:A:2038:HOH:O	1.81	0.79
1:B:276:ASN:HD21	1:B:279:ASP:H	1.29	0.79
1:A:286[C]:HIS:CE1	1:A:299:ASN:OD1	2.37	0.77
1:B:276:ASN:ND2	1:B:279:ASP:H	1.85	0.74
1:A:253:ASN:HD21	1:A:256:GLY:H	1.41	0.67
1:A:286[C]:HIS:HD2	4:A:2236:HOH:O	1.76	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:HIS:N	4:A:2002:HOH:O	2.30	0.65
1:A:173:GLU:HB3	1:A:175:GLN:HE22	1.62	0.64
1:B:173:GLU:HB3	1:B:175:GLN:HE22	1.63	0.64
1:B:253:ASN:HD21	1:B:256:GLY:H	1.45	0.62
1:A:294[A]:SER:CB	4:A:2244:HOH:O	2.13	0.62
1:B:175:GLN:NE2	1:B:357:MET:H	1.99	0.60
1:A:360[B]:THR:CG2	4:A:2070:HOH:O	2.20	0.57
1:B:179:ASN:C	1:B:179:ASN:HD22	2.06	0.56
1:A:360[A]:THR:HG21	4:A:2067:HOH:O	2.06	0.56
1:B:288:HIS:HD2	1:B:297:TYR:OH	1.89	0.55
1:A:175:GLN:NE2	1:A:357:MET:H	2.04	0.55
1:B:300[A]:LEU:HD11	1:B:337:LEU:HD13	1.88	0.55
1:B:276:ASN:C	1:B:276:ASN:HD22	2.09	0.54
1:B:263[A]:GLU:HG3	2:D:3:BGC:O3	2.07	0.54
1:B:175:GLN:HE21	1:B:357:MET:H	1.57	0.53
1:B:286:HIS:HD2	4:B:2173:HOH:O	1.92	0.52
1:A:253:ASN:ND2	1:A:256:GLY:H	2.08	0.52
1:A:286[C]:HIS:HE1	1:A:299:ASN:OD1	1.81	0.52
1:A:294[B]:SER:OG	1:A:295:ASP:N	2.44	0.51
1:B:253:ASN:ND2	1:B:256:GLY:H	2.09	0.51
1:A:339:ASN:HD22	1:A:340:SER:H	1.56	0.51
1:A:286[B]:HIS:HD2	4:A:2236:HOH:O	1.94	0.51
1:A:286[A]:HIS:HD2	4:A:2236:HOH:O	1.95	0.50
1:A:175:GLN:HE21	1:A:357:MET:H	1.59	0.50
1:B:339:ASN:HD22	1:B:340:SER:H	1.58	0.50
1:B:286:HIS:CE1	1:B:299:ASN:OD1	2.65	0.49
1:A:243:THR:OG1	1:A:349:HIS:HD2	1.96	0.49
1:A:276:ASN:HD22	1:A:276:ASN:C	2.15	0.49
1:A:269[B]:SER:OG	4:A:2217:HOH:O	1.65	0.48
1:B:286:HIS:HE1	1:B:299:ASN:OD1	1.96	0.48
1:B:175:GLN:NE2	1:B:175:GLN:H	2.13	0.47
1:A:253:ASN:C	1:A:253:ASN:HD22	2.19	0.45
4:B:2161:HOH:O	2:D:1:GLC:H2	2.15	0.45
1:B:243:THR:OG1	1:B:349:HIS:HD2	1.99	0.45
1:A:238:TRP:CE2	1:A:361:LEU:HB2	2.52	0.44
1:A:253:ASN:C	1:A:253:ASN:ND2	2.71	0.43
1:A:149[B]:GLU:HG3	1:A:187:SER:HA	2.01	0.43
1:B:321:ASN:OD1	1:B:323[A]:SER:HB2	2.19	0.42
1:A:286[C]:HIS:CD2	4:A:2236:HOH:O	2.61	0.42
1:B:340:SER:OG	1:B:342[B]:ASN:ND2	2.52	0.42
1:B:179:ASN:C	1:B:179:ASN:ND2	2.73	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:288:HIS:HE1	4:B:2158:HOH:O	2.02	0.41
1:A:288:HIS:HD2	1:A:297:TYR:OH	2.03	0.41
1:B:182:GLU:O	1:B:194:ARG:HD2	2.20	0.41
1:B:253:ASN:HD22	1:B:253:ASN:C	2.24	0.41
1:A:230:LYS:HE3	1:A:313:PHE:CE2	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:323[B]:SER:OG	4:A:2317:HOH:O[3_544]	2.01	0.19
1:B:166:ASN:OD1	1:B:307:THR:OG1[1_655]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/256 (105%)	259 (97%)	9 (3%)	0	100	100
1	B	258/256 (101%)	248 (96%)	10 (4%)	0	100	100
All	All	526/512 (103%)	507 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	235/221 (106%)	220 (94%)	15 (6%)	17 1
1	B	225/221 (102%)	217 (96%)	8 (4%)	35 4
All	All	460/442 (104%)	437 (95%)	23 (5%)	30 2

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	175	GLN
1	A	178	THR
1	A	179	ASN
1	A	253	ASN
1	A	276	ASN
1	A	282[A]	SER
1	A	282[B]	SER
1	A	286[A]	HIS
1	A	286[B]	HIS
1	A	286[C]	HIS
1	A	294[A]	SER
1	A	294[B]	SER
1	A	339	ASN
1	A	342[A]	ASN
1	A	342[B]	ASN
1	B	175	GLN
1	B	179	ASN
1	B	253	ASN
1	B	276	ASN
1	B	286	HIS
1	B	339	ASN
1	B	367	GLU
1	B	374	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	175	GLN
1	A	179	ASN
1	A	215	ASN
1	A	253	ASN
1	A	276	ASN
1	A	288	HIS
1	A	339	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	349	HIS
1	A	383	GLN
1	B	175	GLN
1	B	179	ASN
1	B	215	ASN
1	B	253	ASN
1	B	276	ASN
1	B	286	HIS
1	B	288	HIS
1	B	298	GLN
1	B	339	ASN
1	B	349	HIS
1	B	383	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	C	1	2	12,12,12	1.93	3 (25%)	17,17,17	1.32	2 (11%)
2	BGC	C	2	2	11,11,12	1.24	1 (9%)	15,15,17	1.10	1 (6%)
2	BGC	C	3	2	11,11,12	0.76	0	15,15,17	1.42	1 (6%)
2	GLC	D	1	2	12,12,12	1.17	1 (8%)	17,17,17	1.51	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	D	2	2	11,11,12	1.48	2 (18%)	15,15,17	0.90	0
2	BGC	D	3	2	11,11,12	1.56	2 (18%)	15,15,17	1.65	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	BGC	C	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	BGC	D	2	2	-	0/2/19/22	0/1/1/1
2	BGC	D	3	2	-	0/2/19/22	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	GLC	O5-C1	-5.14	1.30	1.42
2	D	3	BGC	O5-C1	-4.15	1.37	1.43
2	D	2	BGC	C2-C3	-3.11	1.47	1.52
2	C	1	GLC	O3-C3	2.75	1.49	1.43
2	D	2	BGC	C4-C3	2.27	1.58	1.52
2	C	2	BGC	C4-C3	2.19	1.57	1.52
2	D	1	GLC	C1-C2	2.12	1.57	1.52
2	C	1	GLC	C4-C3	-2.07	1.47	1.52
2	D	3	BGC	O4-C4	-2.02	1.38	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	BGC	O5-C5-C6	4.10	113.63	107.20
2	C	1	GLC	C3-C4-C5	-3.15	104.61	110.24
2	D	3	BGC	O5-C1-C2	3.11	115.57	110.77
2	D	3	BGC	O2-C2-C1	2.65	114.58	109.15
2	D	1	GLC	C1-O5-C5	2.61	118.58	113.66
2	D	1	GLC	O2-C2-C1	-2.53	103.30	109.16
2	D	1	GLC	C3-C4-C5	-2.52	105.74	110.24
2	C	1	GLC	C1-O5-C5	2.37	118.14	113.66
2	D	3	BGC	O2-C2-C3	2.33	114.81	110.14
2	D	1	GLC	O1-C1-O5	2.27	117.20	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	GLC	O5-C1-C2	-2.25	106.27	110.28
2	C	2	BGC	O4-C4-C5	-2.13	104.01	109.30
2	D	3	BGC	C1-O5-C5	-2.09	109.36	112.19
2	D	1	GLC	C4-C3-C2	-2.03	107.28	110.82

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	GLC	1	0
2	D	3	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/256 (98%)	-0.29	5 (1%) 65 62	10, 15, 30, 60	1 (0%)
1	B	252/256 (98%)	-0.16	2 (0%) 86 84	13, 19, 32, 48	1 (0%)
All	All	503/512 (98%)	-0.22	7 (1%) 75 72	10, 17, 31, 60	2 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	132	HIS	3.6
1	A	134	GLY	3.4
1	A	295	ASP	3.3
1	A	135	SER	2.8
1	B	135	SER	2.3
1	A	294[A]	SER	2.2
1	A	136	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	1	12/12	0.95	0.09	13,15,20,29	0
2	BGC	D	3	11/12	0.96	0.12	23,26,35,36	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BGC	C	3	11/12	0.97	0.09	16,20,27,29	0
2	BGC	D	2	11/12	0.97	0.06	17,18,20,20	0
2	GLC	C	1	12/12	0.97	0.06	12,14,21,23	0
2	BGC	C	2	11/12	0.98	0.04	14,15,17,18	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	B	400	1/1	0.99	0.06	23,23,23,23	1
3	CA	A	400	1/1	1.00	0.04	16,16,16,16	1

6.5 Other polymers [i](#)

There are no such residues in this entry.