



Full wwPDB NMR Structure Validation Report ⓘ

Feb 10, 2019 – 05:58 AM EST

PDB ID : 6BP9
Title : HSPB5 alpha-crystallin domain mutant R120G-ACD
Authors : Rajagopal, P.; Klevit, R.E.
Deposited on : 2017-11-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20031633
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031633

2 Ensemble composition and analysis

This entry contains 8 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:66-A:152, B:64-B:152 (176)	1.33	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 7, 8
2	4, 6

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2846 atoms, of which 1424 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Alpha-crystallin B chain.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	89	1423	444	712	128	138	1	0
1	B	89	1423	444	712	128	138	1	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	120	GLY	ARG	engineered mutation	UNP P02511
B	120	GLY	ARG	engineered mutation	UNP P02511

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Alpha-crystallin B chain



- Molecule 1: Alpha-crystallin B chain




There are no outlier residues in this chain.

4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Alpha-crystallin B chain



- Molecule 1: Alpha-crystallin B chain



4.2.2 Score per residue for model 2


- Molecule 1: Alpha-crystallin B chain

Chain A:  97%



- Molecule 1: Alpha-crystallin B chain

Chain B:  99%



4.2.3 Score per residue for model 3

- Molecule 1: Alpha-crystallin B chain

Chain A:  98%



- Molecule 1: Alpha-crystallin B chain


Chain B:  98%



4.2.4 Score per residue for model 4


- Molecule 1: Alpha-crystallin B chain

Chain A:  96%




- Molecule 1: Alpha-crystallin B chain


Chain B:  98%



4.2.5 Score per residue for model 5


- Molecule 1: Alpha-crystallin B chain

Chain A:  91% 6% ..



- Molecule 1: Alpha-crystallin B chain

Chain B:  93% 6% .



4.2.6 Score per residue for model 6

- Molecule 1: Alpha-crystallin B chain

Chain A:  98% .




- Molecule 1: Alpha-crystallin B chain

Chain B:  100%

There are no outlier residues in this chain.

4.2.7 Score per residue for model 7

- Molecule 1: Alpha-crystallin B chain

Chain A:  90% 7% ..



- Molecule 1: Alpha-crystallin B chain


Chain B:  92% 7% .



4.2.8 Score per residue for model 8

- Molecule 1: Alpha-crystallin B chain

Chain A:  96% ..



- Molecule 1: Alpha-crystallin B chain

Chain B:  98% .



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 8 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CSRosetta	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	6bp9_cs.cif
Number of chemical shift lists	1
Total number of shifts	965
Number of shifts mapped to atoms	965
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	43%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	699	696	691	2±2
1	B	711	712	704	2±2
All	All	11280	11264	11160	31

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:108:GLN:O	1:A:109:ASP:CB	0.72	2.38	5	2
1:B:108:GLN:O	1:B:109:ASP:CB	0.71	2.38	5	2
1:A:108:GLN:O	1:A:109:ASP:HB2	0.57	1.99	5	1
1:B:108:GLN:O	1:B:109:ASP:HB2	0.56	1.99	5	1
1:B:65:LEU:O	1:B:66:SER:C	0.51	2.48	4	1
1:B:126:ALA:O	1:B:127:ASP:C	0.50	2.50	7	1
1:A:126:ALA:O	1:A:127:ASP:C	0.49	2.50	7	1
1:A:104:HIS:O	1:A:105:GLU:C	0.48	2.52	1	2
1:A:152:VAL:HG13	1:A:152:VAL:OXT	0.47	2.09	5	1
1:B:152:VAL:HG13	1:B:152:VAL:OXT	0.47	2.09	5	1
1:A:82:LYS:O	1:A:83:HIS:HB2	0.47	2.10	1	3
1:B:82:LYS:O	1:B:83:HIS:HB2	0.47	2.10	5	3
1:B:81:VAL:O	1:B:81:VAL:HG23	0.47	2.10	7	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:104:HIS:O	1:B:105:GLU:C	0.46	2.52	1	2
1:A:81:VAL:HG23	1:A:81:VAL:O	0.46	2.10	7	1
1:A:81:VAL:O	1:A:81:VAL:HG23	0.45	2.12	2	1
1:B:81:VAL:HG23	1:B:81:VAL:O	0.44	2.12	2	1
1:B:129:ASP:O	1:B:132:THR:OG1	0.42	2.34	3	1
1:B:82:LYS:O	1:B:83:HIS:CB	0.41	2.69	1	2
1:A:82:LYS:O	1:A:83:HIS:CB	0.41	2.69	1	2
1:A:96:ASP:OD1	1:A:96:ASP:N	0.40	2.54	4	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	86/89 (97%)	77±2 (89±2%)	9±2 (10±2%)	1±1 (1±1%)	24	70
1	B	87/89 (98%)	78±2 (89±2%)	9±2 (10±2%)	1±1 (1±1%)	24	70
All	All	1384/1424 (97%)	1233 (89%)	139 (10%)	12 (1%)	24	70

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	105	GLU	1
1	A	149	ARG	1
1	A	102	GLY	1
1	A	95	GLY	1
1	A	66	SER	1
1	A	109	ASP	1
1	B	95	GLY	1
1	B	149	ARG	1
1	B	102	GLY	1
1	B	66	SER	1
1	A	105	GLU	1
1	B	109	ASP	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/81 (99%)	80±0 (100±0%)	0±0 (0±0%)	100	100
1	B	81/81 (100%)	81±0 (100±0%)	0±0 (0±0%)	100	100
All	All	1288/1296 (99%)	1288 (100%)	0 (0%)	100	100

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 43% for the well-defined parts and 44% for the entire structure.

7.1 Chemical shift list 1

File name: 6bp9_cs.cif

Chemical shift list name: *assigned_chem_shift_list_0*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	965
Number of shifts mapped to atoms	965
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	89	0.16 ± 0.08	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	81	0.19 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}'$	80	0.43 ± 0.08	None needed (< 0.5 ppm)
^{15}N	84	-1.25 ± 0.21	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 43%, i.e. 969 atoms were assigned a chemical shift out of a possible 2231. 0 out of 35 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	418/864 (48%)	170/344 (49%)	165/352 (47%)	83/168 (49%)
Sidechain	479/1199 (40%)	298/703 (42%)	177/438 (40%)	4/58 (7%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	72/168 (43%)	34/88 (39%)	28/60 (47%)	10/20 (50%)
Overall	969/2231 (43%)	502/1135 (44%)	370/850 (44%)	97/246 (39%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 44%, i.e. 984 atoms were assigned a chemical shift out of a possible 2250. 0 out of 36 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	426/874 (49%)	173/348 (50%)	169/356 (47%)	84/170 (49%)
Sidechain	486/1208 (40%)	302/708 (43%)	180/442 (41%)	4/58 (7%)
Aromatic	72/168 (43%)	34/88 (39%)	28/60 (47%)	10/20 (50%)
Overall	984/2250 (44%)	509/1144 (44%)	377/858 (44%)	98/248 (40%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	68	MET	CG	38.66	38.33 – 25.73	5.3

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

