



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2024 – 05:27 PM EDT

PDB ID : 6BVK
Title : Ras:SOS:Ras in complex with a small molecule activator
Authors : Phan, J.; Abbott, J.; Fesik, S.W.
Deposited on : 2017-12-13
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	 90% 10%
2	B	482	 89% 8% 3%
3	C	167	 86% 11% 11%

2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 8140 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTPase HRas.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	166	1384	855	241	280	8	0	8	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P01112
A	64	ALA	TYR	conflict	UNP P01112

- Molecule 2 is a protein called Son of sevenless homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	469	4081	2606	711	751	13	0	21	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	565	GLY	-	expression tag	UNP Q07889

- Molecule 3 is a protein called GTPase HRas.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	167	1380	857	238	278	7	0	7	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	GLY	-	expression tag	UNP P01112

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter

code: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	32	10	6	13	3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	1	1	0	0

- Molecule 6 is N-{1-[(5-chloro-1H-indol-3-yl)methyl]piperidin-4-yl}-6-methyl-L-tryptophanamide (three-letter code: EAV) (formula: C₂₆H₃₀ClN₅O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
6	B	1	33	26	1	5	1	0	0

- Molecule 7 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
7	B	1	3	1 2	0	0
7	B	1	3	1 2	0	0
7	B	1	3	1 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			3	1	2		
7	B	1	Total	C	O	0	0
			3	1	2		
7	C	1	Total	C	O	0	0
			3	1	2		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	C	1	Total	Na	0	0
			1	1		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	193	Total	O	0	0
			193	193		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	B	762	Total 762	O 762	0	0
10	C	243	Total 243	O 243	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GTPase HRas



- Molecule 2: Son of sevenless homolog 1



- Molecule 3: GTPase HRas



4 Data and refinement statistics i

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	184.42Å 184.42Å 179.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.53 – 1.80 33.53 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (33.53-1.80) 99.9 (33.53-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 1.79Å)	Xtrriage
Refinement program	PHENIX (1.12rc1_2801: ???)	Depositor
R, R_{free}	0.155 , 0.179 0.156 , 0.179	Depositor DCC
R_{free} test set	6880 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtrriage
Anisotropy	0.128	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.005 for l,-k,h 0.008 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8140	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, NA, EAV, FMT, MG, GNP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1394	0.56	0/1877
2	B	0.38	0/4174	0.51	0/5638
3	C	0.37	0/1399	0.57	0/1886
All	All	0.38	0/6967	0.53	0/9401

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	25	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1384	0	1342	11	0
2	B	4081	0	4090	26	0
3	C	1380	0	1345	16	0
4	A	32	0	13	0	0
5	A	1	0	0	0	0
6	B	33	0	0	0	0
7	B	15	0	5	1	0
7	C	3	0	1	1	0
8	B	12	0	16	0	0
9	C	1	0	0	0	0
10	A	193	0	0	4	0
10	B	762	0	0	13	2
10	C	243	0	0	8	1
All	All	8140	0	6812	54	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:68:ARG:NH1	10:C:301:HOH:O	2.04	0.88
2:B:954:GLU:OE1	10:B:2101:HOH:O	1.91	0.88
2:B:621:PRO:O	10:B:2102:HOH:O	1.96	0.81
2:B:892[A]:GLN:OE1	10:B:2104:HOH:O	2.01	0.78
1:A:129:GLN:OE1	10:A:301:HOH:O	2.01	0.76
3:C:68:ARG:NH1	10:C:303:HOH:O	2.18	0.75
2:B:988[B]:LYS:NZ	10:B:2108:HOH:O	2.23	0.72
3:C:70[B]:GLN:NE2	10:C:304:HOH:O	2.25	0.70
7:C:201:FMT:O2	10:C:302:HOH:O	2.12	0.68
3:C:27:HIS:N	10:C:306:HOH:O	2.28	0.65
2:B:680:GLU:OE1	10:B:2106:HOH:O	2.16	0.62
2:B:1041:ARG:NH2	10:B:2112:HOH:O	2.34	0.61
3:C:85:ASN:HD22	3:C:121:ALA:HB3	1.66	0.60
3:C:21:ILE:O	3:C:25:GLN:HB2	2.02	0.59
2:B:1019[B]:ARG:NH1	10:B:2113:HOH:O	2.34	0.58
3:C:117[A]:LYS:NZ	10:C:310:HOH:O	2.41	0.53
2:B:616:HIS:ND1	7:B:2005:FMT:O2	2.39	0.53
2:B:735:LYS:O	2:B:739[A]:ARG:HG3	2.08	0.53
2:B:896[B]:ARG:NH1	10:B:2105:HOH:O	2.12	0.53
1:A:1:MET:HG2	1:A:51:CSO:HA	1.93	0.51
2:B:623:PHE:HE1	2:B:958:PHE:CZ	2.30	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:912:TYR:CG	2:B:932:ILE:HD11	2.46	0.50
2:B:970[B]:GLU:HG3	10:B:2449:HOH:O	2.11	0.49
1:A:49:GLU:HG2	10:A:336:HOH:O	2.11	0.49
1:A:102[B]:ARG:NH2	10:A:302:HOH:O	2.25	0.49
2:B:859[B]:LEU:HD23	2:B:874:VAL:HB	1.95	0.48
2:B:954:GLU:OE2	10:B:2107:HOH:O	2.20	0.47
2:B:828:THR:HG23	2:B:873:GLU:HG2	1.98	0.46
3:C:118:CYS:HB2	10:C:330:HOH:O	2.15	0.46
3:C:122:ALA:HA	10:C:397:HOH:O	2.15	0.45
2:B:645:ILE:O	2:B:649[B]:GLU:HG2	2.17	0.45
3:C:38:ASP:N	3:C:38:ASP:OD1	2.49	0.45
3:C:99:GLN:O	3:C:103[A]:VAL:HG13	2.16	0.45
3:C:24:ILE:HD13	3:C:42:LYS:HB2	1.98	0.45
1:A:61:GLN:HG2	10:A:375:HOH:O	2.17	0.44
2:B:823[B]:LYS:NZ	10:B:2129:HOH:O	2.49	0.44
3:C:27:HIS:CE1	3:C:149:ARG:HD3	2.53	0.44
2:B:587:ILE:HD11	2:B:950:ARG:HG3	2.00	0.43
2:B:810:THR:HG21	2:B:938[B]:LEU:HD12	2.00	0.43
1:A:79:LEU:HG	1:A:114:VAL:HG21	2.00	0.43
2:B:622[B]:ASN:O	2:B:626[B]:THR:HG23	2.18	0.43
1:A:2:THR:O	1:A:51:CSO:HB3	2.17	0.43
3:C:24:ILE:HG21	3:C:39[A]:SER:O	2.19	0.42
1:A:111[B]:MET:HE3	1:A:111[B]:MET:HB3	1.87	0.42
1:A:158:THR:O	1:A:162[A]:GLU:HG3	2.19	0.42
2:B:570:PRO:HB3	2:B:651:PRO:HG2	2.01	0.42
1:A:78:PHE:HB2	1:A:111[B]:MET:HG2	2.02	0.41
2:B:949:LYS:HE2	10:B:2101:HOH:O	2.20	0.41
1:A:148:THR:O	1:A:149:ARG:HB2	2.20	0.41
2:B:895[A]:SER:HB2	10:B:2109:HOH:O	2.19	0.41
3:C:29:VAL:HG12	3:C:31:GLU:HB2	2.02	0.41
2:B:807:SER:HA	2:B:809:TRP:CZ3	2.57	0.40
2:B:859[A]:LEU:HD11	2:B:871:VAL:HG13	2.02	0.40
3:C:24:ILE:HD11	3:C:55:ILE:HD12	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:B:2700:HOH:O	10:B:2718:HOH:O[7_555]	1.90	0.30
10:B:2735:HOH:O	10:C:515:HOH:O[3_555]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/167 (102%)	168 (98%)	3 (2%)	0	100	100
2	B	486/482 (101%)	480 (99%)	6 (1%)	0	100	100
3	C	172/167 (103%)	164 (95%)	7 (4%)	1 (1%)	22	11
All	All	829/816 (102%)	812 (98%)	16 (2%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	26	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/142 (106%)	149 (99%)	1 (1%)	81	79
2	B	460/447 (103%)	457 (99%)	3 (1%)	81	79
3	C	151/144 (105%)	145 (96%)	6 (4%)	27	14
All	All	761/733 (104%)	751 (99%)	10 (1%)	67	59

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	TYR
2	B	753	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	756	SER
2	B	930	PHE
3	C	27	HIS
3	C	31	GLU
3	C	36	ILE
3	C	38	ASP
3	C	127[A]	SER
3	C	127[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	51	1	3,6,7	1.03	0	1,6,8	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	51	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	51	CSO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EAV	B	2001	-	35,37,37	1.20	4 (11%)	42,53,53	1.62	9 (21%)
8	GOL	B	2007	-	5,5,5	0.82	0	5,5,5	0.93	0
4	GNP	A	201	5	29,34,34	1.75	6 (20%)	33,54,54	2.13	7 (21%)
7	FMT	B	2002	-	2,2,2	0.74	0	1,1,1	0.23	0
7	FMT	C	201	-	2,2,2	0.58	0	1,1,1	0.08	0
7	FMT	B	2003	-	2,2,2	0.73	0	1,1,1	0.22	0
7	FMT	B	2006	-	2,2,2	0.72	0	1,1,1	0.23	0
7	FMT	B	2005	-	2,2,2	0.66	0	1,1,1	0.36	0
8	GOL	B	2008	-	5,5,5	0.99	0	5,5,5	1.05	1 (20%)
7	FMT	B	2004	-	2,2,2	0.73	0	1,1,1	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	B	2007	-	-	0/4/4/4	-
4	GNP	A	201	5	-	4/14/38/38	0/3/3/3
6	EAV	B	2001	-	-	0/14/26/26	0/5/5/5
8	GOL	B	2008	-	-	2/4/4/4	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	201	GNP	C6-N1	4.25	1.40	1.33
4	A	201	GNP	PB-O3A	-3.91	1.54	1.59
4	A	201	GNP	PA-O3A	-3.58	1.55	1.59
4	A	201	GNP	PG-O1G	3.24	1.51	1.46
4	A	201	GNP	PB-O2B	-3.02	1.48	1.56
6	B	2001	EAV	C23-N4	2.57	1.41	1.36
6	B	2001	EAV	C8-N1	2.53	1.39	1.34
6	B	2001	EAV	C14-N3	2.41	1.41	1.36
6	B	2001	EAV	C17-C18	2.27	1.42	1.38
4	A	201	GNP	PG-O3G	-2.12	1.51	1.56

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	201	GNP	C5-C6-N1	-8.08	112.62	123.42
6	B	2001	EAV	C1-C25-C24	-5.85	117.07	121.19
4	A	201	GNP	C2-N1-C6	5.78	124.00	115.96
4	A	201	GNP	O2B-PB-O1B	3.31	116.98	109.87
6	B	2001	EAV	C17-C18-C19	2.95	125.24	122.00
6	B	2001	EAV	C17-C16-C15	-2.85	117.39	120.80
4	A	201	GNP	C2-N3-C4	-2.76	112.51	115.48
4	A	201	GNP	N3-C2-N1	-2.55	123.97	127.21
4	A	201	GNP	O1B-PB-N3B	-2.43	108.20	111.77
6	B	2001	EAV	C2-C1-C25	2.40	121.24	118.40
6	B	2001	EAV	C6-C5-C4	-2.39	122.52	126.25
6	B	2001	EAV	C22-C9-C10	-2.35	106.74	110.80
4	A	201	GNP	O1G-PG-N3B	-2.29	108.39	111.77
6	B	2001	EAV	C13-C12-N2	-2.29	110.84	114.07
6	B	2001	EAV	C19-C18-CL	-2.28	116.90	119.65
6	B	2001	EAV	C18-C19-C20	-2.22	116.61	119.66
8	B	2008	GOL	C3-C2-C1	-2.05	104.29	111.80

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	201	GNP	PB-N3B-PG-O1G
4	A	201	GNP	PG-N3B-PB-O1B
8	B	2008	GOL	C1-C2-C3-O3
8	B	2008	GOL	O2-C2-C3-O3
4	A	201	GNP	PA-O3A-PB-O1B
4	A	201	GNP	PG-N3B-PB-O3A


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	201	FMT	1	0
7	B	2005	FMT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/167 (98%)	-0.33	2 (1%) 76 76	9, 19, 42, 55	8 (4%)
2	B	469/482 (97%)	-0.40	15 (3%) 50 48	5, 18, 40, 70	21 (4%)
3	C	167/167 (100%)	0.00	18 (10%) 12 10	8, 19, 52, 84	7 (4%)
All	All	801/816 (98%)	-0.30	35 (4%) 39 37	5, 18, 43, 84	36 (4%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	27	HIS	9.7
3	C	29	VAL	7.9
3	C	26	ASN	6.1
2	B	743	ALA	5.1
2	B	751	ASN	5.0
3	C	30	ASP	4.5
3	C	120	LEU	4.3
3	C	31	GLU	4.2
3	C	38	ASP	3.5
2	B	597	GLY	3.5
3	C	36	ILE	3.4
2	B	755	GLN	3.3
3	C	24	ILE	3.3
2	B	565	GLY	3.2
1	A	165	GLN	3.1
3	C	25	GLN	3.1
3	C	28	PHE	2.9
3	C	0	GLY	2.9
3	C	37	GLU	2.9
2	B	1045	PRO	2.8
2	B	590	GLU	2.8
2	B	573	ASP	2.7
3	C	39[A]	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	723	GLY	2.4
3	C	121	ALA	2.3
3	C	118	CYS	2.2
2	B	1046	ARG	2.2
2	B	752	ILE	2.2
3	C	32	TYR	2.2
2	B	660[A]	ARG	2.1
2	B	574	VAL	2.1
2	B	716	GLU	2.1
1	A	122	ALA	2.0
2	B	724	LYS	2.0
3	C	122	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	A	51	7/8	0.95	0.10	16,17,31,37	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	FMT	B	2006	3/3	0.81	0.18	67,67,69,70	0
6	EAV	B	2001	33/33	0.83	0.18	30,52,71,73	0
7	FMT	C	201	3/3	0.88	0.17	27,27,32,38	0
8	GOL	B	2008	6/6	0.89	0.14	23,43,51,52	0
8	GOL	B	2007	6/6	0.90	0.13	21,37,44,51	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	FMT	B	2003	3/3	0.92	0.12	44,44,48,52	0
7	FMT	B	2002	3/3	0.94	0.15	50,50,52,56	0
7	FMT	B	2004	3/3	0.94	0.14	32,32,33,35	0
7	FMT	B	2005	3/3	0.96	0.14	20,20,21,29	0
9	NA	C	202	1/1	0.98	0.08	33,33,33,33	0
4	GNP	A	201	32/32	0.99	0.03	12,15,18,23	0
5	MG	A	202	1/1	1.00	0.02	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.