



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 02:36 pm BST

PDB ID : 8BWD  
Title : Crystal structure of human Twisted gastrulation protein homolog 1 (TWSG1), crystal form 1  
Authors : Malinauskas, T.; Rudolf, A.F.; Moore, G.; Eggington, H.; Belnoue-Davis, H.; El Omari, K.; Woolley, R.E.; Griffiths, S.C.; Duman, R.; Wagner, A.; Leedham, S.J.; Baldock, C.; Ashe, H.; Siebold, C.  
Deposited on : 2022-12-06  
Resolution : 2.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	 7% 69% 9% 22%
1	B	210	 10% 64% 13% 22%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4812 atoms, of which 2312 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Twisted gastrulation protein homolog 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	164	2403	766	1156	207	243	31	0	0	0
1	B	164	2403	766	1156	207	243	31	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLU	-	expression tag	UNP Q9GZX9
A	24	THR	-	expression tag	UNP Q9GZX9
A	25	GLY	-	expression tag	UNP Q9GZX9
A	224	GLY	-	expression tag	UNP Q9GZX9
A	225	THR	-	expression tag	UNP Q9GZX9
A	226	LYS	-	expression tag	UNP Q9GZX9
A	227	HIS	-	expression tag	UNP Q9GZX9
A	228	HIS	-	expression tag	UNP Q9GZX9
A	229	HIS	-	expression tag	UNP Q9GZX9
A	230	HIS	-	expression tag	UNP Q9GZX9
A	231	HIS	-	expression tag	UNP Q9GZX9
A	232	HIS	-	expression tag	UNP Q9GZX9
B	23	GLU	-	expression tag	UNP Q9GZX9
B	24	THR	-	expression tag	UNP Q9GZX9
B	25	GLY	-	expression tag	UNP Q9GZX9
B	224	GLY	-	expression tag	UNP Q9GZX9
B	225	THR	-	expression tag	UNP Q9GZX9
B	226	LYS	-	expression tag	UNP Q9GZX9
B	227	HIS	-	expression tag	UNP Q9GZX9
B	228	HIS	-	expression tag	UNP Q9GZX9
B	229	HIS	-	expression tag	UNP Q9GZX9
B	230	HIS	-	expression tag	UNP Q9GZX9
B	231	HIS	-	expression tag	UNP Q9GZX9
B	232	HIS	-	expression tag	UNP Q9GZX9

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Twisted gastrulation protein homolog 1



- Molecule 1: Twisted gastrulation protein homolog 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.99Å 97.99Å 187.62Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.43 – 2.63 42.43 – 2.63	Depositor EDS
% Data completeness (in resolution range)	53.3 (42.43-2.63) 53.3 (42.43-2.63)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.73 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.259 , 0.305 0.263 , 0.308	Depositor DCC
$R_{free}$ test set	460 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	82.7	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 46.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	4812	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	111.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1272	0.49	0/1718
1	B	0.27	0/1272	0.50	0/1718
All	All	0.27	0/2544	0.49	0/3436

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1247	1156	1154	13	0
1	B	1247	1156	1154	16	0
2	A	5	0	0	0	0
3	A	1	0	0	0	0
All	All	2500	2312	2308	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:68:ASP:O	1:B:80:ARG:NH2	2.33	0.61
1:B:123:VAL:O	1:B:124:ALA:HB3	2.04	0.58
1:B:174:ASP:OD1	1:B:220:ASN:ND2	2.38	0.57
1:A:126:GLU:OE1	1:A:126:GLU:N	2.33	0.56
1:A:123:VAL:O	1:A:124:ALA:HB3	2.05	0.56
1:A:92:THR:HG21	1:A:178:ILE:HD12	1.87	0.55
1:A:96:LEU:HD12	1:A:190:ALA:O	2.07	0.55
1:B:126:GLU:OE1	1:B:126:GLU:N	2.35	0.52
1:A:84:ASP:OD2	1:B:81:ASN:ND2	2.43	0.52
1:B:92:THR:HG21	1:B:178:ILE:HD12	1.95	0.49
1:A:72:ASP:OD1	1:A:73:CYS:N	2.46	0.48
1:A:171:TYR:HB3	1:A:218:CYS:CB	2.44	0.48
1:B:96:LEU:HD12	1:B:190:ALA:O	2.14	0.48
1:B:171:TYR:HB3	1:B:218:CYS:CB	2.44	0.47
1:B:72:ASP:OD1	1:B:73:CYS:N	2.48	0.46
1:B:41:GLN:O	1:B:42:GLU:HB2	2.16	0.46
1:A:41:GLN:O	1:A:42:GLU:HB2	2.17	0.45
1:B:123:VAL:O	1:B:124:ALA:CB	2.66	0.44
1:A:123:VAL:O	1:A:124:ALA:CB	2.65	0.44
1:A:175:CYS:SG	1:A:199:ALA:HB3	2.59	0.43
1:A:171:TYR:HB3	1:A:218:CYS:HB3	2.01	0.41
1:B:175:CYS:SG	1:B:199:ALA:HB3	2.61	0.41
1:B:46:CYS:SG	1:B:53:CYS:SG	3.19	0.41
1:A:98:GLU:N	1:A:99:PRO:HD3	2.36	0.40
1:A:172:PHE:CZ	1:A:184:SER:HB3	2.57	0.40
1:B:57:LYS:O	1:B:61:LEU:HD23	2.21	0.40
1:B:79:PRO:HB2	1:B:82:TYR:CE2	2.57	0.40
1:B:113:GLN:C	1:B:115:ASN:H	2.24	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/210 (76%)	151 (94%)	9 (6%)	0	100	100
1	B	160/210 (76%)	148 (92%)	12 (8%)	0	100	100
All	All	320/420 (76%)	299 (93%)	21 (7%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/192 (77%)	146 (99%)	1 (1%)	84	91
1	B	147/192 (77%)	145 (99%)	2 (1%)	67	80
All	All	294/384 (77%)	291 (99%)	3 (1%)	76	86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	175	CYS
1	B	114	LEU
1	B	175	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	41	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	301	-	4,4,4	0.14	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	164/210 (78%)	0.83	15 (9%) <b>9</b> <b>7</b>	56, 92, 128, 166	0
1	B	164/210 (78%)	0.91	22 (13%) <b>3</b> <b>2</b>	63, 101, 152, 184	0
All	All	328/420 (78%)	0.87	37 (11%) <b>5</b> <b>3</b>	56, 98, 144, 184	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	114	LEU	6.4
1	A	82	TYR	6.3
1	B	51	GLY	4.8
1	B	48	PRO	4.7
1	B	165	HIS	4.1
1	B	45	GLN	3.8
1	B	126	GLU	3.7
1	B	50	GLU	3.4
1	B	120	SER	3.2
1	A	165	HIS	3.0
1	A	211	TYR	2.9
1	B	170	VAL	2.8
1	B	112	THR	2.8
1	A	121	PHE	2.7
1	B	43	LEU	2.7
1	B	82	TYR	2.7
1	A	217	LYS	2.7
1	B	47	ARG	2.7
1	B	209	ILE	2.7
1	A	63	LEU	2.7
1	B	116	TRP	2.6
1	B	109	GLU	2.5
1	A	52	ASN	2.5
1	B	210	ASP	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	166	MET	2.4
1	B	181	CYS	2.3
1	B	39	LEU	2.2
1	A	45	GLN	2.2
1	A	48	PRO	2.2
1	A	51	GLY	2.2
1	B	123	VAL	2.2
1	A	25	GLY	2.1
1	B	61	LEU	2.0
1	B	124	ALA	2.0
1	A	66	LEU	2.0
1	A	210	ASP	2.0
1	A	107	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	301	5/5	0.88	0.23	118,124,141,150	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.