

# Full wwPDB EM Validation Report (i)

Nov 6, 2023 – 10:07 pm GMT

PDB ID : 8BWR

EMDB ID : EMD-16295

Title: Cryo-EM structure of nanodisc-reconstituted wildtype human MRP4 (in

complex with prostaglandin E2)

Authors : Raj, I.; Bloch, M.; Pape, T.H.; Taylor, N.M.I.

Deposited on : 2022-12-07

Resolution : 4.00 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : FAILED buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : FAILED

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 4.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.



## 2 Entry composition (i)


There are 2 unique types of molecules in this entry. The entry contains 9570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP-binding cassette sub-family C member 4.

Mol	Chain	Residues	Atoms				AltConf	Trace	
1	A	1195	Total 9545	C 6201	N 1602	O 1699	S 43	0	0

• Molecule 2 is (Z)-7-[(1R,2R,3R)-3-hydroxy-2-[(E,3S)-3-hydroxyoct-1-enyl]-5-oxo-cyclopen tyl]hept-5-enoic acid (three-letter code: P2E) (formula:  $C_{20}H_{32}O_5$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	AltConf
2	A	1	Total C O 25 20 5	0

MolProbity failed to run properly - this section is therefore empty.



# 3 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	107793	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	40	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	96000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor



### 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	Chain	Res Link		Во	ond leng	$ ag{ths}$	В	ond ang	les
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	P2E	A	1401	-	25,25,25	1.43	6 (24%)	26,31,31	1.06	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P2E	A	1401	-	-	10/19/35/35	0/1/1/1

#### All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
2	A	1401	P2E	C7-C8	-4.08	1.48	1.54
2	A	1401	P2E	O5-C15	-2.58	1.38	1.43
2	A	1401	P2E	C10-C9	-2.43	1.47	1.51
2	A	1401	P2E	O4-C11	-2.27	1.38	1.43
2	A	1401	P2E	C8-C9	-2.22	1.48	1.52
2	A	1401	P2E	C12-C8	-2.12	1.51	1.55

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^{o})$	$  \ \mathbf{Ideal}(^o)  $
2	A	1401	P2E	C8-C7-C6	2.83	118.85	112.94

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1401	P2E	C14-C15-C16-C17
2	A	1401	P2E	O5-C15-C16-C17
2	A	1401	P2E	C13-C14-C15-C16
2	A	1401	P2E	C13-C14-C15-O5
2	A	1401	P2E	O2-C1-C2-C3
2	A	1401	P2E	C15-C16-C17-C18
2	A	1401	P2E	O1-C1-C2-C3
2	A	1401	P2E	C17-C18-C19-C20

Continued on next page..



Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1401	P2E	C8-C12-C13-C14
2	A	1401	P2E	C11-C12-C13-C14

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 5 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-16295. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 5.1 Orthogonal projections (i)

This section was not generated.

#### 5.2 Central slices (i)

This section was not generated.

### 5.3 Largest variance slices (i)

This section was not generated.

### 5.4 Orthogonal standard-deviation projections (False-color) (i)

This section was not generated.

### 5.5 Orthogonal surface views (i)

This section was not generated.

### 5.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



## 6 Map analysis (i)

This section contains the results of statistical analysis of the map.

### 6.1 Map-value distribution (i)

This section was not generated.

### 6.2 Volume estimate versus contour level (i)

This section was not generated.

### 6.3 Rotationally averaged power spectrum (i)

This section was not generated. The rotationally averaged power spectrum had issues being displayed.



# 7 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



# 8 Map-model fit (i)

This section was not generated.

