



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 16, 2024 – 07:58 AM EST

PDB ID : 6BYU
Title : X-ray crystal structure of Escherichia coli RNA polymerase (RpoB-H526Y) and ppApp complex
Authors : Murakami, K.S.; Molodtsov, V.
Deposited on : 2017-12-21
Resolution : 3.60 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1563 (3.70-3.50)
Clashscore	180529	1665 (3.70-3.50)
Ramachandran outliers	177936	1641 (3.70-3.50)
Sidechain outliers	177891	1640 (3.70-3.50)
RSRZ outliers	164620	1562 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	
1	B	329	
1	G	329	
1	H	329	
2	C	1342	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	1342	 64% 32% .
3	D	1407	 52% 26% 5% 17%
3	J	1407	 52% 26% . 18%
4	E	91	 66% 30% . .
4	K	91	 % 47% 38% . 13%
5	F	613	 51% 23% . 24%
5	L	613	 57% 17% . 23%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 54996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	228	Total 1768	C 1102	N 312	O 348	S 6	0	0	0
1	B	217	Total 1672	C 1044	N 295	O 327	S 6	0	0	0
1	G	224	Total 1730	C 1076	N 308	O 340	S 6	0	0	0
1	H	217	Total 1667	C 1041	N 293	O 327	S 6	0	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	1340	Total 10561	C 6626	N 1837	O 2055	S 43	0	0	0
2	I	1340	Total 10557	C 6624	N 1836	O 2054	S 43	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	526	TYR	HIS	engineered mutation	UNP P0A8V2
I	526	TYR	HIS	engineered mutation	UNP P0A8V2

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	1163	Total 9063	C 5696	N 1622	O 1699	S 46	0	0	0
3	J	1155	Total 8998	C 5656	N 1612	O 1684	S 46	0	0	0


- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	89	Total	C	N	O	S	0	0	0
			685	418	126	140	1			
4	K	79	Total	C	N	O	S	0	0	0
			627	382	118	126	1			

- Molecule 5 is a protein called RNA polymerase sigma factor RpoD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	467	Total	C	N	O	S	0	0	0
			3796	2379	677	717	23			
5	L	469	Total	C	N	O	S	0	0	0
			3796	2379	677	717	23			

- Molecule 6 is (5R)-5-(6-amino-9H-purin-9-yl)-2-({[(S)-hydroxy(phosphonooxy)phosphoryl]oxy}methyl)-4-oxo-4,5-dihydrofuran-3-yl trihydrogen diphosphate (three-letter code: ECJ) (formula: C₁₀H₁₃N₅O₁₆P₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	C	1	Total	C	N	O	P	0	0
			35	10	5	16	4		
6	J	1	Total	C	N	O	P	0	0
			35	10	5	16	4		

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total Mg 1 1	0	0
7	J	1	Total Mg 1 1	0	0

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	2	Total Zn 2 2	0	0
8	J	2	Total Zn 2 2	0	0



- Molecule 2: DNA-directed RNA polymerase subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	186.25Å 203.66Å 308.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.60 30.00 – 3.60	Depositor EDS
% Data completeness (in resolution range)	98.5 (30.00-3.60) 98.3 (30.00-3.60)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 3.56Å)	Xtrriage
Refinement program	PHENIX (1.12_2829)	Depositor
R, R_{free}	0.244 , 0.311 0.245 , 0.308	Depositor DCC
R_{free} test set	133657 reflections (1.48%)	wwPDB-VP
Wilson B-factor (Å ²)	119.5	Xtrriage
Anisotropy	0.210	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 121.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	54996	wwPDB-VP
Average B, all atoms (Å ²)	153.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ECJ, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/1790	0.57	0/2426
1	B	0.27	0/1692	0.54	1/2293 (0.0%)
1	G	0.27	0/1751	0.56	0/2373
1	H	0.27	0/1686	0.55	1/2285 (0.0%)
2	C	0.26	0/10730	0.50	0/14479
2	I	0.26	0/10726	0.49	0/14474
3	D	0.26	0/9201	0.49	0/12420
3	J	0.26	0/9137	0.49	0/12337
4	E	0.26	0/687	0.48	0/928
4	K	0.23	0/629	0.47	0/847
5	F	0.25	0/3847	0.45	0/5171
5	L	0.25	0/3846	0.44	0/5171
All	All	0.26	0/55722	0.50	2/75204 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	I	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	13	LEU	CA-CB-CG	5.80	128.64	115.30
1	B	13	LEU	CA-CB-CG	5.48	127.90	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	1159	VAL	Peptide
2	I	1159	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1768	0	1793	74	0
1	B	1672	0	1694	64	0
1	G	1730	0	1756	73	0
1	H	1667	0	1689	60	0
2	C	10561	0	10555	305	0
2	I	10557	0	10549	301	0
3	D	9063	0	9234	291	0
3	J	8998	0	9154	287	0
4	E	685	0	684	18	0
4	K	627	0	634	19	0
5	F	3796	0	3858	96	1
5	L	3796	0	3848	71	1
6	C	35	0	0	0	0
6	J	35	0	0	1	0
7	D	1	0	0	0	0
7	J	1	0	0	0	0
8	D	2	0	0	0	0
8	J	2	0	0	0	0
All	All	54996	0	55448	1510	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 1510 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:45:ARG:HG2	1:H:38:THR:HB	1.49	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:696:ASP:HB2	2:I:798:GLN:HG2	1.51	0.92
2:C:696:ASP:HB2	2:C:798:GLN:HG2	1.53	0.91
2:I:10:ARG:HD3	2:I:1181:PRO:HG2	1.54	0.88
3:D:418:GLU:HG3	4:E:45:LYS:H	1.42	0.85

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:219:GLU:OE1	5:L:232:ARG:NH2[1_565]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/329 (69%)	202 (89%)	16 (7%)	8 (4%)	3	24
1	B	213/329 (65%)	195 (92%)	16 (8%)	2 (1%)	14	48
1	G	222/329 (68%)	196 (88%)	20 (9%)	6 (3%)	4	28
1	H	213/329 (65%)	194 (91%)	17 (8%)	2 (1%)	14	48
2	C	1338/1342 (100%)	1203 (90%)	114 (8%)	21 (2%)	8	38
2	I	1338/1342 (100%)	1203 (90%)	112 (8%)	23 (2%)	7	37
3	D	1157/1407 (82%)	1038 (90%)	101 (9%)	18 (2%)	8	38
3	J	1151/1407 (82%)	1036 (90%)	100 (9%)	15 (1%)	10	41
4	E	87/91 (96%)	81 (93%)	4 (5%)	2 (2%)	5	31
4	K	77/91 (85%)	72 (94%)	4 (5%)	1 (1%)	10	41
5	F	461/613 (75%)	432 (94%)	25 (5%)	4 (1%)	14	48
5	L	463/613 (76%)	426 (92%)	33 (7%)	4 (1%)	14	48
All	All	6946/8222 (84%)	6278 (90%)	562 (8%)	106 (2%)	8	39

5 of 106 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	167	PRO
1	A	195	ARG
2	C	62	TYR
2	C	170	VAL
2	C	516	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	196/286 (68%)	180 (92%)	16 (8%)	9 34
1	B	184/286 (64%)	173 (94%)	11 (6%)	16 45
1	G	191/286 (67%)	180 (94%)	11 (6%)	17 46
1	H	183/286 (64%)	167 (91%)	16 (9%)	8 33
2	C	1152/1157 (100%)	1047 (91%)	105 (9%)	7 31
2	I	1151/1157 (100%)	1049 (91%)	102 (9%)	8 32
3	D	968/1168 (83%)	869 (90%)	99 (10%)	6 27
3	J	959/1168 (82%)	864 (90%)	95 (10%)	6 28
4	E	71/75 (95%)	66 (93%)	5 (7%)	12 40
4	K	67/75 (89%)	60 (90%)	7 (10%)	5 26
5	F	413/540 (76%)	389 (94%)	24 (6%)	17 46
5	L	410/540 (76%)	384 (94%)	26 (6%)	15 44
All	All	5945/7024 (85%)	5428 (91%)	517 (9%)	8 33

5 of 517 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	J	918	ILE
3	J	1261	LEU
3	J	910	ASN
3	D	797	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	749	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 66 such sidechains are listed below:

Mol	Chain	Res	Type
3	J	594	GLN
3	J	910	ASN
5	L	446	GLN
3	D	716	GLN
3	D	702	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ECJ	C	1401	-	30,37,37	2.97	10 (33%)	34,59,59	2.84	6 (17%)
6	ECJ	J	1501	-	30,37,37	2.97	9 (30%)	34,59,59	3.18	5 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ECJ	C	1401	-	-	5/21/43/43	0/3/3/3
6	ECJ	J	1501	-	-	5/21/43/43	0/3/3/3

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	1401	ECJ	O16-C15	9.99	1.39	1.22
6	J	1501	ECJ	O16-C15	9.86	1.39	1.22
6	C	1401	ECJ	C13-C11	7.23	1.49	1.35
6	J	1501	ECJ	C13-C11	7.10	1.49	1.35
6	J	1501	ECJ	P29-O32	5.86	1.65	1.59

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	J	1501	ECJ	O16-C15-C13	-13.92	109.93	126.89
6	C	1401	ECJ	O16-C15-C13	-12.82	111.27	126.89
6	J	1501	ECJ	C15-C13-C11	-8.53	100.82	109.40
6	C	1401	ECJ	C15-C13-C11	-7.30	102.06	109.40
6	J	1501	ECJ	N27-C25-N24	-6.67	119.61	128.67

There are no chirality outliers.

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	C	1401	ECJ	O09-C10-C11-C13
6	J	1501	ECJ	O09-C10-C11-C13
6	J	1501	ECJ	P01-O05-P06-O09
6	J	1501	ECJ	C10-O09-P06-O05
6	J	1501	ECJ	C10-O09-P06-O08


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	J	1501	ECJ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	228/329 (69%)	-0.78	0 100 100	90, 124, 199, 234	0
1	B	217/329 (65%)	-0.65	0 100 100	85, 176, 231, 252	0
1	G	224/329 (68%)	-0.69	0 100 100	137, 176, 229, 280	0
1	H	217/329 (65%)	-0.60	0 100 100	150, 198, 235, 274	0
2	C	1340/1342 (99%)	-0.72	0 100 100	67, 119, 204, 265	0
2	I	1340/1342 (99%)	-0.70	1 (0%) 92 88	100, 147, 231, 373	0
3	D	1163/1407 (82%)	-0.73	0 100 100	63, 114, 199, 265	0
3	J	1155/1407 (82%)	-0.71	0 100 100	82, 145, 217, 279	0
4	E	89/91 (97%)	-0.63	0 100 100	107, 148, 174, 186	0
4	K	79/91 (86%)	-0.51	1 (1%) 74 53	182, 265, 298, 321	0
5	F	467/613 (76%)	-0.66	0 100 100	97, 182, 330, 392	0
5	L	469/613 (76%)	-0.71	0 100 100	121, 192, 300, 372	0
All	All	6988/8222 (84%)	-0.70	2 (0%) 100 100	63, 145, 240, 392	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	K	16	ARG	2.4
2	I	1264	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	ECJ	J	1501	35/35	0.84	0.08	63,90,174,177	35
8	ZN	D	2003	1/1	0.86	0.30	377,377,377,377	0
6	ECJ	C	1401	35/35	0.89	0.08	74,99,183,185	35
7	MG	J	1502	1/1	0.96	0.05	81,81,81,81	0
8	ZN	D	2002	1/1	0.99	0.06	152,152,152,152	0
7	MG	D	2001	1/1	0.99	0.03	47,47,47,47	0
8	ZN	J	1503	1/1	0.99	0.01	155,155,155,155	0
8	ZN	J	1504	1/1	0.99	0.03	128,128,128,128	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ECJ J 1501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.