



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 10:44 am GMT

PDB ID : 6BZ0
Title : 1.83 Angstrom Resolution Crystal Structure of Dihydrolipoyl Dehydrogenase from *Acinetobacter baumannii* in Complex with FAD.
Authors : Minasov, G.; Shuvalova, L.; Dubrovskaya, I.; Kiryukhina, O.; Grimshaw, S.; Kwon, K.; Anderson, W.F.; Satchell, K.J.F.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2017-12-21
Resolution : 1.83 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3313 (1.86-1.82)
Clashscore	122126	3530 (1.86-1.82)
Ramachandran outliers	120053	3495 (1.86-1.82)
Sidechain outliers	120020	3496 (1.86-1.82)
RSRZ outliers	108989	3265 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	
1	B	480	
1	C	480	
1	D	480	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrolipoyl dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	469	Total 3625	C 2298	N 625	O 688	S 14	0	13	0
1	B	469	Total 3595	C 2277	N 623	O 682	S 13	0	9	0
1	C	468	Total 3573	C 2264	N 619	O 677	S 13	0	8	0
1	D	469	Total 3576	C 2263	N 618	O 682	S 13	0	7	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP D0CDT4
A	-1	ASN	-	expression tag	UNP D0CDT4
A	0	ALA	-	expression tag	UNP D0CDT4
A	355	ILE	VAL	engineered mutation	UNP D0CDT4
B	-2	SER	-	expression tag	UNP D0CDT4
B	-1	ASN	-	expression tag	UNP D0CDT4
B	0	ALA	-	expression tag	UNP D0CDT4
B	355	ILE	VAL	engineered mutation	UNP D0CDT4
C	-2	SER	-	expression tag	UNP D0CDT4
C	-1	ASN	-	expression tag	UNP D0CDT4
C	0	ALA	-	expression tag	UNP D0CDT4
C	355	ILE	VAL	engineered mutation	UNP D0CDT4
D	-2	SER	-	expression tag	UNP D0CDT4
D	-1	ASN	-	expression tag	UNP D0CDT4
D	0	ALA	-	expression tag	UNP D0CDT4
D	355	ILE	VAL	engineered mutation	UNP D0CDT4

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Cl	0	0
			3	3		
3	A	3	Total	Cl	0	0
			3	3		
3	D	1	Total	Cl	0	0
			1	1		
3	C	4	Total	Cl	0	0
			4	4		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	359	Total 365	O 365	0	8
5	B	248	Total 254	O 254	0	6
5	C	332	Total 342	O 342	0	10
5	D	295	Total 299	O 299	0	4


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dihydrolipoyl dehydrogenase




- Molecule 1: Dihydrolipoyl dehydrogenase




- Molecule 1: Dihydrolipoyl dehydrogenase



- Molecule 1: Dihydrolipoyl dehydrogenase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.97Å 82.77Å 96.26Å 80.38° 80.21° 89.49°	Depositor
Resolution (Å)	29.63 – 1.83 29.63 – 1.83	Depositor EDS
% Data completeness (in resolution range)	97.6 (29.63-1.83) 97.6 (29.63-1.83)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.83Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.179 , 0.222 0.187 , 0.229	Depositor DCC
R_{free} test set	7557 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtrriage
Anisotropy	0.244	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15854	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3686	0.68	0/4985
1	B	0.39	0/3656	0.65	0/4945
1	C	0.43	0/3634	0.68	0/4916
1	D	0.41	0/3637	0.66	0/4920
All	All	0.42	0/14613	0.67	0/19766

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3625	0	3652	15	0
1	B	3595	0	3618	12	0
1	C	3573	0	3603	20	0
1	D	3576	0	3589	14	0
2	A	53	0	31	1	0
2	B	53	0	31	0	0
2	C	53	0	31	0	0
2	D	53	0	31	0	0
3	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	3	0	0	0	0
3	C	4	0	0	1	0
3	D	1	0	0	0	0
4	A	2	0	0	0	0
5	A	365	0	0	0	0
5	B	254	0	0	0	0
5	C	342	0	0	0	0
5	D	299	0	0	0	0
All	All	15854	0	14586	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:VAL:HG13	1:B:246:GLY:HA2	1.72	0.72
1:B:48:CYS:HG	1:B:53:CYS:HG	1.38	0.70
1:C:53:CYS:O	1:C:57:LYS:HE2	2.01	0.61
1:C:459[A]:GLU:HA	1:C:459[A]:GLU:OE1	2.02	0.60
1:D:210[B]:GLU:HA	1:D:210[B]:GLU:OE1	2.02	0.60
1:A:48:CYS:HG	1:A:53:CYS:HG	1.46	0.60
1:D:57:LYS:HD3	1:D:57:LYS:N	2.17	0.59
1:C:234:GLN:NE2	1:C:359:HIS:O	2.28	0.59
1:D:159:VAL:HG11	1:D:244:VAL:HG12	1.86	0.58
1:C:404:PHE:CD1	1:C:413:LEU:HD11	2.40	0.56
1:D:386:PHE:CD1	1:D:456:VAL:HG21	2.41	0.56
1:C:370:GLU:HB2	3:C:505:CL:CL	2.43	0.56
1:A:18[B]:GLU:HG3	1:A:330:MET:HG2	1.87	0.55
1:D:159:VAL:HG13	1:D:246:GLY:HA2	1.88	0.55
1:B:49:LEU:HD22	1:B:102:THR:HG23	1.89	0.54
1:D:57:LYS:HE3	1:D:357:TYR:CD1	2.43	0.53
1:C:48:CYS:SG	1:C:53:CYS:SG	3.02	0.53
1:D:404:PHE:CD1	1:D:413:LEU:HD11	2.45	0.51
1:B:53:CYS:O	1:B:57:LYS:HE2	2.12	0.50
1:B:404:PHE:CD1	1:B:413:LEU:HD11	2.47	0.49
1:C:55:PRO:HB3	1:C:94:LYS:HD3	1.95	0.48
1:B:152:SER:HB3	1:B:318:ASP:HB3	1.96	0.48
1:C:48:CYS:HG	1:C:53:CYS:HG	1.49	0.48
1:C:124:LEU:HB2	1:C:128:LYS:HB2	1.96	0.48
1:C:472:ILE:H	1:C:472:ILE:HD12	1.80	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:CYS:O	1:A:57:LYS:HE2	2.15	0.46
1:B:166:ILE:HD11	1:B:252:ARG:CZ	2.46	0.46
1:C:57:LYS:HD2	1:C:357:TYR:CG	2.51	0.46
1:C:452:THR:O	1:C:455:GLU:HG2	2.16	0.45
1:D:159:VAL:HG13	1:D:246:GLY:CA	2.46	0.45
1:C:240:ILE:O	1:C:240:ILE:HD12	2.15	0.45
1:B:49:LEU:HD21	1:B:101:LEU:HB2	1.99	0.45
1:D:159:VAL:CG1	1:D:244:VAL:HG12	2.46	0.45
1:B:14:PRO:O	1:B:18:GLU:HB2	2.18	0.44
1:D:190:VAL:HG23	1:D:356:ILE:HG23	1.99	0.44
1:C:190[B]:VAL:HG22	1:C:356:ILE:HG12	1.99	0.44
1:A:218[B]:MET:CE	1:A:218[B]:MET:HA	2.47	0.44
1:A:48:CYS:SG	1:A:53:CYS:SG	3.05	0.44
1:D:412:ARG:HE	1:D:437:VAL:HG13	1.82	0.44
1:C:472:ILE:N	1:C:472:ILE:HD12	2.33	0.44
1:A:48:CYS:HG	1:A:53:CYS:HB2	1.83	0.43
1:B:124:LEU:HB2	1:B:128:LYS:HB2	2.00	0.43
1:D:166:ILE:HD11	1:D:252:ARG:CZ	2.48	0.43
1:C:89:LYS:HD3	1:D:77:GLY:HA2	2.01	0.43
1:D:117:LEU:HB3	1:D:131:PHE:CZ	2.53	0.43
1:C:117:LEU:HB3	1:C:131:PHE:CZ	2.54	0.42
1:A:117:LEU:HB3	1:A:131:PHE:CZ	2.55	0.42
1:B:117:LEU:HB3	1:B:131:PHE:CZ	2.54	0.42
1:C:386:PHE:CD1	1:C:456:VAL:HG21	2.54	0.42
1:C:326:ALA:O	1:C:330:MET:HG3	2.19	0.42
1:A:152:SER:HB3	1:A:318:ASP:HB3	2.02	0.41
1:B:57:LYS:HD2	1:B:357:TYR:CG	2.55	0.41
1:A:218[B]:MET:HA	1:A:218[B]:MET:HE2	2.01	0.41
1:A:386:PHE:CD1	1:A:456:VAL:HG21	2.55	0.41
1:A:404:PHE:CD1	1:A:413:LEU:HD11	2.55	0.41
1:A:53:CYS:SG	2:A:501:FAD:C10	3.09	0.41
1:A:48:CYS:HG	1:A:53:CYS:CB	2.34	0.41
1:A:49:LEU:CD1	1:A:105[A]:ILE:HD11	2.51	0.40
1:C:296:GLU:N	1:C:296:GLU:OE2	2.41	0.40
1:A:55:PRO:HB2	1:A:174:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	480/480 (100%)	469 (98%)	11 (2%)	0	100	100
1	B	476/480 (99%)	460 (97%)	15 (3%)	1 (0%)	49	34
1	C	474/480 (99%)	461 (97%)	13 (3%)	0	100	100
1	D	474/480 (99%)	462 (98%)	12 (2%)	0	100	100
All	All	1904/1920 (99%)	1852 (97%)	51 (3%)	1 (0%)	53	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	298	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/375 (101%)	375 (99%)	3 (1%)	83	77
1	B	374/375 (100%)	370 (99%)	4 (1%)	76	67
1	C	372/375 (99%)	369 (99%)	3 (1%)	83	77
1	D	372/375 (99%)	367 (99%)	5 (1%)	71	61
All	All	1496/1500 (100%)	1481 (99%)	15 (1%)	78	71

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LYS
1	A	94	LYS
1	A	321	ARG
1	B	57	LYS
1	B	94	LYS
1	B	260	GLN
1	B	321	ARG
1	C	57	LYS
1	C	94	LYS
1	C	321	ARG
1	D	68	ASP
1	D	94	LYS
1	D	164	GLN
1	D	321	ARG
1	D	467	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 13 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	501	-	51,58,58	1.55	9 (17%)	57,89,89	2.21	10 (17%)
2	FAD	B	501	-	51,58,58	1.56	8 (15%)	57,89,89	2.07	11 (19%)
2	FAD	C	501	-	51,58,58	1.52	7 (13%)	57,89,89	2.08	9 (15%)
2	FAD	D	501	-	51,58,58	1.52	8 (15%)	57,89,89	2.11	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/28/50/50	0/6/6/6
2	FAD	B	501	-	-	0/28/50/50	0/6/6/6
2	FAD	C	501	-	-	0/28/50/50	0/6/6/6
2	FAD	D	501	-	-	0/28/50/50	0/6/6/6

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FAD	C8A-N9A	-2.07	1.34	1.36
2	A	501	FAD	C10-N1	2.01	1.35	1.33
2	B	501	FAD	O4B-C1B	2.03	1.44	1.41
2	D	501	FAD	C10-N1	2.04	1.36	1.33
2	C	501	FAD	C2A-N3A	2.05	1.35	1.32
2	B	501	FAD	C2A-N3A	2.15	1.35	1.32
2	D	501	FAD	C2A-N3A	2.49	1.36	1.32
2	A	501	FAD	C4X-N5	2.57	1.37	1.33
2	B	501	FAD	C5A-C4A	2.72	1.46	1.40
2	A	501	FAD	C5A-C4A	2.77	1.46	1.40
2	D	501	FAD	C9A-N10	2.90	1.42	1.38
2	C	501	FAD	C5A-C4A	2.99	1.47	1.40
2	D	501	FAD	C5A-C4A	3.00	1.47	1.40
2	C	501	FAD	C9A-N10	3.02	1.42	1.38
2	A	501	FAD	C9A-N10	3.08	1.42	1.38
2	A	501	FAD	C8-C7	3.16	1.48	1.40
2	D	501	FAD	C8-C7	3.30	1.49	1.40
2	B	501	FAD	C8-C7	3.38	1.49	1.40
2	C	501	FAD	C8-C7	3.44	1.49	1.40
2	B	501	FAD	C9A-N10	3.48	1.43	1.38
2	A	501	FAD	C9A-C5X	3.73	1.50	1.42
2	C	501	FAD	C9A-C5X	3.74	1.50	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	FAD	C9A-C5X	3.87	1.50	1.42
2	D	501	FAD	C4-C4X	3.92	1.48	1.41
2	B	501	FAD	C9A-C5X	3.98	1.50	1.42
2	A	501	FAD	C4-C4X	4.12	1.48	1.41
2	C	501	FAD	C4-C4X	4.30	1.49	1.41
2	B	501	FAD	C4-C4X	4.44	1.49	1.41
2	C	501	FAD	C4X-C10	4.50	1.48	1.41
2	D	501	FAD	C4X-C10	4.56	1.48	1.41
2	B	501	FAD	C4X-C10	4.75	1.48	1.41
2	A	501	FAD	C4X-C10	4.93	1.49	1.41

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FAD	N3A-C2A-N1A	-7.68	122.29	128.86
2	C	501	FAD	N3A-C2A-N1A	-7.22	122.68	128.86
2	A	501	FAD	N3A-C2A-N1A	-7.15	122.74	128.86
2	D	501	FAD	N3A-C2A-N1A	-6.89	122.97	128.86
2	A	501	FAD	C4-C4X-C10	-6.59	115.03	119.95
2	D	501	FAD	C4-C4X-C10	-5.40	115.92	119.95
2	C	501	FAD	C4-C4X-C10	-4.70	116.44	119.95
2	B	501	FAD	C4-C4X-C10	-4.39	116.67	119.95
2	B	501	FAD	C4X-C4-N3	-3.16	118.97	123.47
2	C	501	FAD	C4X-C4-N3	-3.04	119.15	123.47
2	D	501	FAD	C4X-C4-N3	-2.94	119.29	123.47
2	A	501	FAD	C4X-C4-N3	-2.65	119.70	123.47
2	B	501	FAD	C4A-C5A-N7A	-2.60	106.90	109.41
2	B	501	FAD	C9A-N10-C10	-2.50	118.43	121.77
2	D	501	FAD	C4A-C5A-N7A	-2.46	107.03	109.41
2	A	501	FAD	C1B-N9A-C4A	-2.40	122.48	126.64
2	B	501	FAD	C1B-N9A-C4A	-2.40	122.49	126.64
2	A	501	FAD	C4A-C5A-N7A	-2.23	107.26	109.41
2	C	501	FAD	C9A-N10-C10	-2.13	118.93	121.77
2	D	501	FAD	C1B-N9A-C4A	-2.07	123.05	126.64
2	B	501	FAD	C2A-N1A-C6A	2.01	122.17	118.75
2	A	501	FAD	C2A-N1A-C6A	2.16	122.41	118.75
2	C	501	FAD	C2A-N1A-C6A	2.45	122.91	118.75
2	B	501	FAD	C4-C4X-N5	3.12	122.21	118.70
2	A	501	FAD	C4X-N5-C5X	3.53	120.46	116.76
2	C	501	FAD	C4-C4X-N5	3.58	122.73	118.70
2	D	501	FAD	C4-C4X-N5	3.66	122.81	118.70
2	D	501	FAD	C4X-N5-C5X	3.83	120.77	116.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FAD	C4X-N5-C5X	4.13	121.08	116.76
2	A	501	FAD	C4-C4X-N5	4.31	123.55	118.70
2	C	501	FAD	C4X-N5-C5X	4.36	121.33	116.76
2	A	501	FAD	C1'-N10-C9A	5.32	123.04	118.31
2	B	501	FAD	C1'-N10-C9A	5.37	123.08	118.31
2	C	501	FAD	C1'-N10-C9A	5.69	123.36	118.31
2	D	501	FAD	C1'-N10-C9A	5.81	123.47	118.31
2	C	501	FAD	C4-N3-C2	7.43	121.47	115.14
2	B	501	FAD	C4-N3-C2	7.61	121.62	115.14
2	D	501	FAD	C4-N3-C2	8.19	122.11	115.14
2	A	501	FAD	C4-N3-C2	8.19	122.12	115.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	469/480 (97%)	-0.18	9 (1%) 66 65	17, 27, 43, 98	0
1	B	469/480 (97%)	0.03	15 (3%) 47 45	20, 34, 63, 106	0
1	C	468/480 (97%)	-0.14	10 (2%) 63 62	16, 27, 43, 95	0
1	D	469/480 (97%)	0.00	21 (4%) 33 31	19, 30, 55, 97	0
All	All	1875/1920 (97%)	-0.07	55 (2%) 51 49	16, 29, 54, 106	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	297	ARG	6.3
1	B	297	ARG	6.3
1	D	471	ALA	6.2
1	A	298	GLY	5.3
1	D	467	ARG	5.2
1	D	469	ILE	4.8
1	B	295	THR	4.7
1	C	296	GLU	4.6
1	D	468	ALA	4.5
1	D	294	LEU	4.4
1	C	298	GLY	4.2
1	B	472	ILE	4.0
1	B	296	GLU	3.9
1	C	295	THR	3.8
1	A	296	GLU	3.7
1	A	294	LEU	3.6
1	D	3	GLN	3.5
1	C	294	LEU	3.4
1	D	39	ASN	3.3
1	B	298	GLY	3.3
1	B	261	ALA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	298	GLY	3.2
1	D	296	GLU	3.1
1	D	375	LYS	3.1
1	C	297	ARG	3.1
1	A	472	ILE	3.0
1	D	297	ARG	3.0
1	A	295	THR	3.0
1	D	465	ASP	3.0
1	C	211	ALA	2.9
1	B	4	GLN	2.9
1	D	261	ALA	2.9
1	B	39	ASN	2.8
1	B	137	GLU	2.8
1	B	142	GLU	2.8
1	D	374	GLU	2.7
1	B	37	ILE	2.7
1	A	343	HIS	2.6
1	B	299	LEU	2.5
1	C	472	ILE	2.5
1	D	466	GLY	2.4
1	B	136	GLY	2.3
1	A	388	VAL	2.3
1	D	344	ALA	2.2
1	C	23	ALA	2.2
1	D	38	HIS	2.2
1	D	41	LYS	2.2
1	A	374	GLU	2.2
1	B	135	GLU	2.1
1	D	40	GLY	2.1
1	D	293	LYS	2.1
1	B	288	GLU	2.1
1	D	470	HIS	2.1
1	C	374	GLU	2.1
1	C	293	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	B	502	1/1	0.79	0.14	50,50,50,50	1
3	CL	C	504	1/1	0.80	0.24	47,47,47,47	1
3	CL	B	503	1/1	0.84	0.07	73,73,73,73	0
4	MG	A	506	1/1	0.93	0.19	47,47,47,47	0
2	FAD	B	501	53/53	0.94	0.11	27,35,39,41	0
3	CL	A	503	1/1	0.95	0.06	44,44,44,44	0
3	CL	A	504	1/1	0.95	0.05	39,39,39,39	0
2	FAD	C	501	53/53	0.96	0.09	17,21,29,35	0
2	FAD	A	501	53/53	0.96	0.09	19,21,24,27	0
2	FAD	D	501	53/53	0.96	0.10	22,26,30,35	0
3	CL	B	504	1/1	0.96	0.11	36,36,36,36	0
4	MG	A	505	1/1	0.97	0.08	36,36,36,36	0
3	CL	C	503	1/1	0.98	0.04	43,43,43,43	0
3	CL	C	505	1/1	0.98	0.07	35,35,35,35	0
3	CL	D	502	1/1	0.98	0.07	42,42,42,42	0
3	CL	A	502	1/1	0.99	0.03	30,30,30,30	0
3	CL	C	502	1/1	0.99	0.05	29,29,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.