



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2023 – 10:37 PM JST

PDB ID : 7C0S  
Title : Crystal structure of a dinucleotide-binding protein (F79A) of ABC transporter endogenously bound to uridylyl-3'-5'-phospho-guanosine (Form II)  
Authors : Kanaujia, S.P.; Chandravanshi, M.; Samanta, R.  
Deposited on : 2020-05-01  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	 % 88% 9% ..
1	B	406	 % 87% 10% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	504	-	-	X	-
7	PGR	B	505	-	-	-	X
8	PGE	B	506	-	-	X	-

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sugar ABC transporter, periplasmic sugar-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	396	Total	C	N	O	S	0	2	0
			3053	1973	513	563	4			
1	B	397	Total	C	N	O	S	0	6	0
			3091	1996	524	567	4			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP Q5SLB4
A	0	MET	-	expression tag	UNP Q5SLB4
A	79	ALA	PHE	engineered mutation	UNP Q5SLB4
A	399	HIS	-	expression tag	UNP Q5SLB4
A	400	HIS	-	expression tag	UNP Q5SLB4
A	401	HIS	-	expression tag	UNP Q5SLB4
A	402	HIS	-	expression tag	UNP Q5SLB4
A	403	HIS	-	expression tag	UNP Q5SLB4
A	404	HIS	-	expression tag	UNP Q5SLB4
B	-1	MET	-	initiating methionine	UNP Q5SLB4
B	0	MET	-	expression tag	UNP Q5SLB4
B	79	ALA	PHE	engineered mutation	UNP Q5SLB4
B	399	HIS	-	expression tag	UNP Q5SLB4
B	400	HIS	-	expression tag	UNP Q5SLB4
B	401	HIS	-	expression tag	UNP Q5SLB4
B	402	HIS	-	expression tag	UNP Q5SLB4
B	403	HIS	-	expression tag	UNP Q5SLB4
B	404	HIS	-	expression tag	UNP Q5SLB4

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is [(1S,3R,3aR,6aS)-3-(2-azanyl-6-oxidanylidene-1H-purin-9-yl)-5,5-bis(oxidanyl)-1,3,3a,4,6,6a-hexahydrocyclopenta[c]furan-1-yl]methyl [(2R,3S,4R,5R)-5-[2,4-bis(oxidanylidene)pyrimidin-1-yl]-2-(hydroxymethyl)-4-oxidanyl-oxolan-3-yl] hydrogen phosphate (three-letter code: FGO) (formula: C<sub>22</sub>H<sub>28</sub>N<sub>7</sub>O<sub>13</sub>P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
4	A	1	43	22	7	13	1	0	0
4	B	1	43	22	7	13	1	0	0

- Molecule 5 is CARBON DIOXIDE (three-letter code: CO2) (formula: CO<sub>2</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
5	B	1	3	1 2	0	0

- Molecule 6 is CARBONATE ION (three-letter code: CO3) (formula: CO<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	1	3		

- Molecule 7 is R-1,2-PROPANEDIOL (three-letter code: PGR) (formula:  $C_3H_8O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			5	3	2		

- Molecule 8 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	379	Total	O	0	0
			379	379		
9	B	345	Total	O	0	0
			345	345		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sugar ABC transporter, periplasmic sugar-binding protein



- Molecule 1: Sugar ABC transporter, periplasmic sugar-binding protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.54Å 58.36Å 123.82Å 90.00° 94.81° 90.00°	Depositor
Resolution (Å)	60.56 – 1.85 60.56 – 1.85	Depositor EDS
% Data completeness (in resolution range)	91.3 (60.56-1.85) 91.3 (60.56-1.85)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.49 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.205 , 0.240 0.213 , 0.248	Depositor DCC
$R_{free}$ test set	3553 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtrriage
Anisotropy	0.251	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6986	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FGO, CO2, PGE, PGR, CO3, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.90	3/3128 (0.1%)	0.98	6/4244 (0.1%)
1	B	0.94	5/3178 (0.2%)	1.01	10/4309 (0.2%)
All	All	0.92	8/6306 (0.1%)	1.00	16/8553 (0.2%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	287[A]	GLU	CD-OE2	9.40	1.35	1.25
1	B	287[B]	GLU	CD-OE2	9.40	1.35	1.25
1	A	370	GLU	CD-OE2	6.15	1.32	1.25
1	B	370[A]	GLU	CD-OE2	5.77	1.31	1.25
1	B	370[B]	GLU	CD-OE2	5.77	1.31	1.25
1	A	131	LEU	C-O	5.20	1.33	1.23
1	A	80	GLU	CD-OE2	5.20	1.31	1.25
1	B	80	GLU	CD-OE2	5.07	1.31	1.25

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	342	ARG	NE-CZ-NH2	-8.14	116.23	120.30
1	B	342	ARG	NE-CZ-NH2	-7.29	116.66	120.30
1	B	287[A]	GLU	CB-CA-C	7.17	124.74	110.40
1	B	287[B]	GLU	CB-CA-C	7.17	124.74	110.40
1	A	342	ARG	NE-CZ-NH1	7.11	123.85	120.30
1	B	342	ARG	NE-CZ-NH1	6.88	123.74	120.30
1	B	108	THR	CA-CB-CG2	6.55	121.56	112.40
1	A	108	THR	CA-CB-CG2	6.07	120.89	112.40
1	A	236	PHE	CB-CA-C	-5.87	98.65	110.40
1	B	61	THR	CA-CB-OG1	-5.76	96.90	109.00
1	B	327	TYR	CB-CG-CD1	-5.58	117.65	121.00

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	327	TYR	CB-CG-CD1	-5.54	117.68	121.00
1	B	360	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	B	249	ARG	NE-CZ-NH2	5.39	122.99	120.30
1	B	236	PHE	CB-CA-C	-5.12	100.16	110.40
1	A	49	ARG	NE-CZ-NH2	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3053	0	3085	22	0
1	B	3091	0	3136	36	0
2	A	1	0	0	1	0
2	B	1	0	0	1	0
3	A	4	0	6	2	0
3	B	4	0	6	7	0
4	A	43	0	0	0	0
4	B	43	0	0	4	0
5	B	3	0	0	0	0
6	B	4	0	0	0	0
7	B	5	0	8	0	0
8	B	10	0	14	12	0
9	A	379	0	0	10	0
9	B	345	0	0	11	0
All	All	6986	0	6255	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:LEU:HD12	9:A:801:HOH:O	1.71	0.89

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:504:EDO:H22	4:B:507:FGO:C14	2.02	0.89
1:A:106:ASN:OD1	1:A:108:THR:HB	1.74	0.87
2:A:501:CL:CL	9:A:776:HOH:O	2.31	0.85
1:B:106:ASN:OD1	1:B:108:THR:HB	1.79	0.83
1:A:370:GLU:HG2	9:A:740:HOH:O	1.82	0.78
1:B:154:ALA:HA	8:B:506:PGE:H5	1.66	0.77
1:B:355:GLU:OE2	1:B:396:ARG:NE	2.25	0.70
1:A:151:GLU:OE2	1:A:265:LYS:NZ	2.26	0.68
1:B:154:ALA:HA	8:B:506:PGE:C5	2.23	0.68
1:B:158:LYS:HB2	8:B:506:PGE:H42	1.77	0.66
1:A:297:GLU:OE2	9:A:601:HOH:O	2.14	0.65
1:A:108:THR:HG21	9:A:625:HOH:O	1.97	0.64
1:B:274:GLN:HB3	3:B:504:EDO:H21	1.79	0.64
1:B:158:LYS:HD2	8:B:506:PGE:H2	1.81	0.62
1:A:0:MET:HB3	1:A:4:ASP:HB2	1.81	0.62
1:B:64:LYS:HE2	1:B:85:LEU:CD2	2.29	0.61
1:B:249:ARG:NE	9:B:608:HOH:O	2.33	0.61
1:B:154:ALA:HB1	8:B:506:PGE:H52	1.84	0.59
1:A:263:ARG:NH1	9:A:603:HOH:O	2.30	0.58
1:A:268:PRO:HG3	3:A:502:EDO:H22	1.85	0.58
1:B:126:LYS:NZ	9:B:611:HOH:O	2.37	0.57
1:B:275:GLY:O	3:B:504:EDO:H12	2.04	0.56
1:B:93:LEU:HD11	1:B:289:GLN:CD	2.25	0.56
8:B:506:PGE:C6	9:B:870:HOH:O	2.54	0.56
8:B:506:PGE:H6	9:B:870:HOH:O	2.06	0.56
1:B:154:ALA:O	8:B:506:PGE:H4	2.06	0.55
1:B:38:GLU:HG3	9:B:875:HOH:O	2.06	0.55
1:B:1:LYS:HE2	9:B:887:HOH:O	2.07	0.55
1:B:105:VAL:HG11	1:B:110:LEU:HD13	1.90	0.54
1:B:154:ALA:CB	8:B:506:PGE:H52	2.38	0.54
1:A:80:GLU:HG2	1:A:112:ALA:HB1	1.91	0.53
1:B:64:LYS:HE2	1:B:85:LEU:HD23	1.93	0.50
1:A:105:VAL:HG11	1:A:110:LEU:HD13	1.94	0.50
1:A:333[A]:GLU:HG2	9:A:736:HOH:O	2.11	0.49
1:A:105:VAL:CG1	1:A:110:LEU:HD13	2.42	0.48
1:B:105:VAL:CG1	1:B:110:LEU:HD13	2.43	0.48
1:B:6:ILE:HG21	1:B:50:PRO:HD2	1.95	0.47
1:B:158:LYS:HB2	8:B:506:PGE:C4	2.44	0.47
4:B:507:FGO:C19	9:B:607:HOH:O	2.63	0.47
3:B:504:EDO:H22	4:B:507:FGO:N3	2.26	0.47
1:A:6:ILE:HG21	1:A:50:PRO:HD2	1.96	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:ARG:HD2	9:A:931:HOH:O	2.15	0.46
1:A:125:ASN:O	1:A:275:GLY:HA3	2.16	0.45
1:B:336:ASP:OD1	9:B:601:HOH:O	2.21	0.45
1:B:151:GLU:OE2	1:B:265:LYS:NZ	2.50	0.45
1:A:170:PHE:HB2	1:A:176:THR:HG21	1.99	0.45
1:B:80:GLU:HG2	1:B:112:ALA:HB1	1.99	0.44
2:B:501:CL:CL	9:B:711:HOH:O	2.58	0.44
1:B:170:PHE:HB2	1:B:176:THR:HG21	2.00	0.43
1:B:275:GLY:H	3:B:504:EDO:H21	1.82	0.43
1:B:125:ASN:O	1:B:275:GLY:HA3	2.19	0.42
1:A:7:LYS:HE3	9:A:606:HOH:O	2.18	0.42
1:B:158:LYS:CD	8:B:506:PGE:H2	2.49	0.42
1:A:15:VAL:HG22	1:A:291:VAL:HG21	2.02	0.42
8:B:506:PGE:H62	9:B:870:HOH:O	2.18	0.42
1:A:37:VAL:HG12	1:A:41:LYS:HE3	2.02	0.42
1:B:213:LYS:HB3	1:B:213:LYS:HE3	1.93	0.41
1:A:46:ARG:HB3	1:A:291:VAL:HG22	2.03	0.41
1:B:64:LYS:HE2	1:B:85:LEU:HD21	2.03	0.41
1:B:88:GLU:OE2	1:B:396:ARG:NH2	2.48	0.41
1:B:275:GLY:H	3:B:504:EDO:C2	2.34	0.41
1:B:355:GLU:OE2	1:B:396:ARG:CZ	2.68	0.41
3:B:504:EDO:H22	4:B:507:FGO:N2	2.34	0.41
3:A:502:EDO:C2	9:A:909:HOH:O	2.69	0.40
1:B:82:ASN:ND2	9:B:645:HOH:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	396/406 (98%)	389 (98%)	7 (2%)	0	<b>100</b> <b>100</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	401/406 (99%)	395 (98%)	6 (2%)	0	100	100
All	All	797/812 (98%)	784 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/318 (98%)	303 (98%)	7 (2%)	50	34
1	B	315/318 (99%)	309 (98%)	6 (2%)	57	43
All	All	625/636 (98%)	612 (98%)	13 (2%)	53	38

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	MET
1	A	8	GLU
1	A	14	LYS
1	A	15	VAL
1	A	98	LEU
1	A	149	LEU
1	A	226	ASN
1	B	14	LYS
1	B	15	VAL
1	B	71	LYS
1	B	98	LEU
1	B	149	LEU
1	B	226	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	82	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	PGR	B	505	-	3,4,4	0.09	0	1,4,4	0.16	0
8	PGE	B	506	-	9,9,9	0.33	0	8,8,8	0.23	0
4	FGO	A	503	-	44,48,48	1.46	9 (20%)	50,75,75	2.03	9 (18%)
3	EDO	B	504	-	3,3,3	0.40	0	2,2,2	0.56	0
4	FGO	B	507	-	44,48,48	1.47	4 (9%)	50,75,75	2.12	16 (32%)
6	CO3	B	503	-	2,3,3	0.65	0	2,3,3	0.18	0
5	CO2	B	502	-	2,2,2	0.20	0	1,1,1	0.72	0
3	EDO	A	502	-	3,3,3	0.32	0	2,2,2	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PGR	B	505	-	-	2/2/2/2	-
8	PGE	B	506	-	-	5/7/7/7	-
4	FGO	A	503	-	-	0/17/63/63	0/6/6/6
3	EDO	B	504	-	-	1/1/1/1	-
4	FGO	B	507	-	-	0/17/63/63	0/6/6/6
3	EDO	A	502	-	-	1/1/1/1	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	507	FGO	C5-N1	-3.94	1.30	1.38
4	A	503	FGO	O5-C6	3.63	1.31	1.24
4	B	507	FGO	C19-C18	-3.56	1.48	1.53
4	A	503	FGO	C5-N1	-3.45	1.31	1.38
4	B	507	FGO	O5-C6	3.45	1.31	1.24
4	B	507	FGO	C8-C7	2.95	1.41	1.35
4	A	503	FGO	C21-C17	2.70	1.58	1.53
4	A	503	FGO	C8-C7	2.63	1.41	1.35
4	A	503	FGO	C18-C17	-2.58	1.48	1.55
4	A	503	FGO	C16-N5	2.38	1.38	1.33
4	A	503	FGO	C5-N	2.17	1.41	1.38
4	A	503	FGO	O7-C11	2.11	1.44	1.41
4	A	503	FGO	O9-C20	2.05	1.46	1.40

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	507	FGO	O10-C20-C21	6.57	132.28	110.56
4	A	503	FGO	N1-C5-N	6.23	123.16	114.89
4	A	503	FGO	C6-N1-C5	-5.21	119.71	126.58
4	A	503	FGO	O8-C15-C13	-4.83	114.94	124.37
4	B	507	FGO	N1-C5-N	4.78	121.23	114.89
4	A	503	FGO	O9-C20-C21	4.76	126.30	110.56
4	A	503	FGO	O8-C15-N4	4.53	126.00	120.65
4	B	507	FGO	C17-C18-C10	4.25	111.68	103.72
4	B	507	FGO	O8-C15-C13	-4.18	116.22	124.37
4	B	507	FGO	C6-N1-C5	-3.63	121.79	126.58
4	B	507	FGO	C7-C6-N1	3.60	120.23	114.84
4	A	503	FGO	C7-C6-N1	3.34	119.84	114.84
4	B	507	FGO	O8-C15-N4	3.23	124.47	120.65
4	B	507	FGO	C13-C15-N4	3.00	119.25	113.95
4	B	507	FGO	O5-C6-N1	-2.91	115.03	119.31
4	B	507	FGO	C16-N4-C15	-2.83	119.88	125.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	FGO	C13-C15-N4	2.83	118.95	113.95
4	A	503	FGO	O4-C5-N1	-2.62	116.62	121.50
4	B	507	FGO	O10-C20-C19	-2.58	102.02	110.56
4	B	507	FGO	O9-C20-C19	2.52	118.92	110.56
4	A	503	FGO	C2-N-C8	2.37	126.01	120.84
4	B	507	FGO	C2-N-C8	2.32	125.90	120.84
4	B	507	FGO	O4-C5-N1	-2.18	117.43	121.50
4	B	507	FGO	C7-C8-N	-2.18	118.16	121.81
4	B	507	FGO	C21-C17-C11	2.05	116.95	113.62

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	505	PGR	O1-C1-C2-C3
7	B	505	PGR	O1-C1-C2-O2
8	B	506	PGE	O2-C3-C4-O3
8	B	506	PGE	O3-C5-C6-O4
3	A	502	EDO	O1-C1-C2-O2
3	B	504	EDO	O1-C1-C2-O2
8	B	506	PGE	C4-C3-O2-C2
8	B	506	PGE	C6-C5-O3-C4
8	B	506	PGE	C1-C2-O2-C3


There are no ring outliers.

4 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	506	PGE	12	0
3	B	504	EDO	7	0
4	B	507	FGO	4	0
3	A	502	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	396/406 (97%)	-0.12	4 (1%) 82 82	10, 21, 38, 67	0
1	B	397/406 (97%)	-0.11	6 (1%) 73 74	11, 22, 40, 69	0
All	All	793/812 (97%)	-0.12	10 (1%) 77 78	10, 21, 39, 69	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	44	GLN	4.9
1	A	44	GLN	4.6
1	A	0	MET	4.3
1	B	68	ALA	3.2
1	A	8	GLU	2.9
1	B	67	PHE	2.6
1	B	0	MET	2.6
1	A	12	ARG	2.5
1	B	161[A]	ARG	2.2
1	B	396	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	PGR	B	505	5/5	0.61	0.53	62,65,68,69	0
8	PGE	B	506	10/10	0.81	0.27	23,36,45,50	0
3	EDO	A	502	4/4	0.84	0.17	32,36,38,41	0
6	CO3	B	503	4/4	0.84	0.23	49,53,55,56	0
3	EDO	B	504	4/4	0.90	0.28	28,28,31,33	0
5	CO2	B	502	3/3	0.92	0.11	33,33,35,38	0
4	FGO	A	503	43/43	0.93	0.12	13,20,30,36	0
4	FGO	B	507	43/43	0.93	0.13	15,21,32,39	0
2	CL	B	501	1/1	0.97	0.07	37,37,37,37	0
2	CL	A	501	1/1	0.98	0.08	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.