



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 04:13 am BST

PDB ID : 2C19  
Title : 5-(4-Carboxy-2-oxo-butylsulfanyl)-4-oxo-pentanoic acid acid bound to Porphobilinogen synthase from *Pseudomonas aeruginosa*  
Authors : Frere, F.; Nentwich, M.; Gacond, S.; Heinz, D.W.; Neier, R.; Frankenberg-Dinkel, N.  
Deposited on : 2005-09-11  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	 86% 12% ..
1	B	337	 83% 13% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	C1X	A	205	-	-	X	-
1	C1X	B	205	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6273 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DELTA-AMINOLEVULINIC ACID DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	2808	1759	503	532	14	0	24	1
1	B	329	2802	1752	498	538	14	0	28	1

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	VAL	ILE	conflict	UNP Q59643
B	199	VAL	ILE	conflict	UNP Q59643

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	327	Total	O	0	0
			327	327		
4	B	332	Total	O	0	0
			332	332		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.20Å 127.20Å 85.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	89.80 – 2.05 34.26 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.8 (89.80-2.05) 97.8 (34.26-2.05)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.95 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.144 , 0.201 0.145 , 0.203	Depositor DCC
$R_{free}$ test set	2220 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.0	Xtrriage
Anisotropy	0.232	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 52.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6273	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, C1X

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	0/2833	0.75	1/3836 (0.0%)
1	B	0.75	0/2827	0.80	3/3831 (0.1%)
All	All	0.76	0/5660	0.77	4/7667 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	198	ARG	NE-CZ-NH2	-11.23	114.69	120.30
1	B	198	ARG	NE-CZ-NH1	7.35	123.97	120.30
1	B	271	ARG	NE-CZ-NH2	-5.49	117.55	120.30
1	A	301	LEU	CB-CG-CD1	5.25	119.92	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	204	ALA	Mainchain
1	B	204	ALA	Mainchain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	B	205	C1X	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2808	0	2764	58	0
1	B	2802	0	2723	55	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	2	0	0	0	0
4	A	327	0	0	22	0
4	B	332	0	0	8	0
All	All	6273	0	5487	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:C1X:C8	1:A:260:LYS:HZ1	1.05	1.56
1:B:205:C1X:C8	1:B:260:LYS:HZ1	0.97	1.54
1:B:136[A]:HIS:ND1	1:B:138[A]:GLN:HB2	1.49	1.27
1:A:205:C1X:H8	1:A:260:LYS:NZ	0.89	1.22
1:B:205:C1X:H8	1:B:260:LYS:NZ	0.86	1.18
1:A:263[B]:MET:HG2	1:B:263[B]:MET:HG2	1.23	1.12
1:A:307[B]:LEU:HD21	4:A:2153:HOH:O	1.46	1.12
1:A:263[B]:MET:SD	1:B:263[B]:MET:SD	2.49	1.11
1:A:275[A]:GLU:HG2	4:A:2291:HOH:O	1.55	1.06
1:B:303[B]:GLU:OE2	4:B:2318:HOH:O	1.77	1.00
1:A:74[B]:GLU:HG2	4:A:2117:HOH:O	1.64	0.96
1:A:263[B]:MET:CG	1:B:263[B]:MET:HG2	1.95	0.96
1:A:205:C1X:H8	1:A:260:LYS:HZ2	1.35	0.90
1:B:205:C1X:H8	1:B:260:LYS:HZ2	1.34	0.90
1:A:263[B]:MET:HG2	1:B:263[B]:MET:CG	2.03	0.87

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74[B]:GLU:CD	4:A:2117:HOH:O	2.13	0.87
1:B:136[A]:HIS:ND1	1:B:138[A]:GLN:CB	2.38	0.85
1:B:205:C1X:H8	1:B:260:LYS:HZ3	1.39	0.84
1:A:263[A]:MET:HG3	4:A:2313:HOH:O	1.78	0.84
1:B:205:C1X:H8	1:B:260:LYS:CE	2.09	0.82
1:A:74[B]:GLU:CG	4:A:2117:HOH:O	2.21	0.82
1:A:94[B]:LYS:NZ	4:A:2134:HOH:O	2.07	0.81
1:A:205:C1X:H8	1:A:260:LYS:CE	2.09	0.81
1:A:205:C1X:H8	1:A:260:LYS:HZ3	1.44	0.78
1:A:205:C1X:C3	1:A:260:LYS:HZ1	1.94	0.78
1:B:205:C1X:C3	1:B:260:LYS:HZ1	1.95	0.77
1:A:205:C1X:C8	1:A:260:LYS:NZ	1.85	0.77
1:B:118[A]:ARG:NH2	4:B:2188:HOH:O	2.17	0.75
1:B:66[B]:ASP:OD1	4:B:2128:HOH:O	2.03	0.75
1:A:307[B]:LEU:CD2	4:A:2153:HOH:O	2.16	0.72
1:B:198:ARG:N	1:B:198:ARG:HD3	2.05	0.70
1:A:74[A]:GLU:HG2	4:A:2117:HOH:O	1.91	0.69
1:B:205:C1X:C8	1:B:260:LYS:NZ	1.83	0.69
1:B:189:LEU:HD22	1:B:197[B]:VAL:HG11	1.75	0.67
1:B:143:ASP:HB3	1:B:149:LEU:HD21	1.77	0.67
1:A:275[A]:GLU:OE2	4:A:2291:HOH:O	2.11	0.67
1:A:275[A]:GLU:CG	4:A:2291:HOH:O	2.25	0.66
1:A:35[A]:VAL:HG11	4:A:2122:HOH:O	1.94	0.66
1:B:205:C1X:C9	1:B:260:LYS:HZ1	2.02	0.65
1:A:307[B]:LEU:HG	4:A:2314:HOH:O	1.95	0.65
1:B:118[A]:ARG:NE	4:B:2188:HOH:O	2.28	0.64
1:A:136[A]:HIS:HD2	1:A:138:GLN:H	1.44	0.63
1:B:194:HIS:O	1:B:197[B]:VAL:HG12	1.98	0.63
1:A:263[A]:MET:CG	4:A:2313:HOH:O	2.39	0.62
1:A:263[B]:MET:SD	1:B:263[B]:MET:CG	2.87	0.62
1:B:38:LEU:O	1:B:81:PRO:HD2	2.01	0.60
1:A:293[B]:MET:O	1:A:297:GLN:HG3	2.02	0.59
1:A:136[A]:HIS:HE1	4:A:2139:HOH:O	1.85	0.58
1:A:263[B]:MET:SD	1:B:263[B]:MET:HG2	2.43	0.58
1:A:83:LEU:O	1:A:125[B]:ILE:HG12	2.03	0.58
1:B:335:ARG:HG3	1:B:336:GLY:N	2.19	0.58
1:A:94[B]:LYS:HE2	1:A:223:ASN:ND2	2.21	0.56
1:A:205:C1X:C7	1:A:260:LYS:NZ	2.66	0.55
1:A:8:ARG:HH11	1:B:244[B]:HIS:CD2	2.25	0.55
1:B:166[B]:GLU:OE2	4:B:2234:HOH:O	2.17	0.55
1:B:136[A]:HIS:C	1:B:138[A]:GLN:H	2.11	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:205:C1X:C7	1:B:260:LYS:NZ	2.69	0.53
1:A:105:GLU:O	1:A:110[A]:ARG:HD3	2.09	0.53
1:A:159[B]:ARG:CZ	4:A:2203:HOH:O	2.58	0.52
1:A:78:LEU:HD23	1:A:334:ARG:HD3	1.93	0.51
1:A:301:LEU:HG	1:A:305:VAL:HG11	1.93	0.50
1:A:72:ALA:HA	1:A:75:TRP:CE3	2.46	0.50
1:A:130:LEU:HD12	1:A:140:GLY:HA2	1.92	0.50
1:A:42[B]:VAL:HG21	1:A:68:LEU:HD13	1.94	0.49
1:A:303[B]:GLU:OE1	4:A:2305:HOH:O	2.20	0.48
1:A:94[B]:LYS:HE2	1:A:223:ASN:HD21	1.77	0.48
1:A:266:LEU:HB2	1:B:266:LEU:HB2	1.96	0.48
1:B:288:GLU:HA	1:B:291:MET:HE2	1.95	0.48
1:B:118[A]:ARG:CZ	4:B:2188:HOH:O	2.53	0.48
1:B:136[A]:HIS:C	1:B:138[A]:GLN:N	2.67	0.48
4:A:2007:HOH:O	1:B:244[B]:HIS:HD2	1.98	0.46
1:B:136[A]:HIS:CE1	1:B:138[A]:GLN:HB2	2.39	0.46
1:B:332:GLN:O	1:B:336:GLY:N	2.48	0.46
1:B:131:ASP:N	1:B:132:PRO:CD	2.79	0.45
1:B:205:C1X:C3	1:B:260:LYS:NZ	2.67	0.45
1:A:143:ASP:HB3	1:A:149:LEU:HD21	1.98	0.45
1:B:198:ARG:HA	1:B:255:ASP:OD2	2.17	0.45
1:A:159[B]:ARG:NH2	4:A:2203:HOH:O	2.51	0.44
1:A:3:PHE:N	4:A:2002:HOH:O	2.50	0.44
1:B:135[B]:THR:HG22	1:B:136[B]:HIS:CD2	2.53	0.44
1:B:130:LEU:HD12	1:B:140:GLY:HA2	1.99	0.44
1:A:205:C1X:H7C2	1:A:214:PHE:CZ	2.53	0.44
1:A:263[B]:MET:CG	1:B:263[B]:MET:CG	2.76	0.44
1:A:136[A]:HIS:CD2	1:A:138:GLN:H	2.31	0.43
1:B:198:ARG:N	1:B:198:ARG:CD	2.77	0.43
1:B:131:ASP:N	1:B:132:PRO:HD3	2.33	0.43
1:B:72:ALA:HA	1:B:75:TRP:CE3	2.53	0.43
1:A:187[B]:GLU:CD	4:A:2221:HOH:O	2.56	0.43
1:B:335:ARG:CG	1:B:336:GLY:N	2.82	0.43
1:A:42[B]:VAL:HG21	1:A:68:LEU:CD1	2.49	0.42
1:B:118[A]:ARG:HD3	4:B:2186:HOH:O	2.19	0.42
1:A:178:MET:O	1:A:181:ARG:HD2	2.20	0.42
1:A:263[B]:MET:CG	1:B:263[B]:MET:SD	3.08	0.42
1:A:35[A]:VAL:HG22	4:A:2077:HOH:O	2.20	0.42
1:A:45:LEU:HD21	1:A:62[A]:ARG:HD3	2.01	0.42
1:B:200:MET:HG3	1:B:256[A]:MET:HG2	2.01	0.42
1:A:136[A]:HIS:CD2	1:A:138:GLN:HB2	2.55	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:327:LYS:O	1:B:331[A]:GLU:HG3	2.20	0.41
1:B:136[A]:HIS:CE1	1:B:138[A]:GLN:CB	3.02	0.41
1:A:205:C1X:C9	1:A:260:LYS:NZ	2.76	0.40
1:B:71[B]:GLU:OE1	4:B:2133:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/337 (105%)	346 (98%)	8 (2%)	1 (0%)	41	31
1	B	352/337 (104%)	342 (97%)	10 (3%)	0	100	100
All	All	707/674 (105%)	688 (97%)	18 (2%)	1 (0%)	47	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	335	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/268 (108%)	285 (99%)	4 (1%)	67	65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	289/268 (108%)	275 (95%)	14 (5%)	25	18
All	All	578/536 (108%)	560 (97%)	18 (3%)	42	33

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	PHE
1	A	181	ARG
1	A	283	TYR
1	A	301	LEU
1	B	32[A]	VAL
1	B	32[B]	VAL
1	B	50	GLN
1	B	74	GLU
1	B	139	ASP
1	B	181	ARG
1	B	198	ARG
1	B	256[A]	MET
1	B	256[B]	MET
1	B	275[A]	GLU
1	B	275[B]	GLU
1	B	283	TYR
1	B	307	LEU
1	B	334	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	ASN
1	B	50	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	C1X	A	205	1	16,24,25	1.01	2 (12%)	11,30,32	1.24	1 (9%)
1	C1X	B	205	1	16,24,25	0.89	0	11,30,32	1.23	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	C1X	A	205	1	-	1/13/33/35	0/0/1/1
1	C1X	B	205	1	-	2/13/33/35	0/0/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	205	C1X	C5-C4	2.25	1.53	1.50
1	A	205	C1X	C3-C4	2.15	1.53	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	C1X	CE-NZ-C4	2.17	128.69	122.49
1	B	205	C1X	CE-NZ-C4	2.09	128.46	122.49

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	205	C1X	C5-C4-NZ-CE
1	B	205	C1X	C5-C4-NZ-CE
1	B	205	C1X	C1-C2-C3-C4

There are no ring outliers.

2 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	205	C1X	10	0
1	B	205	C1X	10	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	205:C1X	C	206:TYR	N	1.64

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	333/337 (98%)	-0.67	2 (0%) 89 91	10, 16, 26, 55	1 (0%)
1	B	328/337 (97%)	-0.56	7 (2%) 63 67	12, 16, 29, 52	0
All	All	661/674 (98%)	-0.62	9 (1%) 75 78	10, 16, 27, 55	1 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	336	GLY	13.4
1	B	336	GLY	7.3
1	B	227	GLY	6.3
1	B	144	ASP	3.5
1	A	335	ARG	3.4
1	B	335	ARG	3.0
1	B	221	ALA	2.9
1	B	334	ARG	2.8
1	B	228	ASN	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	C1X	A	205	24/25	0.97	0.13	11,14,18,22	0
1	C1X	B	205	24/25	0.97	0.10	10,15,23,26	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	A	1338	1/1	0.99	0.04	18,18,18,18	0
3	NA	A	1337	1/1	0.99	0.03	19,19,19,19	0
2	MG	A	1336	1/1	0.99	0.06	14,14,14,14	0
2	MG	B	1336	1/1	0.99	0.04	13,13,13,13	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.