



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2024 – 10:27 AM EDT

PDB ID : 9C4P
Title : Crystal structure of wild-type arabidopsis thaliana acetohydroxyacid synthase in complex with commercial herbicide triasulfuron
Authors : Cheng, Y.; Guddat, L.W.
Deposited on : 2024-06-05
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	590	 3% 93% 5%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 4884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetolactate synthase, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	582	4472	2837	773	837	25	0	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	668	LEU	-	expression tag	UNP P17597
A	669	GLU	-	expression tag	UNP P17597
A	670	HIS	-	expression tag	UNP P17597
A	671	HIS	-	expression tag	UNP P17597
A	672	HIS	-	expression tag	UNP P17597
A	673	HIS	-	expression tag	UNP P17597
A	674	HIS	-	expression tag	UNP P17597
A	675	HIS	-	expression tag	UNP P17597

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	53	27	9	15	2	0	0


- Molecule 4 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (three-letter code: NHE) (formula: $C_8H_{17}NO_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	13	8	1	3	1	0	0


- Molecule 5 is 2-(2-chloroethoxy)-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)carbamoyl]benzene-1-sulfonamide (three-letter code: A1AUE) (formula: $C_{14}H_{16}ClN_5O_5S$) (labeled as

"Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Cl	N	O			S
5	A	1	26	14	1	5	5	1	0	0

- Molecule 6 is S,R MESO-TARTARIC ACID (three-letter code: SRT) (formula: C₄H₆O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
6	A	1	10	4 6	0	0

- Molecule 7 is 2-[3-[(4-azanyl-2-methyl-pyrimidin-5-yl)methyl]-2-[(1 {S})-1-(dioxidanyl)-1-oxidanyl-ethyl]-4-methyl-1,3-thiazol-5-yl]ethyl phosphono hydrogen phosphate (three-letter code: AUJ) (formula: C₁₄H₂₃N₄O₁₀P₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
7	A	1	31	14	4	10	2	1	0	0

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
8	A	1	5	4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	262	Total	O	0	0
			262	262		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetolactate synthase, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	178.70Å 178.70Å 184.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.19 – 2.40 48.19 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.19-2.40) 99.9 (48.19-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 2.39Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.172 , 0.188 0.172 , 0.189	Depositor DCC
R_{free} test set	66251 reflections (2.92%)	wwPDB-VP
Wilson B-factor (Å ²)	53.7	Xtrriage
Anisotropy	0.269	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4884	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SRT, FAD, MG, NHE, AUJ, SO4, CSD, A1AUE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/4563	0.48	0/6193

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4472	0	4473	16	0
2	A	2	0	0	0	0
3	A	53	0	31	0	0
4	A	13	0	17	0	0
5	A	26	0	0	0	0
6	A	10	0	4	1	0
7	A	31	0	0	0	0
8	A	15	0	0	0	0
9	A	262	0	0	1	1
All	All	4884	0	4525	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:467:ALA:HA	1:A:621:LEU:HD21	1.88	0.56
1:A:316:GLU:OE2	1:A:416:LYS:NZ	2.41	0.51
1:A:103:LEU:HD21	1:A:192:ILE:HD13	1.94	0.50
1:A:526:ALA:HA	6:A:705:SRT:H2	1.95	0.49
1:A:114:THR:HG21	1:A:525:VAL:HG11	1.93	0.48
1:A:394:ILE:HG12	1:A:411:VAL:HB	1.95	0.48
1:A:91:PHE:CG	1:A:97:ARG:HG2	2.49	0.47
1:A:332:LEU:HD13	1:A:504:TRP:HE1	1.80	0.46
1:A:485:VAL:HG21	1:A:511:GLY:C	2.38	0.44
1:A:199[B]:ARG:NH2	9:A:809:HOH:O	2.49	0.44
1:A:92:ALA:HB3	1:A:95:GLN:HB2	2.01	0.43
1:A:644:GLN:O	1:A:665:ASP:HB2	2.20	0.42
1:A:562:LEU:HD21	1:A:601:MET:HG3	2.03	0.41
1:A:342:ASP:OD1	1:A:343:GLU:N	2.53	0.41
1:A:577:ARG:HG3	1:A:650:MET:HE3	2.03	0.41
1:A:316:GLU:HG2	1:A:420:GLN:HG3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:801:HOH:O	9:A:801:HOH:O[4_545]	2.12	0.08

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	581/590 (98%)	571 (98%)	10 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	479/486 (99%)	475 (99%)	4 (1%)	79 90

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	ARG
1	A	189	LEU
1	A	344	LEU
1	A	375	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSD	A	340	1	4,7,8	0.97	0	1,8,10	7.93	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	340	1	-	0/2/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	340	CSD	OD1-SG-CB	7.93	120.20	105.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	SO4	A	709	-	4,4,4	0.23	0	6,6,6	0.08	0
6	SRT	A	705	-	9,9,9	1.03	0	12,12,12	1.30	2 (16%)
8	SO4	A	708	-	4,4,4	0.24	0	6,6,6	0.08	0
8	SO4	A	710	-	4,4,4	0.25	0	6,6,6	0.06	0
7	AUJ	A	706	2	27,32,32	2.18	7 (25%)	31,49,49	1.69	10 (32%)
4	NHE	A	703	-	13,13,13	1.38	3 (23%)	16,17,17	1.80	5 (31%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FAD	A	702	-	54,58,58	2.99	21 (38%)	71,89,89	1.47	12 (16%)
5	A1AUE	A	704	-	27,27,27	1.94	8 (29%)	37,37,37	3.44	11 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SRT	A	705	-	-	10/12/12/12	-
7	AUJ	A	706	2	-	5/17/26/26	0/2/2/2
4	NHE	A	703	-	-	5/7/15/15	0/1/1/1
3	FAD	A	702	-	-	6/30/50/50	0/6/6/6
5	A1AUE	A	704	-	-	8/21/21/21	0/2/2/2

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	FAD	O4-C4	9.24	1.41	1.23
3	A	702	FAD	PA-O3P	8.34	1.68	1.59
3	A	702	FAD	P-O3P	8.14	1.68	1.59
3	A	702	FAD	O2-C2	7.82	1.39	1.24
3	A	702	FAD	O4B-C1B	-6.79	1.32	1.40
7	A	706	AUJ	C5-C4	5.24	1.51	1.42
7	A	706	AUJ	C2-N3	5.04	1.42	1.34
5	A	704	A1AUE	C22-N09	4.71	1.45	1.38
7	A	706	AUJ	C6-N1	4.06	1.42	1.34
5	A	704	A1AUE	C20-N09	3.64	1.45	1.37
3	A	702	FAD	C5B-C4B	-3.63	1.40	1.51
3	A	702	FAD	C6A-N6A	3.37	1.46	1.34
4	A	703	NHE	C2-S	3.36	1.82	1.77
5	A	704	A1AUE	C13-S02	3.34	1.82	1.77
7	A	706	AUJ	CBA-NBB	3.32	1.42	1.35
5	A	704	A1AUE	O05-S02	3.18	1.47	1.43
7	A	706	AUJ	CAP-NBB	-3.18	1.42	1.49
3	A	702	FAD	O4B-C4B	3.08	1.51	1.45
3	A	702	FAD	C4-N3	-3.07	1.33	1.38
3	A	702	FAD	O3B-C3B	3.02	1.50	1.43
7	A	706	AUJ	C4-NAD	2.99	1.41	1.34
3	A	702	FAD	C9A-N10	-2.99	1.36	1.41
7	A	706	AUJ	C4-N3	-2.98	1.31	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	704	A1AUE	O07-C24	2.96	1.40	1.33
3	A	702	FAD	C3B-C4B	2.83	1.60	1.53
5	A	704	A1AUE	O04-S02	2.76	1.46	1.43
3	A	702	FAD	C1'-C2'	2.76	1.56	1.52
3	A	702	FAD	C1B-N9A	-2.68	1.43	1.49
5	A	704	A1AUE	S02-N08	2.59	1.70	1.64
5	A	704	A1AUE	C20-N08	2.46	1.44	1.39
3	A	702	FAD	C2-N3	-2.44	1.33	1.39
3	A	702	FAD	PA-O5B	2.40	1.68	1.59
3	A	702	FAD	C4X-N5	2.30	1.35	1.30
4	A	703	NHE	O2-S	2.25	1.51	1.45
3	A	702	FAD	P-O5'	2.24	1.68	1.59
3	A	702	FAD	O4'-C4'	-2.23	1.38	1.43
3	A	702	FAD	C5'-C4'	2.19	1.54	1.51
4	A	703	NHE	O1-S	2.12	1.51	1.45
3	A	702	FAD	O2B-C2B	2.09	1.48	1.43

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	704	A1AUE	O05-S02-O04	-14.52	101.88	119.52
5	A	704	A1AUE	C24-N11-C22	8.10	119.47	112.86
5	A	704	A1AUE	C22-N09-C20	-6.01	124.15	130.34
3	A	702	FAD	N3A-C2A-N1A	-5.81	120.78	128.67
5	A	704	A1AUE	C13-S02-N08	5.46	112.75	105.96
5	A	704	A1AUE	N11-C24-N12	-4.13	120.14	127.65
4	A	703	NHE	O2-S-O1	-3.77	101.57	113.82
5	A	704	A1AUE	C22-N10-C23	3.68	120.10	114.61
5	A	704	A1AUE	N11-C22-N10	-3.58	120.39	126.26
7	A	706	AUJ	C6-N1-C2	3.45	121.75	116.07
5	A	704	A1AUE	N10-C23-N12	-3.38	119.81	125.77
4	A	703	NHE	O2-S-C2	3.16	111.50	106.73
3	A	702	FAD	O2'-C2'-C3'	3.12	116.55	109.25
3	A	702	FAD	C4-C4X-N5	3.10	122.49	118.21
3	A	702	FAD	C5'-C4'-C3'	-2.96	106.64	112.22
7	A	706	AUJ	CAZ-CAX-NBB	2.95	113.84	107.66
4	A	703	NHE	O1-S-C2	2.93	111.16	106.73
7	A	706	AUJ	N1-C2-N3	-2.90	120.70	125.53
3	A	702	FAD	C4X-C10-N10	2.88	120.60	116.48
3	A	702	FAD	C4-N3-C2	-2.81	120.64	125.64
3	A	702	FAD	C4A-C5A-N7A	-2.77	106.41	109.34
3	A	702	FAD	C1'-C2'-C3'	2.60	116.70	109.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	706	AUJ	C5-C6-N1	-2.59	119.62	123.83
6	A	705	SRT	O1-C1-C2	2.52	120.32	113.31
3	A	702	FAD	C4X-C4-N3	2.47	119.54	113.25
7	A	706	AUJ	CAB-CAX-CAZ	-2.47	122.21	127.60
7	A	706	AUJ	OAK-PBE-OAG	-2.42	101.19	112.44
7	A	706	AUJ	CAA-C2-N1	2.37	119.72	117.20
5	A	704	A1AUE	C25-C23-N12	2.36	120.66	117.13
7	A	706	AUJ	OAI-PBD-OAF	-2.34	101.74	110.83
6	A	705	SRT	O41-C4-C3	2.32	119.75	113.31
3	A	702	FAD	C10-C4X-N5	-2.28	120.14	124.81
4	A	703	NHE	O3-S-C2	2.28	110.47	106.00
5	A	704	A1AUE	C26-O07-C24	-2.28	113.97	117.56
3	A	702	FAD	O4-C4-C4X	-2.25	120.60	126.53
4	A	703	NHE	C1-N-C1'	-2.14	110.07	114.18
7	A	706	AUJ	NAD-C4-N3	2.12	119.89	117.03
5	A	704	A1AUE	O03-C14-C13	2.08	119.92	116.70
7	A	706	AUJ	OAK-PBE-OAT	2.06	112.84	107.27
3	A	702	FAD	C9A-N10-C10	-2.03	117.65	120.75

There are no chirality outliers.

All (34) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	FAD	C3'-C4'-C5'-O5'
3	A	702	FAD	O4'-C4'-C5'-O5'
4	A	703	NHE	N-C1-C2-S
4	A	703	NHE	C1-C2-S-O1
4	A	703	NHE	C1-C2-S-O3
5	A	704	A1AUE	C14-C13-S02-O05
6	A	705	SRT	C2-C3-C4-O41
7	A	706	AUJ	CAN-CAO-CAZ-CAX
7	A	706	AUJ	CAN-OAS-PBE-OAG
7	A	706	AUJ	PBE-OAT-PBD-OAI
7	A	706	AUJ	CAC-CBC-OBC1-OC11
6	A	705	SRT	O1-C1-C2-O2
6	A	705	SRT	O11-C1-C2-O2
6	A	705	SRT	O1-C1-C2-C3
6	A	705	SRT	O11-C1-C2-C3
6	A	705	SRT	C2-C3-C4-O4
6	A	705	SRT	O3-C3-C4-O4
6	A	705	SRT	O3-C3-C4-O41
5	A	704	A1AUE	C15-C13-S02-O05

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	A	705	SRT	C1-C2-C3-O3
3	A	702	FAD	O2'-C2'-C3'-O3'
5	A	704	A1AUE	C14-C13-S02-N08
6	A	705	SRT	C1-C2-C3-C4
3	A	702	FAD	O2'-C2'-C3'-C4'
4	A	703	NHE	C2-C1-N-C1'
5	A	704	A1AUE	C15-C13-S02-N08
3	A	702	FAD	PA-O3P-P-O5'
4	A	703	NHE	C1-C2-S-O2
7	A	706	AUJ	OAS-CAN-CAO-CAZ
5	A	704	A1AUE	O03-C19-C21-CL01
5	A	704	A1AUE	C20-N08-S02-O04
3	A	702	FAD	C4B-C5B-O5B-PA
5	A	704	A1AUE	C21-C19-O03-C14
5	A	704	A1AUE	N08-C20-N09-C22



There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	705	SRT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	581/590 (98%)	-0.25	16 (2%) 55 51	26, 53, 80, 104	2 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	431	GLU	4.9
1	A	86	THR	4.1
1	A	434	LYS	3.4
1	A	408	HIS	3.4
1	A	341	ASP	2.8
1	A	449	GLN	2.6
1	A	287	SER	2.6
1	A	279	ARG	2.5
1	A	269	GLN	2.4
1	A	617	LYS	2.3
1	A	300	LYS	2.2
1	A	343	GLU	2.1
1	A	442	ASN	2.1
1	A	430	ALA	2.0
1	A	388	ARG	2.0
1	A	469	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	340	8/9	0.93	0.15	66,76,86,89	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	SO4	A	710	5/5	0.64	0.32	79,80,85,112	5
6	SRT	A	705	10/10	0.74	0.23	71,77,83,90	10
8	SO4	A	709	5/5	0.91	0.15	64,69,88,91	5
8	SO4	A	708	5/5	0.92	0.17	62,76,82,90	5
4	NHE	A	703	13/13	0.96	0.11	49,60,70,74	0
5	A1AUE	A	704	26/26	0.98	0.07	41,49,55,61	7
3	FAD	A	702	53/53	0.98	0.06	37,49,54,60	0
7	AUJ	A	706	31/31	0.98	0.09	36,47,61,72	20
2	MG	A	701	1/1	0.99	0.03	46,46,46,46	0
2	MG	A	707	1/1	0.99	0.17	74,74,74,74	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SRT A 705:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NHE A 703:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around A1AUE A 704:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.