



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 08:54 am BST

PDB ID : 4C8O  
Title : Binary complex of the large fragment of DNA polymerase I from *Thermus Aquaticus* with the artificial base pair dNaM-d5SICS at the postinsertion site (sequence context 2)  
Authors : Betz, K.; Malyshev, D.A.; Lavergne, T.; Welte, W.; Diederichs, K.; Romesberg, F.E.; Marx, A.  
Deposited on : 2013-10-01  
Resolution : 1.75 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	
2	B	11	
3	C	14	

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9712 atoms, of which 4644 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE I, THERMOSTABLE.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S				
1	A	540	8680	2742	4365	772	788	13	0	5	0	

- Molecule 2 is a DNA chain called 5'-D(\*GP\*CP\*CP\*AP\*CP\*GP\*GP\*CP\*GP\*CP\*LHOP)-3'.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	S			
2	B	11	352	110	128	41	62	10	1	0	0	0

- Molecule 3 is a DNA chain called 5'-D(\*TP\*TP\*CP\*BMNP\*GP\*CP\*GP\*CP\*CP\*GP\*TP\*GP\*GP\*CP)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
3	C	12	387	121	139	42	73	12	0	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
7	C	1	20	4	12	1	3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	220	Total	O	0	0
			220	220		
8	B	10	Total	O	0	0
			10	10		
8	C	6	Total	O	0	0
			6	6		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA POLYMERASE I, THERMOSTABLE



- Molecule 2: 5'-D(\*GP\*CP\*CP\*AP\*CP\*GP\*GP\*CP\*GP\*CP\*LHOP)-3'



- Molecule 3: 5'-D(\*TP\*TP\*CP\*BMNP\*GP\*CP\*GP\*CP\*CP\*GP\*TP\*GP\*GP\*CP)-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.80Å 99.19Å 203.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.88 – 1.75 47.88 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.88-1.75) 99.6 (47.88-1.75)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 1.75Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.185 , 0.212 0.188 , 0.214	Depositor DCC
$R_{free}$ test set	3300 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.5	Xtrriage
Anisotropy	0.535	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 47.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9712	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: BMN, MG, SO4, CL, TRS, LHO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.28	0/4420	0.48	0/5988
2	B	0.52	0/225	0.78	0/345
3	C	0.84	2/250 (0.8%)	1.15	4/381 (1.0%)
All	All	0.35	2/4895 (0.0%)	0.56	4/6714 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	210	DC	C4'-O4'	-7.07	1.38	1.45
3	C	210	DC	C2'-C1'	6.57	1.58	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	210	DC	O4'-C1'-C2'	9.70	113.66	105.90
3	C	210	DC	C3'-C2'-C1'	-9.04	91.65	102.50
3	C	210	DC	O4'-C4'-C3'	-5.75	102.20	104.50
3	C	210	DC	C4'-C3'-C2'	5.62	108.15	103.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4315	4365	4369	12	0
2	B	224	128	127	2	0
3	C	248	139	139	4	0
4	A	35	0	0	1	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	C	8	12	12	0	0
8	A	220	0	0	3	0
8	B	10	0	0	0	0
8	C	6	0	0	0	0
All	All	5068	4644	4647	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:759:ASP:OD2	8:A:2193:HOH:O	2.16	0.63
2:B:112:LHO:H210	3:C:204:DC:H1'	1.82	0.62
1:A:686:TYR:OH	4:A:1839:SO4:O1	2.24	0.55
1:A:746:ARG:NE	8:A:2173:HOH:O	2.38	0.54
3:C:210:DC:H2'	3:C:211:DG:C8	2.45	0.52
1:A:611:TYR:HB3	1:A:614:ILE:HB	1.93	0.49
1:A:549:LEU:N	8:A:2120:HOH:O	2.33	0.47
1:A:524:GLU:N	1:A:524:GLU:OE1	2.47	0.47
1:A:299:TRP:CG	1:A:300:PRO:HA	2.49	0.47
1:A:547:ASP:HB2	1:A:548:PRO:HD3	2.00	0.44
2:B:104:DC:H2'	2:B:105:DA:C8	2.53	0.42
1:A:302:PRO:HG2	1:A:328:ARG:HD3	2.02	0.42
1:A:437:LEU:HD21	1:A:574:LEU:HD21	2.02	0.41
1:A:466:GLU:OE1	1:A:469:ARG:NH1	2.51	0.41
3:C:209:DC:H2'	3:C:210:DC:C6	2.56	0.41
1:A:535:TYR:CZ	1:A:539[A]:THR:HG21	2.57	0.40
3:C:204:DC:C6	3:C:205:BMN:H23	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	543/540 (101%)	532 (98%)	10 (2%)	1 (0%)	47 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	503	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	446/441 (101%)	440 (99%)	6 (1%)	69 54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	354	LYS
1	A	411	ARG
1	A	484	LEU
1	A	487	ARG
1	A	636	ARG
1	A	805	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LHO	B	112	2	20,25,26	0.16	0	27,36,39	0.43	0
3	BMN	C	205	3	22,25,26	0.16	0	31,35,38	0.89	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LHO	B	112	2	-	0/7/21/22	0/3/3/3
3	BMN	C	205	3	-	1/9/23/24	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	205	BMN	C2-C1-C6	3.44	121.94	117.06

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	205	BMN	C2-C1-C1'-C2'

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	112	LHO	1	0
3	C	205	BMN	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	A	1833	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	A	1837	-	4,4,4	0.15	0	6,6,6	0.08	0
4	SO4	A	1834	-	4,4,4	0.14	0	6,6,6	0.09	0
4	SO4	A	1839	-	4,4,4	0.13	0	6,6,6	0.08	0
4	SO4	A	1840	-	4,4,4	0.13	0	6,6,6	0.06	0
4	SO4	A	1841	-	4,4,4	0.13	0	6,6,6	0.05	0
7	TRS	C	1216	-	7,7,7	0.36	0	9,9,9	0.29	0
4	SO4	A	1838	-	4,4,4	0.15	0	6,6,6	0.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	TRS	C	1216	-	-	0/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1839	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	540/540 (100%)	0.56	43 (7%) <span style="border: 1px solid red; padding: 2px;">12</span> <span style="border: 1px solid red; padding: 2px;">16</span>	33, 47, 83, 131	0
2	B	10/11 (90%)	0.16	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	51, 57, 67, 71	0
3	C	11/14 (78%)	0.25	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	44, 58, 99, 132	0
All	All	561/565 (99%)	0.55	43 (7%) <span style="border: 1px solid red; padding: 2px;">13</span> <span style="border: 1px solid red; padding: 2px;">18</span>	33, 47, 83, 132	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	636	ARG	6.4
1	A	635	GLY	5.9
1	A	736	ARG	5.7
1	A	735	ALA	5.3
1	A	737	VAL	5.2
1	A	525	ALA	5.0
1	A	740	VAL	4.7
1	A	587	ARG	4.3
1	A	343	ARG	4.1
1	A	738	LYS	4.1
1	A	832	GLU	4.1
1	A	734	GLU	3.9
1	A	519	LEU	3.9
1	A	477	LEU	3.5
1	A	293	ALA	3.5
1	A	508	LYS	3.3
1	A	581	LEU	3.3
1	A	509	THR	3.3
1	A	564	PHE	3.0
1	A	507	GLU	2.9
1	A	634	GLU	2.7
1	A	552	LEU	2.7
1	A	506	THR	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	294	LEU	2.5
1	A	783	VAL	2.5
1	A	577	SER	2.4
1	A	562	THR	2.4
1	A	549	LEU	2.4
1	A	570	ALA	2.4
1	A	579	PRO	2.3
1	A	594	ILE	2.3
1	A	590	LEU	2.3
1	A	365	LEU	2.2
1	A	511	LYS	2.2
1	A	524	GLU	2.2
1	A	529	VAL	2.2
1	A	739	SER	2.2
1	A	475	PHE	2.2
1	A	584	ILE	2.2
1	A	492	ARG	2.2
1	A	667	PHE	2.1
1	A	533	LEU	2.1
1	A	503	ILE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BMN	C	205	23/24	0.76	0.16	97,135,174,190	0
2	LHO	B	112	23/24	0.95	0.13	46,62,87,96	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	TRS	C	1216	8/8	0.75	0.28	69,91,112,118	0
4	SO4	A	1841	5/5	0.88	0.17	79,81,97,107	5
4	SO4	A	1837	5/5	0.89	0.28	35,52,56,57	5
4	SO4	A	1838	5/5	0.89	0.23	90,90,100,100	5
4	SO4	A	1834	5/5	0.90	0.17	58,59,65,68	5
4	SO4	A	1840	5/5	0.91	0.12	73,82,95,97	5
4	SO4	A	1839	5/5	0.92	0.17	61,64,69,71	5
6	CL	A	1836	1/1	0.93	0.14	74,74,74,74	0
4	SO4	A	1833	5/5	0.94	0.49	112,112,112,116	5
5	MG	A	1835	1/1	0.98	0.09	66,66,66,66	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.