



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 4, 2023 – 08:48 pm GMT

PDB ID : 2CAR
Title : Crystal Structure Of Human Inosine Triphosphatase
Authors : Stenmark, P.; Kursula, P.; Arrowsmith, C.; Berglund, H.; Edwards, A.; Ehn, M.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Hallberg, B.M.; Hogbom, M.; Holmberg Schiavone, L.; Kotenyova, T.; Nilsson-Ehle, P.; Nyman, T.; Ogg, D.; Persson, C.; Sagemark, J.; Schuler, H.; Sundstrom, M.; Thorsell, A.G.; van den Berg, S.; Weigelt, J.; Nordlund, P.
Deposited on : 2005-12-22
Resolution : 1.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Rfmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	196	 76% 18% . .
1	B	196	 77% 20% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INOSINE TRIPHOSPHATE PYROPHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	196	1646	1053	275	306	12	0	20	0
1	B	194	1611	1032	269	301	9	0	15	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	319	Total 319	O 319	0	0
2	B	329	Total 329	O 329	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INOSINE TRIPHOSPHATE PYROPHOSPHATASE



- Molecule 1: INOSINE TRIPHOSPHATE PYROPHOSPHATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	31.23Å 105.00Å 50.14Å 90.00° 90.01° 90.00°	Depositor
Resolution (Å)	40.00 – 1.09 36.26 – 1.09	Depositor EDS
% Data completeness (in resolution range)	90.7 (40.00-1.09) 91.8 (36.26-1.09)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 1.09Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.097 , 0.153 0.102 , 0.155	Depositor DCC
R_{free} test set	2636 reflections (2.13%)	wwPDB-VP
Wilson B-factor (Å ²)	7.7	Xtrriage
Anisotropy	0.174	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 58.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.488 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	3905	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	0/1721	1.92	43/2323 (1.9%)
1	B	0.81	0/1679	1.92	47/2269 (2.1%)
All	All	0.81	0/3400	1.92	90/4592 (2.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (90) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	ASP	CB-CG-OD2	21.28	137.45	118.30
1	B	133[A]	ARG	NE-CZ-NH2	18.30	129.45	120.30
1	B	133[B]	ARG	NE-CZ-NH2	18.30	129.45	120.30
1	A	144	ARG	NE-CZ-NH2	16.82	128.71	120.30
1	A	124	ASP	CB-CG-OD1	-15.51	104.34	118.30
1	A	178[A]	ARG	NE-CZ-NH2	-14.49	113.06	120.30
1	A	178[B]	ARG	NE-CZ-NH2	-14.49	113.06	120.30
1	A	144	ARG	NE-CZ-NH1	-13.39	113.61	120.30
1	A	161[A]	GLU	OE1-CD-OE2	-12.97	107.73	123.30
1	A	161[B]	GLU	OE1-CD-OE2	-12.97	107.73	123.30
1	B	161[A]	GLU	CG-CD-OE1	12.31	142.93	118.30
1	B	161[B]	GLU	CG-CD-OE1	12.31	142.93	118.30
1	B	135	ARG	NE-CZ-NH1	-12.12	114.24	120.30
1	A	135	ARG	NE-CZ-NH1	11.72	126.16	120.30
1	A	124	ASP	CA-CB-CG	-11.44	88.23	113.40
1	A	124	ASP	CB-CA-C	11.25	132.90	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	164	TYR	CB-CG-CD1	10.55	127.33	121.00
1	A	125	PRO	CA-C-N	10.41	140.11	117.20
1	B	144[A]	ARG	NE-CZ-NH2	-10.38	115.11	120.30
1	B	144[B]	ARG	NE-CZ-NH2	-10.38	115.11	120.30
1	B	48[A]	GLU	OE1-CD-OE2	10.37	135.74	123.30
1	B	48[B]	GLU	OE1-CD-OE2	10.37	135.74	123.30
1	A	133[A]	ARG	NE-CZ-NH2	10.24	125.42	120.30
1	A	133[B]	ARG	NE-CZ-NH2	10.24	125.42	120.30
1	A	164	TYR	CB-CG-CD2	-10.23	114.86	121.00
1	A	125	PRO	C-N-CA	10.07	146.88	121.70
1	B	161[A]	GLU	OE1-CD-OE2	-9.95	111.36	123.30
1	B	161[B]	GLU	OE1-CD-OE2	-9.95	111.36	123.30
1	B	130[A]	ARG	CD-NE-CZ	9.94	137.51	123.60
1	B	130[B]	ARG	CD-NE-CZ	9.94	137.51	123.60
1	B	178	ARG	CD-NE-CZ	9.90	137.47	123.60
1	B	144[A]	ARG	NE-CZ-NH1	9.66	125.13	120.30
1	B	144[B]	ARG	NE-CZ-NH1	9.66	125.13	120.30
1	B	178	ARG	NE-CZ-NH2	-9.62	115.49	120.30
1	A	139	ARG	NE-CZ-NH1	8.98	124.79	120.30
1	B	161[A]	GLU	CB-CG-CD	8.47	137.07	114.20
1	B	161[B]	GLU	CB-CG-CD	8.47	137.07	114.20
1	B	158[A]	ASP	CA-CB-CG	8.17	131.37	113.40
1	B	158[B]	ASP	CA-CB-CG	8.17	131.37	113.40
1	A	161[A]	GLU	CG-CD-OE1	8.07	134.45	118.30
1	A	161[B]	GLU	CG-CD-OE1	8.07	134.45	118.30
1	B	62	ARG	NE-CZ-NH2	7.88	124.24	120.30
1	B	50	ASP	CB-CG-OD1	7.88	125.39	118.30
1	B	77	PHE	CB-CG-CD1	7.34	125.94	120.80
1	B	135	ARG	NH1-CZ-NH2	7.26	127.38	119.40
1	B	122	THR	CA-CB-CG2	7.13	122.38	112.40
1	A	160	TYR	CD1-CE1-CZ	6.83	125.95	119.80
1	B	87	TYR	CB-CG-CD2	-6.76	116.94	121.00
1	A	109	ASP	CB-CG-OD1	6.73	124.35	118.30
1	A	125	PRO	CA-C-O	-6.71	104.10	120.20
1	B	158[A]	ASP	CB-CG-OD1	6.70	124.33	118.30
1	B	158[B]	ASP	CB-CG-OD1	6.70	124.33	118.30
1	B	144[A]	ARG	CD-NE-CZ	6.69	132.96	123.60
1	B	144[B]	ARG	CD-NE-CZ	6.69	132.96	123.60
1	A	133[A]	ARG	NE-CZ-NH1	-6.51	117.04	120.30
1	A	133[B]	ARG	NE-CZ-NH1	-6.51	117.04	120.30
1	A	93	GLU	OE1-CD-OE2	-6.41	115.61	123.30
1	A	126	SER	C-N-CA	6.39	137.69	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178[A]	ARG	NE-CZ-NH1	6.39	123.50	120.30
1	A	178[B]	ARG	NE-CZ-NH1	6.39	123.50	120.30
1	B	161[A]	GLU	CG-CD-OE2	-6.32	105.66	118.30
1	B	161[B]	GLU	CG-CD-OE2	-6.32	105.66	118.30
1	B	133[A]	ARG	NH1-CZ-NH2	-6.12	112.67	119.40
1	B	133[B]	ARG	NH1-CZ-NH2	-6.12	112.67	119.40
1	B	139	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	B	130[A]	ARG	NE-CZ-NH2	-6.07	117.26	120.30
1	B	130[B]	ARG	NE-CZ-NH2	-6.07	117.26	120.30
1	B	164	TYR	CB-CG-CD1	5.94	124.56	121.00
1	B	50	ASP	CB-CG-OD2	-5.88	113.00	118.30
1	B	62	ARG	NH1-CZ-NH2	-5.85	112.97	119.40
1	A	0	SER	CA-C-O	5.74	132.16	120.10
1	A	77	PHE	CB-CG-CD1	5.65	124.75	120.80
1	A	2	ALA	N-CA-CB	-5.51	102.39	110.10
1	A	125	PRO	O-C-N	-5.32	114.19	122.70
1	B	162	GLN	CA-CB-CG	5.24	124.94	113.40
1	B	41	ASP	CA-CB-CG	5.22	124.88	113.40
1	A	133[A]	ARG	CD-NE-CZ	5.21	130.90	123.60
1	A	133[B]	ARG	CD-NE-CZ	5.21	130.90	123.60
1	A	175	VAL	CG1-CB-CG2	5.21	119.24	110.90
1	B	68	VAL	CG1-CB-CG2	5.21	119.23	110.90
1	A	135	ARG	NH1-CZ-NH2	-5.20	113.68	119.40
1	A	62	ARG	CD-NE-CZ	5.18	130.85	123.60
1	A	29[A]	ASP	CB-CG-OD2	5.14	122.93	118.30
1	A	29[B]	ASP	CB-CG-OD2	5.14	122.93	118.30
1	A	124	ASP	N-CA-CB	-5.14	101.35	110.60
1	B	87	TYR	CB-CG-CD1	5.08	124.05	121.00
1	B	187	GLU	CB-CG-CD	5.08	127.90	114.20
1	A	0	SER	O-C-N	-5.07	114.59	122.70
1	B	183	LEU	CB-CG-CD2	-5.04	102.43	111.00
1	A	91	PHE	CB-CG-CD1	-5.01	117.29	120.80

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	124	ASP	Peptide
1	A	125	PRO	Peptide,Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1646	0	1661	30	0
1	B	1611	0	1622	20	0
2	A	319	0	0	7	0
2	B	329	0	0	11	0
All	All	3905	0	3283	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71[B]:GLU:OE1	1:A:178[B]:ARG:NH1	1.76	1.15
1:B:51:GLU:O	1:B:55:GLN:HG3	1.82	0.80
1:A:52:ILE:HG23	2:A:2128:HOH:O	1.86	0.75
1:B:22[B]:GLU:HB2	2:B:2073:HOH:O	1.86	0.74
1:B:22[A]:GLU:HB2	2:B:2073:HOH:O	1.87	0.74
1:B:40:ILE:HG23	2:B:2172:HOH:O	1.98	0.64
1:A:124:ASP:HB3	1:A:125:PRO:O	1.98	0.63
1:A:127:GLN:OE1	1:A:128:PRO:HD2	2.03	0.57
1:A:45:TYR:HD2	2:A:2128:HOH:O	1.87	0.57
1:B:56:LYS:HE3	2:B:2126:HOH:O	2.04	0.57
1:B:3:ALA:HB3	2:B:2009:HOH:O	2.05	0.56
1:A:124:ASP:CG	1:A:127:GLN:H	2.10	0.54
1:A:71[B]:GLU:OE1	1:A:178[B]:ARG:CZ	2.53	0.53
1:A:143:PRO:HA	1:A:153[B]:PRO:O	2.09	0.53
1:B:38:GLN:HG3	2:B:2044:HOH:O	2.09	0.52
1:B:16:ASN:OD1	1:B:18:LYS:HB3	2.10	0.52
1:A:125:PRO:HG3	2:A:2216:HOH:O	2.08	0.52
1:A:162[A]:GLN:HG3	2:A:2261:HOH:O	2.11	0.49
1:B:93:GLU:HG2	2:B:2097:HOH:O	2.12	0.49
1:B:1:MET:HB3	1:B:188:TYR:OH	2.13	0.48
1:A:1[B]:MET:CE	1:A:130[B]:ARG:HG3	2.44	0.48
1:A:97:PRO:HB3	1:A:154[B]:CYS:SG	2.54	0.47
1:A:98[B]:GLU:OE2	1:A:147:GLN:NE2	2.46	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:PRO:HB2	1:A:126:SER:OG	2.15	0.46
1:B:63:GLN:NE2	2:B:2172:HOH:O	2.48	0.46
1:B:62:ARG:NH1	2:B:2167:HOH:O	2.49	0.46
1:B:4:SER:N	2:B:2009:HOH:O	2.48	0.45
1:B:130[B]:ARG:HG2	1:B:132:PHE:CZ	2.52	0.45
1:A:153[A]:PRO:O	1:A:163:THR:HG21	2.17	0.45
1:A:124:ASP:OD1	1:A:127:GLN:N	2.45	0.44
1:B:59:GLU:OE2	1:B:62:ARG:NH2	2.49	0.44
1:A:149:PHE:O	1:A:152:ASP:HB2	2.17	0.44
1:A:67:PRO:HA	1:A:121:SER:O	2.18	0.43
1:A:178[A]:ARG:NH2	2:A:2290:HOH:O	2.45	0.43
1:B:130[B]:ARG:HD3	1:B:132:PHE:CZ	2.54	0.43
1:A:4:SER:N	2:A:2001:HOH:O	2.50	0.43
1:A:124:ASP:OD2	1:A:127:GLN:O	2.36	0.43
1:A:163:THR:O	1:A:167[A]:MET:HG3	2.19	0.43
1:B:8:LYS:HD3	2:B:2124:HOH:O	2.18	0.42
1:A:125:PRO:HB2	1:A:126:SER:CB	2.49	0.42
1:B:168:PRO:HG2	1:B:171:GLU:OE2	2.20	0.42
1:A:50:ASP:OD1	1:A:133[B]:ARG:NH2	2.50	0.42
1:B:42:LEU:HD11	1:B:60:ALA:HB2	2.00	0.42
1:A:66:GLY:O	1:A:68[B]:VAL:HG23	2.20	0.41
1:A:5:LEU:HA	1:A:8:LYS:HG3	2.03	0.41
1:A:1[A]:MET:SD	1:A:122:THR:HG23	2.60	0.41
1:A:124:ASP:OD1	1:A:127:GLN:HB3	2.21	0.41
1:A:139:ARG:NH2	2:A:2234:HOH:O	2.53	0.40
1:A:163:THR:O	1:A:167[B]:MET:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/196 (109%)	208 (97%)	3 (1%)	3 (1%)	11	1
1	B	207/196 (106%)	206 (100%)	0	1 (0%)	29	6
All	All	421/392 (107%)	414 (98%)	3 (1%)	4 (1%)	15	1

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	124	ASP
1	A	126	SER
1	A	125	PRO
1	B	96	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/161 (112%)	171 (94%)	10 (6%)	21	2
1	B	175/161 (109%)	171 (98%)	4 (2%)	50	12
All	All	356/322 (111%)	342 (96%)	14 (4%)	37	3

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1[A]	MET
1	A	1[B]	MET
1	A	97	PRO
1	A	98[A]	GLU
1	A	98[B]	GLU
1	A	124	ASP
1	A	139	ARG
1	A	162[A]	GLN
1	A	162[B]	GLN
1	A	164	TYR
1	B	1	MET
1	B	16	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	126	SER
1	B	164	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	B	63	GLN
1	B	162	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/196 (100%)	-0.75	1 (0%) 91 89	5, 10, 24, 39	0
1	B	194/196 (98%)	-0.81	0 100 100	6, 10, 24, 39	0
All	All	390/392 (99%)	-0.78	1 (0%) 94 91	5, 10, 24, 39	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	125	PRO	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.