



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 28, 2024 – 02:00 am BST

PDB ID : 8CB7  
Title : The Transcriptional Regulator PrfA from *Listeria Monocytogenes* in complex with tetrapeptide Glu-Val-Phe-Leu  
Authors : Oelker, M.; Sauer-Eriksson, A.E.  
Deposited on : 2023-01-25  
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	239	
1	B	239	
2	C	4	
2	D	4	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7591 atoms, of which 3796 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Listeriolysin regulatory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	229	3742	1221	1873	292	349	7	0	0	0
1	B	226	3706	1210	1855	289	345	7	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P22262
A	0	ALA	-	expression tag	UNP P22262
B	-1	GLY	-	expression tag	UNP P22262
B	0	ALA	-	expression tag	UNP P22262

- Molecule 2 is a protein called tetrapeptide GLU-VAL-PHE-LEU.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	C	4	70	25	34	4	7	0	0	0
2	D	4	70	25	34	4	7	0	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total 3 3	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS failed to run properly.


- Molecule 1: Listeriolysin regulatory protein

Chain A: 




- Molecule 1: Listeriolysin regulatory protein

Chain B: 



- Molecule 2: tetrapeptide GLU-VAL-PHE-LEU

Chain C: 



- Molecule 2: tetrapeptide GLU-VAL-PHE-LEU

Chain D: 



## 4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.62Å 94.24Å 101.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.70 – 2.85	Depositor
% Data completeness (in resolution range)	100.0 (47.70-2.85)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.39 (at 2.86Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.259 , 0.307	Depositor
Wilson B-factor (Å <sup>2</sup> )	68.2	Xtrriage
Anisotropy	0.051	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7591	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/1910	0.65	0/2577
1	B	0.63	0/1892	0.74	0/2553
2	C	1.09	0/36	1.00	0/46
2	D	0.27	0/36	0.41	0/46
All	All	0.52	0/3874	0.70	0/5222

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1869	1873	1874	30	0
1	B	1851	1855	1855	23	0
2	C	36	34	34	1	0
2	D	36	34	34	2	0
3	A	3	0	0	0	0
All	All	3795	3796	3797	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:165:THR:O	1:B:165:THR:HG22	1.94	0.67
1:A:198:VAL:HG11	1:A:213:TYR:CD1	2.31	0.66
1:A:151:THR:HG21	1:A:208:VAL:HG21	1.79	0.64
1:A:193:LEU:O	1:A:198:VAL:HG22	1.98	0.63
1:A:143:ILE:HD11	1:A:186:VAL:CG1	2.31	0.61
1:B:209:GLN:O	1:B:209:GLN:HG3	1.99	0.61
1:B:68:VAL:HG21	1:B:116:VAL:HG22	1.82	0.60
1:A:150:LEU:HD11	2:C:303:LEU:HD11	1.84	0.59
1:B:125:SER:HG	1:B:224:TRP:HE1	1.51	0.58
1:B:96:VAL:O	1:B:96:VAL:HG23	2.05	0.57
1:A:197:LYS:O	1:A:197:LYS:HG2	2.05	0.56
1:A:113:PHE:CD1	1:B:113:PHE:CZ	2.94	0.56
1:B:171:MET:HA	1:B:171:MET:CE	2.35	0.54
1:A:143:ILE:HD11	1:A:186:VAL:HG11	1.90	0.54
1:A:147:LEU:O	1:A:151:THR:HG23	2.07	0.54
1:B:149:ILE:HD11	2:D:301:VAL:O	2.07	0.54
1:B:163:LYS:HE3	1:B:207:TYR:OH	2.07	0.53
1:A:137:ASN:ND2	1:A:217:TYR:O	2.42	0.52
1:A:146:GLN:HA	1:A:146:GLN:OE1	2.10	0.51
1:A:20:LYS:HE2	1:A:96:VAL:HG11	1.92	0.51
1:A:140:LEU:HD13	1:A:193:LEU:HD21	1.93	0.51
1:B:24:LYS:HB2	1:B:89:SER:O	2.11	0.50
1:A:143:ILE:HD11	1:A:186:VAL:HG13	1.94	0.48
1:B:54:ASN:OD1	1:B:55:GLY:N	2.47	0.47
1:B:148:LEU:CD1	1:B:211:LEU:HD11	2.44	0.47
1:A:151:THR:CG2	1:A:208:VAL:HG21	2.45	0.47
1:B:78:THR:HG22	1:B:79:SER:N	2.28	0.47
1:B:126:TYR:OH	2:D:303:LEU:N	2.48	0.46
1:A:76:THR:O	1:A:77:GLU:HB2	2.15	0.46
1:A:143:ILE:CD1	1:A:186:VAL:HG13	2.46	0.46
1:A:113:PHE:CE1	1:B:113:PHE:CZ	3.04	0.46
1:B:68:VAL:HG11	1:B:116:VAL:HG13	1.97	0.46
1:A:74:ILE:HA	1:A:99:ILE:HD11	1.97	0.46
1:A:108:LYS:HA	1:A:108:LYS:HD3	1.69	0.46
1:B:148:LEU:HD12	1:B:211:LEU:HD11	1.98	0.46
1:B:149:ILE:O	1:B:153:VAL:HG12	2.16	0.45
1:B:102:LEU:HG	1:B:106:LEU:HD22	1.99	0.44
1:A:106:LEU:HD22	1:A:113:PHE:HA	1.99	0.43
1:A:67:PHE:CE1	1:A:119:THR:HG23	2.54	0.43
1:A:110:LEU:HD21	1:A:114:PHE:CZ	2.53	0.43
1:A:18:LYS:HG2	1:A:19:PRO:HD2	2.01	0.43
1:A:143:ILE:CD1	1:A:186:VAL:CG1	2.96	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:TYR:HB2	1:A:224:TRP:CZ2	2.53	0.43
1:B:41:LEU:O	1:B:41:LEU:HD23	2.19	0.42
1:B:212:ASP:HA	1:B:215:LYS:HB2	2.01	0.42
1:A:78:THR:HG22	1:A:79:SER:N	2.35	0.42
1:B:215:LYS:HE3	1:B:215:LYS:HB3	1.86	0.42
1:A:163:LYS:HE3	1:A:205:CYS:CB	2.50	0.41
1:B:22:PHE:N	1:B:92:ALA:O	2.52	0.41
1:A:132:ASN:C	1:A:132:ASN:HD22	2.23	0.41
1:A:197:LYS:O	1:A:197:LYS:CG	2.69	0.41

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/239 (94%)	222 (99%)	3 (1%)	0	100	100
1	B	222/239 (93%)	216 (97%)	6 (3%)	0	100	100
2	C	2/4 (50%)	2 (100%)	0	0	100	100
2	D	2/4 (50%)	2 (100%)	0	0	100	100
All	All	451/486 (93%)	442 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/212 (97%)	201 (98%)	5 (2%)	49	77
1	B	204/212 (96%)	198 (97%)	6 (3%)	42	72
2	C	4/4 (100%)	4 (100%)	0	100	100
2	D	4/4 (100%)	4 (100%)	0	100	100
All	All	418/432 (97%)	407 (97%)	11 (3%)	46	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	113	PHE
1	A	132	ASN
1	A	173	GLU
1	A	195	GLN
1	A	202	LYS
1	B	9	LYS
1	B	23	HIS
1	B	113	PHE
1	B	157	GLU
1	B	171	MET
1	B	220	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 6.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 6.5 Other polymers

EDS failed to run properly - this section is therefore empty.