



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 03:16 PM UTC

PDB ID : 9CB7 / pdb_00009cb7
EMDB ID : EMD-45418
Title : DeltaNhp10 INO80 bound to S.c 0/40 nucleosome, Ino80-Nucleosome
Authors : Wu, H.; Kaur, U.; Narlikar, G.J.; Cheng, Y.F.
Deposited on : 2024-06-18
Resolution : 4.04 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6625 (3.54 - 4.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	 54% 17% 29%
1	E	136	 53% 18% 29%
2	B	103	 65% 14% 21%
2	F	103	 69% 8% 23%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	132	
3	G	132	
4	D	131	
4	H	131	
5	I	227	
6	J	227	
7	Q	1489	
8	Z	320	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 15448 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	A	97	Total	C	N	O	0	0
			801	508	155	138		
1	E	97	Total	C	N	O	0	0
			801	508	155	138		

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	81	Total	C	N	O	S	0	0
			643	404	126	112	1		
2	F	79	Total	C	N	O	S	0	0
			620	389	120	110	1		

- Molecule 3 is a protein called Histone H2A.1.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	106	Total	C	N	O	0	0
			819	514	161	144		
3	G	108	Total	C	N	O	0	0
			827	517	164	146		

- Molecule 4 is a protein called Histone H2B.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	93	Total	C	N	O	S	0	0
			726	456	127	142	1		
4	H	93	Total	C	N	O	S	0	0
			726	456	127	142	1		

- Molecule 5 is a DNA chain called DNA (227-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	146	Total	C	N	O	P	0	0
			3010	1424	565	875	146		

- Molecule 6 is a DNA chain called DNA (227-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	146	Total	C	N	O	P	0	0
			2976	1413	540	877	146		

- Molecule 7 is a protein called Chromatin-remodeling ATPase INO80.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	Q	425	Total	C	N	O	S	0	0
			3390	2170	588	614	18		

- Molecule 8 is a protein called Ino eighty subunit 2.


Mol	Chain	Residues	Atoms				AltConf	Trace
8	Z	22	Total	C	N	O	0	0
			109	65	22	22		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone H3

Chain A: 



• Molecule 1: Histone H3

Chain E: 



• Molecule 2: Histone H4

Chain B: 




• Molecule 2: Histone H4

Chain F: 



• Molecule 3: Histone H2A.1


Chain C: 




Chain G:  70% 12% 18%




Chain D:  60% 11% 29%



Chain H:  52% 19% 29%



Chain I:  54% 10% 36%



Chain J:  53% 11% 36%

- Molecule 8: Ino eighty subunit 2

Chain Z: 7% 93%

PHE	ARG	SER	ASP	ASP	GLY	ASP	TLE	MET
	ARG	ASP	ASP	ASP	GLY	ARG	ARG	ASP
	ALA	GLU	GLU	ILE	ILE	ASP	ASP	SER
	GLU	LEU	LEU	LEU	LEU	GLY	GLY	ALA
	ASN	VAL	GLU	GLU	GLU	LEU	SER	SER
	ALA	SER	GLU	GLU	GLU	VAL	TLE	ILE
	ARG	VAL	VAL	VAL	GLU	GLU	GLU	GLU
	ARG	ASN	SER	SER	SER	ASP	ASP	ASP
	LYS	GLY	GLY	LEU	LEU	VAL	VAL	ALA
	ASN	ASN	GLU	GLU	GLU	ASP	ASP	GLU
	LEU	GLY	LYS	LYS	LYS	GLU	GLU	LEU
	SER	ASN	GLU	GLU	GLU	ASP	ASP	SER
	GLU	GLU	GLU	ASN	LEU	TYR	ASP	ASP
	LYS	GLU	GLU	ASN	ARG	GLU	GLU	VAL
R256		ASP	ASP	GLY	GLY	GLU	SER	VAL
		ASP	GLU	GLY	GLY	GLU	PRO	GLU
R277		VAL	VAL	GLY	GLY	ASP	GLY	ALA
	SER	GLU	GLU	GLY	GLY	GLU	GLU	GLY
	HIS	GLU	GLU	LYS	LYS	VAL	VAL	GLY
	LEU	ALA	ALA	GLU	GLU	LEU	LEU	GLU
	PRO	THR	THR	VAL	VAL	SER	SER	GLU
	ASN	LYS	GLU	GLU	GLU	PRO	TYR	TYR
	ASP	GLU	LYS	LYS	LYS	SER	HIS	ASP
	ASP	ASN	ASP	GLU	GLU	ASP	ASP	ASP
	GLU	THR	THR	THR	GLU	ASP	ASP	ASP
	LYS	THR	THR	GLU	GLU	ASP	ASP	ASP
	ASN	ASN	THR	THR	GLU	ASP	ASP	ASP
	ASP	SER	SER	SER	THR	VAL	GLN	GLN
	GLY	THR	THR	THR	THR	GLY	LYS	GLN
	SER	ARG	ARG	GLN	GLM	ARG	ILE	ILE
	SER	THR	THR	ASN	LYS	VAL	VAL	THR
	PHE	THR	THR	ASP	GLY	ALA	ALA	LYS
	VAL	THR	THR	VAL	VAL	ASP	ASP	SER
	LYS	ARG	ARG	GLY	GLY	ALA	THR	SER
	PRO	SER	LEU	GLM	GLM	ALA	ALA	ARG
	ARG	MET	LEU	GLY	GLY	SER	THR	ARG
	PRO	MET	LEU	GLY	GLY	LYS	LYS	ARG
	PRO	LEU	LEU	GLU	GLU	THR	THR	ALA
	TYR	ASP	ASP	GLU	GLU	ASP	ASP	ARG
	ASN	LEU	LEU	GLU	GLU	ILE	ILE	ARG
	SER	LEU	LEU	GLM	GLM	ASP	ASP	ARG
	GLU	GLU	GLU	GLY	GLY	GLU	GLU	ARG
	GLY	ASP	ASP	GLY	GLY	THR	THR	ALA
	MET	GLY	GLY	SER	SER	ALA	ALA	ARG
	THR	GLY	GLY	GLY	GLY	ASP	ASP	ARG
	ARG	GLY	GLY	GLY	GLY	ILE	ILE	ARG
	ILE	SER	SER	LYS	LYS	TYR	TYR	ARG
	LEU	LYS	LYS	GLY	GLY	ASP	ASP	VAL
	ARG	ARG	LYS	GLU	GLU	VAL	VAL	VAL
	TYR	LEU	LYS	ASN	ASN	SER	SER	LYS
	GLU	THR	THR	GLU	GLU	GLY	GLY	GLY
	GLU	ASP	ASP	PRO	PRO	ASN	ASN	ASN
	GLU	GLU	GLU	GLU	GLU	ASP	ASP	ASP
	LEU	GLU	GLU	ILE	ILE	GLY	GLY	THR
	LEU	GLU	GLU	THR	THR	GLU	GLU	THR
	PHE	ILE	ILE	THR	THR	SER	SER	SER
	THR	CYS	GLN	LYS	LYS	THR	THR	THR
	THR	LEU	GLU	GLU	GLU	THR	THR	THR

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	40107	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.032	Depositor
Minimum map value	-0.012	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.0033	Depositor
Map size (Å)	421.12, 421.12, 421.12	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.94, 0.94, 0.94	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/812	0.66	3/1086 (0.3%)
1	E	0.26	0/812	0.68	1/1086 (0.1%)
2	B	0.31	0/650	0.86	2/869 (0.2%)
2	F	0.25	0/627	0.67	0/840
3	C	0.25	0/830	0.61	0/1121
3	G	0.26	0/838	0.68	2/1131 (0.2%)
4	D	0.24	0/736	0.67	0/991
4	H	0.31	0/736	0.76	0/991
5	I	0.24	0/3380	0.49	0/5219
6	J	0.23	0/3334	0.42	0/5139
7	Q	0.21	0/3462	0.56	0/4692
8	Z	0.12	0/108	0.37	0/149
All	All	0.24	0/16325	0.57	8/23314 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	H	0	1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	108	VAL	CA-C-N	6.57	129.59	120.65
3	G	108	VAL	C-N-CA	6.57	129.59	120.65
2	B	64	GLU	N-CA-CB	5.94	119.45	110.30
2	B	75	GLU	N-CA-CB	5.72	119.69	110.42
1	A	97	GLU	N-CA-CB	5.55	118.88	110.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	H	106	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	851	18	0
1	E	801	0	851	18	0
2	B	643	0	678	10	0
2	F	620	0	643	7	0
3	C	819	0	865	19	0
3	G	827	0	865	16	0
4	D	726	0	748	12	0
4	H	726	0	748	18	0
5	I	3010	0	1638	23	0
6	J	2976	0	1639	21	0
7	Q	3390	0	3276	50	0
8	Z	109	0	47	0	0
All	All	15448	0	12849	158	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:Q:963:LEU:HD13	7:Q:1010:VAL:HB	1.76	0.67
3:G:51:TYR:HB3	4:H:97:ILE:HD11	1.77	0.66
1:E:63:ARG:HH21	6:J:18:DC:H5''	1.65	0.61
4:H:36:ARG:HH22	6:J:49:DC:H4'	1.65	0.61
7:Q:963:LEU:HA	7:Q:966:VAL:HG22	1.85	0.59

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/136 (70%)	92 (97%)	3 (3%)	0	100	100
1	E	95/136 (70%)	94 (99%)	1 (1%)	0	100	100
2	B	79/103 (77%)	76 (96%)	3 (4%)	0	100	100
2	F	77/103 (75%)	73 (95%)	4 (5%)	0	100	100
3	C	104/132 (79%)	100 (96%)	4 (4%)	0	100	100
3	G	106/132 (80%)	103 (97%)	3 (3%)	0	100	100
4	D	91/131 (70%)	87 (96%)	4 (4%)	0	100	100
4	H	91/131 (70%)	91 (100%)	0	0	100	100
7	Q	415/1489 (28%)	388 (94%)	27 (6%)	0	100	100
8	Z	20/320 (6%)	20 (100%)	0	0	100	100
All	All	1173/2813 (42%)	1124 (96%)	49 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/113 (75%)	85 (100%)	0	100	100
1	E	85/113 (75%)	85 (100%)	0	100	100
2	B	65/79 (82%)	65 (100%)	0	100	100
2	F	62/79 (78%)	62 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	85/99 (86%)	85 (100%)	0	100	100
3	G	84/99 (85%)	84 (100%)	0	100	100
4	D	81/109 (74%)	81 (100%)	0	100	100
4	H	81/109 (74%)	81 (100%)	0	100	100
7	Q	356/1350 (26%)	356 (100%)	0	100	100
All	All	984/2150 (46%)	984 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
7	Q	878	GLN
7	Q	1323	GLN
7	Q	1364	ASN
7	Q	1359	HIS
7	Q	769	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-45418. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 224



Y Index: 224




Z Index: 224


6.2.2 Raw map



X Index: 224



Y Index: 224




Z Index: 224


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 229




Y Index: 279




Z Index: 321


6.3.2 Raw map



X Index: 229



Y Index: 255




Z Index: 289


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0033. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 456 nm^3 ; this corresponds to an approximate mass of 412 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.248 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.248 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.04	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.86	9.75	7.27

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.86 differs from the reported value 4.04 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-45418 and PDB model 9CB7. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.0033 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 96% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0033) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.9650	<div><div></div></div> 0.2880
A	<div><div></div></div> 0.9740	<div><div></div></div> 0.3380
B	<div><div></div></div> 0.9790	<div><div></div></div> 0.3590
C	<div><div></div></div> 0.9700	<div><div></div></div> 0.3390
D	<div><div></div></div> 0.9800	<div><div></div></div> 0.3480
E	<div><div></div></div> 0.9660	<div><div></div></div> 0.3370
F	<div><div></div></div> 0.9820	<div><div></div></div> 0.3960
G	<div><div></div></div> 0.9590	<div><div></div></div> 0.3800
H	<div><div></div></div> 0.9720	<div><div></div></div> 0.3740
I	<div><div></div></div> 0.9580	<div><div></div></div> 0.2810
J	<div><div></div></div> 0.9720	<div><div></div></div> 0.2820
Q	<div><div></div></div> 0.9520	<div><div></div></div> 0.1790
Z	<div><div></div></div> 0.9820	<div><div></div></div> 0.2350

1.0

0.0

<0.0