



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2023 – 06:59 PM EDT

PDB ID : 5CCZ
Title : Crystal structure of Mycobacterium tuberculosis malate synthase in complex with 3-(4-fluorophenyl)-4-methyl-1H-pyrazol-5-amine
Authors : Huang, H.-L.; Sacchettini, J.C.
Deposited on : 2015-07-02
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	741	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Malate synthase G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	713	5451	3420	962	1048	21	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	619	ALA	CYS	engineered mutation	UNP P9WK17

- Molecule 2 is 3-(4-fluorophenyl)-4-methyl-1H-pyrazol-5-amine (three-letter code: 50C) (formula: C₁₀H₁₀FN₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	N		
2	A	1	14	10	1	3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	489	Total	O	0	0
			489	489		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Malate synthase G



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	81.51Å 81.51Å 227.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.88 – 2.14 35.88 – 2.14	Depositor EDS
% Data completeness (in resolution range)	97.5 (35.88-2.14) 97.5 (35.88-2.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.46 (at 2.14Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.193 , 0.250 0.196 , 0.251	Depositor DCC
R_{free} test set	2130 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	33.8	Xtrriage
Anisotropy	0.386	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5955	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 50C, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/5554	0.57	0/7552

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5451	0	5392	44	0
2	A	14	0	0	0	0
3	A	1	0	0	0	0
4	A	489	0	0	12	0
All	All	5955	0	5392	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:704:GLN:HG2	4:A:1367:HOH:O	1.88	0.72
1:A:297:GLY:HA3	1:A:385:LEU:HB2	1.72	0.71
1:A:410:CYS:HB3	4:A:1287:HOH:O	1.96	0.63
1:A:719:ARG:NH1	4:A:911:HOH:O	2.32	0.63
1:A:719:ARG:NH2	4:A:903:HOH:O	2.24	0.62
1:A:414:SER:HB2	4:A:1062:HOH:O	2.02	0.59
1:A:506:SER:HB3	1:A:564:GLN:OE1	2.04	0.56
1:A:524:ASP:O	1:A:528:THR:HG23	2.06	0.55
1:A:479:ARG:NH1	4:A:907:HOH:O	2.28	0.55
1:A:82:ARG:NH1	4:A:922:HOH:O	2.39	0.55
1:A:697:GLU:HG2	1:A:713:ILE:HD13	1.89	0.54
1:A:94:GLU:OE1	1:A:571:ARG:NH2	2.41	0.54
1:A:342:GLY:HA2	1:A:397:LYS:HE3	1.93	0.51
1:A:235:HIS:NE2	4:A:913:HOH:O	2.34	0.50
1:A:317:ASP:HB3	1:A:329:THR:HG23	1.95	0.49
1:A:637:LEU:HB3	1:A:710:THR:HG21	1.95	0.49
1:A:611:ARG:NH2	1:A:673:ASN:OD1	2.46	0.48
1:A:447:LYS:HE3	4:A:1193:HOH:O	2.13	0.48
1:A:7:VAL:HG11	1:A:36:TRP:HB3	1.96	0.47
1:A:469:HIS:CD2	1:A:707:ASN:HA	2.49	0.47
1:A:296:LYS:HA	1:A:382:ASN:O	2.14	0.47
1:A:715:HIS:O	1:A:719:ARG:HG2	2.14	0.47
1:A:16:LEU:HD22	1:A:349:ALA:HB1	1.96	0.46
1:A:281:ALA:HB2	1:A:349:ALA:HA	1.97	0.46
1:A:724:ARG:HB3	4:A:1368:HOH:O	2.15	0.46
1:A:612:TRP:CE3	1:A:714:LEU:HD22	2.50	0.46
1:A:518:MET:HE2	4:A:1261:HOH:O	2.16	0.46
1:A:455:PHE:CZ	1:A:457:ASN:HB3	2.52	0.45
1:A:648:LEU:HD13	1:A:653:ILE:HG13	1.99	0.44
1:A:33:ASP:OD1	1:A:33:ASP:N	2.41	0.44
1:A:648:LEU:HD21	1:A:658:VAL:HG21	1.98	0.43
1:A:670:ASP:OD1	1:A:679:TYR:OH	2.23	0.43
1:A:515:MET:HE1	1:A:541:TRP:HB3	2.00	0.43
1:A:31:ASP:HA	1:A:32:PRO:HD3	1.89	0.43
1:A:683:ALA:HA	1:A:684:PRO:HA	1.79	0.43
1:A:5:VAL:HG21	1:A:17:TYR:CD2	2.54	0.42
1:A:461:LEU:HD22	1:A:639:ILE:HG21	2.00	0.42
1:A:108:GLU:OE1	1:A:389:ARG:NH2	2.50	0.42
1:A:431:ILE:O	1:A:456:ILE:HA	2.19	0.42
1:A:83:GLN:HG2	4:A:1139:HOH:O	2.18	0.42
1:A:674:ALA:C	1:A:676:ASP:H	2.24	0.41
1:A:94:GLU:HA	1:A:95:PRO:HD3	1.90	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:465:GLY:HA3	1:A:638:ARG:HG2	2.01	0.41
1:A:595:ARG:NH1	1:A:599:ASP:OD2	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	707/741 (95%)	687 (97%)	19 (3%)	1 (0%)	51 51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	325	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	573/593 (97%)	548 (96%)	25 (4%)	28 24

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	33	ASP
1	A	50	ASN
1	A	71	ARG
1	A	76	ILE
1	A	82	ARG
1	A	121	VAL
1	A	128	LEU
1	A	186	SER
1	A	199	GLN
1	A	226	SER
1	A	248	GLN
1	A	299	LEU
1	A	385	LEU
1	A	457	ASN
1	A	462	ASP
1	A	528	THR
1	A	583	LEU
1	A	595	ARG
1	A	603	GLN
1	A	637	LEU
1	A	648	LEU
1	A	677	VAL
1	A	698	LEU
1	A	704	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	50C	A	801	-	12,15,15	1.70	5 (41%)	11,21,21	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	50C	A	801	-	-	2/4/4/4	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	50C	C01-C02	-2.67	1.46	1.51
2	A	801	50C	C08-C07	-2.40	1.46	1.49
2	A	801	50C	C07-C02	2.32	1.45	1.40
2	A	801	50C	C10-C11	2.18	1.41	1.37
2	A	801	50C	N06-N05	-2.06	1.33	1.37

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	50C	N06-C07-C08-C09
2	A	801	50C	N06-C07-C08-C14

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	713/741 (96%)	0.06	24 (3%) 45 52	18, 29, 48, 64	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	269	ILE	4.2
1	A	727	GLU	3.9
1	A	118	VAL	3.5
1	A	381	VAL	3.4
1	A	398	PRO	3.3
1	A	432	MET	3.3
1	A	541	TRP	3.0
1	A	338	VAL	3.0
1	A	2	THR	3.0
1	A	337	PHE	2.9
1	A	677	VAL	2.9
1	A	380	ASP	2.8
1	A	587	LEU	2.8
1	A	396	VAL	2.6
1	A	339	ARG	2.6
1	A	151	THR	2.6
1	A	683	ALA	2.3
1	A	341	VAL	2.2
1	A	457	ASN	2.2
1	A	684	PRO	2.1
1	A	675	GLY	2.1
1	A	272	PHE	2.1
1	A	206	ASP	2.1
1	A	431	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	50C	A	801	14/14	0.93	0.15	23,28,37,44	0
3	MG	A	802	1/1	0.99	0.21	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.