



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 4, 2024 – 04:17 PM EST

PDB ID : 3CDL  
Title : Crystal structure of a TetR family transcriptional regulator from *Pseudomonas syringae* pv. tomato str. DC3000  
Authors : Tan, K.; Bigelow, L.; Gu, M.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2008-02-27  
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1460 (2.36-2.36)
Clashscore	180529	1571 (2.36-2.36)
Ramachandran outliers	177936	1559 (2.36-2.36)
Sidechain outliers	177891	1559 (2.36-2.36)
RSRZ outliers	164620	1460 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	203	
1	B	203	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator AefR.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	187	1507	967	261	273	1	5	0	0	0
1	B	188	1520	974	266	273	1	6	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q87Z81
A	-1	ASN	-	expression tag	UNP Q87Z81
A	0	ALA	-	expression tag	UNP Q87Z81
B	-2	SER	-	expression tag	UNP Q87Z81
B	-1	ASN	-	expression tag	UNP Q87Z81
B	0	ALA	-	expression tag	UNP Q87Z81


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	24	Total 24	O 24	0	0
2	B	17	Total 17	O 17	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional regulator AefR



- Molecule 1: Transcriptional regulator AefR



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	132.31Å 53.41Å 55.83Å 90.00° 113.25° 90.00°	Depositor
Resolution (Å)	33.26 – 2.36 33.26 – 2.36	Depositor EDS
% Data completeness (in resolution range)	96.3 (33.26-2.36) 96.5 (33.26-2.36)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.08 (at 2.36Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.207 , 0.275 0.207 , 0.276	Depositor DCC
$R_{free}$ test set	726 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.3	Xtrriage
Anisotropy	0.072	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 53.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.025 for -h+k-l,-l,-k 0.000 for -h-k-l,l,k 0.029 for -h-2*l,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3068	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.74	0/1536	0.75	0/2073
1	B	0.69	4/1552 (0.3%)	0.75	3/2092 (0.1%)
All	All	0.71	4/3088 (0.1%)	0.75	3/4165 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	41	ARG	CZ-NH1	5.95	1.40	1.33
1	B	37	GLU	CD-OE2	5.79	1.32	1.25
1	B	41	ARG	NE-CZ	5.51	1.40	1.33
1	B	37	GLU	CD-OE1	5.11	1.31	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	41	ARG	NE-CZ-NH2	-8.04	116.28	120.30
1	B	41	ARG	NE-CZ-NH1	5.93	123.27	120.30
1	B	121	LEU	CA-CB-CG	5.07	126.96	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	76	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1507	0	1493	49	0
1	B	1520	0	1511	48	0
2	A	24	0	0	0	0
2	B	17	0	0	2	0
All	All	3068	0	3004	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:GLN:H	1:B:179:GLN:HE21	1.17	0.93
1:A:16:ILE:HA	1:A:57:MSE:HE1	1.50	0.92
1:A:7:LYS:O	1:A:11:ILE:HG23	1.71	0.90
1:A:111:ILE:HD11	1:B:108:GLY:HA2	1.52	0.89
1:B:63:ASN:HB3	2:B:207:HOH:O	1.80	0.82
1:A:182:SER:O	1:A:186:GLU:HG2	1.82	0.78
1:B:38:VAL:HG22	1:B:42:THR:OG1	1.83	0.77
1:B:110:THR:HG21	1:B:120:TRP:HD1	1.49	0.75
1:B:83:LEU:HD11	1:B:156[B]:MSE:HE3	1.67	0.74
1:A:111:ILE:CD1	1:B:108:GLY:HA2	2.19	0.71
1:A:191:MSE:HE3	1:B:196:TYR:CE2	2.26	0.70
1:A:45:ASN:C	1:A:45:ASN:HD22	1.96	0.69
1:B:106:VAL:O	1:B:110:THR:HG23	1.93	0.69
1:A:16:ILE:HG12	1:A:57:MSE:HE1	1.76	0.67
1:B:113:SER:OG	1:B:116:ARG:HB2	1.95	0.67
1:A:191:MSE:HE3	1:B:196:TYR:HE2	1.59	0.66
1:B:179:GLN:H	1:B:179:GLN:NE2	1.91	0.65
1:A:79:LEU:HD12	1:A:197:GLU:HG3	1.80	0.64
1:B:182:SER:O	1:B:186:GLU:HB2	1.96	0.64
1:B:83:LEU:HD21	1:B:156[B]:MSE:CE	2.28	0.63
1:A:180:GLU:O	1:A:184:VAL:HG13	2.00	0.62
1:B:55:ALA:O	1:B:59:GLN:HG2	1.99	0.61
1:B:51:GLU:HG3	2:B:216:HOH:O	2.01	0.61

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:116:ARG:HH11	1:B:116:ARG:HG2	1.64	0.61
1:A:51:GLU:HA	1:A:51:GLU:OE1	1.99	0.60
1:A:57:MSE:HE3	1:A:102:LEU:HD13	1.84	0.59
1:B:14:ALA:HB1	1:B:35:ARG:HG2	1.85	0.58
1:B:59:GLN:O	1:B:63:ASN:HB2	2.03	0.58
1:A:119:VAL:HG22	1:A:120:TRP:CD1	2.39	0.58
1:A:116:ARG:HA	1:A:119:VAL:HG13	1.87	0.56
1:A:45:ASN:HD22	1:A:46:HIS:N	2.05	0.55
1:B:38:VAL:CG1	1:B:43:VAL:HG23	2.37	0.55
1:A:151:PHE:HZ	1:B:187:SER:HG	1.56	0.54
1:A:16:ILE:HG12	1:A:57:MSE:CE	2.38	0.53
1:B:87:LEU:HB3	1:B:185:VAL:HG13	1.91	0.53
1:B:83:LEU:HD21	1:B:156[B]:MSE:HE2	1.91	0.53
1:B:59:GLN:HE22	1:B:123:ARG:HD3	1.74	0.53
1:A:139:GLN:NE2	1:A:147:VAL:HG12	2.24	0.52
1:B:88:TRP:HZ2	1:B:182:SER:HB3	1.75	0.52
1:A:79:LEU:CD1	1:A:197:GLU:HG3	2.38	0.52
1:B:26:ILE:O	1:B:31:ARG:NH1	2.42	0.52
1:A:16:ILE:CA	1:A:57:MSE:HE1	2.31	0.51
1:A:144:LEU:HD22	1:A:196:TYR:HB3	1.92	0.51
1:B:177:THR:OG1	1:B:180:GLU:HG3	2.10	0.51
1:A:16:ILE:CG1	1:A:57:MSE:HE1	2.39	0.51
1:A:179:GLN:HE21	1:A:179:GLN:H	1.58	0.50
1:A:16:ILE:HA	1:A:57:MSE:CE	2.32	0.50
1:B:110:THR:HG21	1:B:120:TRP:CD1	2.39	0.50
1:A:144:LEU:HD23	1:A:197:GLU:HA	1.93	0.49
1:A:11:ILE:HD13	1:A:43:VAL:HG13	1.95	0.49
1:B:109:ALA:O	1:B:116:ARG:NH1	2.46	0.49
1:B:179:GLN:HE21	1:B:179:GLN:N	1.99	0.49
1:A:95:THR:CG2	1:A:95:THR:O	2.61	0.48
1:B:91:MSE:HE2	1:B:160:LEU:HG	1.94	0.48
1:A:30:ASP:HA	1:A:33:ALA:HB3	1.95	0.47
1:B:180:GLU:O	1:B:184:VAL:HG13	2.13	0.47
1:A:82:GLN:HG3	1:A:134:TRP:HZ2	1.78	0.47
1:A:30:ASP:HB3	1:A:40:LYS:HG3	1.97	0.47
1:A:107:VAL:O	1:A:111:ILE:HG23	2.16	0.46
1:B:100:LEU:HD12	1:B:175:LEU:HD11	1.98	0.45
1:B:38:VAL:CG2	1:B:42:THR:OG1	2.59	0.45
1:A:101:ASP:OD1	1:A:104:ARG:NH2	2.50	0.45
1:B:54:PHE:CE2	1:B:116:ARG:HG3	2.51	0.45
1:A:191:MSE:HE1	1:B:192:PHE:CD1	2.52	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:LYS:HE2	1:A:44:TYR:HE1	1.82	0.44
1:B:94:LEU:HD23	1:B:164:ALA:HB3	1.99	0.44
1:A:177:THR:O	1:A:181:GLN:HG3	2.18	0.44
1:B:29:MSE:HE3	1:B:43:VAL:HG12	1.99	0.43
1:A:82:GLN:HG3	1:A:134:TRP:CZ2	2.53	0.43
1:B:18:GLU:HG2	1:B:35:ARG:HD3	2.00	0.43
1:B:193:LEU:HD23	1:B:193:LEU:HA	1.79	0.43
1:A:179:GLN:H	1:A:179:GLN:NE2	2.16	0.43
1:A:45:ASN:C	1:A:45:ASN:ND2	2.69	0.43
1:A:84:LEU:HD22	1:A:88:TRP:HD1	1.84	0.43
1:B:131:PHE:CD1	1:B:157:HIS:CE1	3.07	0.43
1:A:11:ILE:HD11	1:A:47:PHE:CD2	2.54	0.42
1:A:138:ALA:HB3	1:A:144:LEU:HD12	2.00	0.42
1:A:159:LEU:HD13	1:A:188:ALA:HA	2.02	0.42
1:A:111:ILE:HG12	1:A:112:HIS:N	2.34	0.42
1:B:181:GLN:O	1:B:185:VAL:HG23	2.20	0.41
1:B:23:GLY:O	1:B:27:THR:OG1	2.23	0.41
1:B:38:VAL:HG13	1:B:43:VAL:HG23	2.01	0.41
1:B:54:PHE:CZ	1:B:116:ARG:HG3	2.56	0.41
1:A:95:THR:O	1:A:95:THR:HG22	2.20	0.41
1:A:110:THR:HB	1:A:117:ALA:HB2	2.03	0.41
1:A:111:ILE:CG1	1:A:112:HIS:N	2.84	0.41
1:B:22:ARG:HB3	1:B:26:ILE:HD11	2.03	0.41
1:B:121:LEU:HA	1:B:124:ILE:HG22	2.02	0.40
1:A:116:ARG:O	1:A:120:TRP:HD1	2.05	0.40
1:B:87:LEU:HD21	1:B:156[B]:MSE:CE	2.51	0.40
1:A:30:ASP:HB3	1:A:40:LYS:CG	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/203 (89%)	172 (95%)	8 (4%)	1 (1%)	22	24
1	B	183/203 (90%)	174 (95%)	8 (4%)	1 (0%)	25	28
All	All	364/406 (90%)	346 (95%)	16 (4%)	2 (0%)	22	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	71	VAL
1	B	75	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/166 (96%)	129 (81%)	30 (19%)	1	1
1	B	160/166 (96%)	129 (81%)	31 (19%)	1	1
All	All	319/332 (96%)	258 (81%)	61 (19%)	1	1

All (61) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	4	THR
1	A	8	ARG
1	A	11	ILE
1	A	29	MSE
1	A	44	TYR
1	A	45	ASN
1	A	52	GLU
1	A	53	LEU
1	A	56	GLU
1	A	57	MSE
1	A	72	VAL
1	A	84	LEU
1	A	90	LYS

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	94	LEU
1	A	102	LEU
1	A	105	VAL
1	A	111	ILE
1	A	119	VAL
1	A	121	LEU
1	A	129	GLU
1	A	132	SER
1	A	140	LYS
1	A	156	MSE
1	A	172	ASN
1	A	175	LEU
1	A	179	GLN
1	A	182	SER
1	A	184	VAL
1	A	191	MSE
1	B	8	ARG
1	B	30	ASP
1	B	35	ARG
1	B	52	GLU
1	B	56	GLU
1	B	62	TRP
1	B	63	ASN
1	B	76	LEU
1	B	77	VAL
1	B	84	LEU
1	B	87	LEU
1	B	102	LEU
1	B	105	VAL
1	B	110	THR
1	B	116	ARG
1	B	118	GLN
1	B	120	TRP
1	B	121	LEU
1	B	123	ARG
1	B	129	GLU
1	B	130	THR
1	B	140	LYS
1	B	145	LYS
1	B	156[A]	MSE
1	B	156[B]	MSE
1	B	159	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	161	LYS
1	B	175	LEU
1	B	179	GLN
1	B	183	ASN
1	B	189	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	155	GLN
1	A	179	GLN
1	A	183	ASN
1	B	45	ASN
1	B	93	ASN
1	B	118	GLN
1	B	139	GLN
1	B	155	GLN
1	B	157	HIS
1	B	179	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	182/203 (89%)	0.62	16 (8%) <span style="border: 1px solid red; padding: 2px;">17</span> <span style="border: 1px solid red; padding: 2px;">20</span>	40, 50, 63, 74	0
1	B	183/203 (90%)	0.38	4 (2%) <span style="border: 1px solid blue; padding: 2px;">62</span> <span style="border: 1px solid blue; padding: 2px;">67</span>	41, 49, 66, 83	0
All	All	365/406 (89%)	0.50	20 (5%) <span style="border: 1px solid red; padding: 2px;">32</span> <span style="border: 1px solid red; padding: 2px;">37</span>	40, 50, 64, 83	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3	LEU	5.8
1	A	3	LEU	4.9
1	A	120	TRP	3.4
1	A	26	ILE	3.3
1	A	15	ALA	2.6
1	A	23	GLY	2.5
1	A	17	ALA	2.5
1	A	16	ILE	2.4
1	A	24	PHE	2.4
1	B	76	LEU	2.3
1	B	64	CYS	2.3
1	A	43	VAL	2.3
1	A	22	ARG	2.2
1	A	121	LEU	2.2
1	A	36	ALA	2.2
1	A	128	GLU	2.1
1	B	126	GLU	2.1
1	A	131	PHE	2.0
1	A	20	GLY	2.0
1	A	61	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.