



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2023 – 01:22 AM JST

PDB ID : 7CD0  
Title : Crystal structure of the 2-iodoporphobilinogen-bound ES2 intermediate form of human hydroxymethylbilane synthase  
Authors : Sato, H.; Sugishima, M.; Wada, K.; Hirabayashi, K.; Tsukaguchi, M.  
Deposited on : 2020-06-18  
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	361	 77% 12% 11%
1	B	361	 73% 16% 11%

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Porphobilinogen deaminase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	Total	C	N	O	S	0	0	0
			2460	1549	449	453	9			
1	B	320	Total	C	N	O	S	0	0	0
			2434	1533	446	446	9			

- Molecule 2 is 3-[4-(2-hydroxy-2-oxoethyl)-5-[[4-(2-hydroxy-2-oxoethyl)-5-[[4-(2-hydroxy-2-oxoethyl)-5-[[4-(2-hydroxy-2-oxoethyl)-3-(3-hydroxy-3-oxopropyl)-5-methyl-1 {H}-pyrrol-2-yl]methyl]-3-(3-hydroxy-3-oxopropyl)-1 {H}-pyrrol-2-yl]methyl]-3-(3-hydroxy-3-oxopropyl)-1 {H}-pyrrol-2-yl]methyl]-1 {H}-pyrrol-3-yl]propanoic acid (three-letter code: 7J8) (formula: C<sub>40</sub>H<sub>46</sub>N<sub>4</sub>O<sub>16</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total	C	N	O	0	0
			60	40	4	16		
2	B	1	Total	C	N	O	0	0
			60	40	4	16		

- Molecule 3 is 3-[5-(aminomethyl)-4-(carboxymethyl)-2-iodo-1H-pyrrol-3-yl]propanoic acid (three-letter code: FWL) (formula: C<sub>10</sub>H<sub>13</sub>IN<sub>2</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	I	N	O		
3	B	1	17	10	1	2	4	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	54	54	54	0	0
4	B	22	22	22	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Porphobilinogen deaminase

Chain A: 



- Molecule 1: Porphobilinogen deaminase

Chain B: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.42Å 81.37Å 108.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.25 – 2.31 45.25 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.25-2.31) 99.5 (45.25-2.31)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 2.32Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.227 , 0.277 0.227 , 0.276	Depositor DCC
$R_{free}$ test set	1614 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.8	Xtrriage
Anisotropy	0.312	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5107	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.5549e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7J8, FWL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/2502	0.51	0/3394
1	B	0.39	1/2476 (0.0%)	0.62	4/3360 (0.1%)
All	All	0.33	1/4978 (0.0%)	0.57	4/6754 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	115	LYS	CD-CE	5.38	1.64	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	56	MET	CB-CG-SD	-5.38	96.25	112.40
1	B	132	LYS	CA-CB-CG	5.34	125.15	113.40
1	B	115	LYS	CG-CD-CE	-5.16	96.43	111.90
1	B	132	LYS	CB-CG-CD	-5.15	98.22	111.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	26	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2460	0	2490	36	0
1	B	2434	0	2461	37	0
2	A	60	0	0	2	0
2	B	60	0	0	1	0
3	B	17	0	0	1	0
4	A	54	0	0	1	0
4	B	22	0	0	1	0
All	All	5107	0	4951	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:VAL:HG13	1:B:162:GLU:HG2	1.25	1.15
1:A:194:GLN:HE21	1:A:199:HIS:HD2	1.10	0.94
1:B:251:ARG:NH1	1:B:316:VAL:O	2.12	0.81
1:B:142:VAL:HA	1:B:162:GLU:HB3	1.62	0.81
1:A:194:GLN:HE21	1:A:199:HIS:CD2	2.02	0.73
1:B:352:ASP:O	1:B:356:GLN:HG2	1.92	0.69
1:A:255:ARG:NH1	4:A:502:HOH:O	2.26	0.67
1:A:295:MET:HE1	1:A:337:SER:HB2	1.76	0.66
1:B:29:GLN:NE2	4:B:501:HOH:O	2.29	0.66
1:B:172:THR:HA	1:B:175:ARG:HE	1.62	0.65
1:A:352:ASP:OD1	1:A:355:ARG:NH2	2.31	0.64
1:A:139:GLU:OE2	1:A:140:LYS:NZ	2.32	0.60
1:A:126:HIS:ND1	1:A:128:LYS:HG2	2.16	0.60
1:B:348:LYS:NZ	1:B:352:ASP:OD1	2.33	0.59
1:A:194:GLN:NE2	1:A:199:HIS:HD2	1.92	0.58
1:B:101:PRO:O	1:B:251:ARG:NH2	2.36	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:GLU:CG	1:A:140:LYS:HG3	2.34	0.57
1:A:139:GLU:HG3	1:A:140:LYS:HG3	1.85	0.57
1:A:275:GLN:NE2	1:A:296:GLN:OE1	2.37	0.57
1:A:98:LYS:O	2:A:401:7J8:N2	2.38	0.55
1:B:127:PRO:HA	1:B:130:VAL:HG23	1.88	0.55
1:B:285:LEU:HD12	1:B:357:LEU:HD22	1.88	0.55
1:A:245:LEU:HB2	1:A:271:MET:HE1	1.89	0.53
1:B:126:HIS:HD2	1:B:127:PRO:N	2.07	0.52
1:A:265:VAL:HG22	1:A:282:VAL:HG12	1.90	0.52
1:B:112:ALA:HB1	1:B:238:LEU:HB2	1.92	0.51
1:B:142:VAL:HA	1:B:162:GLU:CB	2.36	0.50
1:B:347:ALA:O	1:B:351:LEU:HG	2.11	0.50
1:B:22:ARG:NH2	1:B:88:ASN:O	2.44	0.50
1:B:191:ALA:O	1:B:195:ARG:HG3	2.12	0.49
1:A:170:LEU:HD21	1:A:193:LEU:HD23	1.94	0.48
1:A:245:LEU:HB2	1:A:271:MET:CE	2.43	0.48
1:A:315:LEU:HD11	1:A:329:LEU:HD12	1.94	0.48
1:A:295:MET:HE1	1:A:337:SER:CB	2.44	0.47
1:A:139:GLU:HG2	1:A:140:LYS:NZ	2.29	0.47
1:A:348:LYS:HE3	1:A:352:ASP:OD2	2.14	0.47
1:B:117:GLU:OE1	1:B:195:ARG:NH2	2.48	0.47
1:B:142:VAL:CG1	1:B:162:GLU:HG2	2.19	0.47
1:B:154:LEU:HB3	1:B:163:PHE:CE2	2.50	0.46
1:A:112:ALA:HB1	1:A:238:LEU:HB2	1.96	0.46
1:A:146:SER:HB2	2:A:401:7J8:O11	2.16	0.46
1:B:26:ARG:HG3	1:B:81:LEU:HD21	1.97	0.46
1:A:26:ARG:HD2	1:A:77:PHE:CE1	2.50	0.46
1:B:147:SER:HB3	1:B:150:ARG:HB2	1.96	0.46
1:B:168:GLY:O	1:B:173:ARG:NE	2.47	0.45
1:A:105:PRO:HB2	1:A:108:PHE:CD2	2.51	0.45
1:A:43:LYS:HZ3	1:B:130:VAL:HB	1.81	0.45
1:B:104:LEU:HD11	1:B:110:ILE:HG13	1.98	0.45
1:B:265:VAL:HG22	1:B:282:VAL:HG12	1.99	0.45
1:B:275:GLN:NE2	1:B:296:GLN:OE1	2.49	0.44
1:B:169:ASN:HB2	3:B:402:FWL:O2A	2.18	0.44
1:B:117:GLU:OE1	1:B:195:ARG:NE	2.51	0.43
1:B:123:VAL:CG1	1:B:204:GLN:HB3	2.49	0.43
1:A:128:LYS:NZ	1:A:129:PHE:CE2	2.87	0.43
1:A:39:VAL:HG12	1:A:43:LYS:HD2	2.01	0.42
1:B:259:GLY:O	1:B:355:ARG:HD3	2.18	0.42
1:A:139:GLU:HG2	1:A:140:LYS:HZ3	1.83	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:PRO:HB2	1:A:108:PHE:HD2	1.85	0.42
1:B:86:GLU:HB2	1:B:108:PHE:HZ	1.84	0.42
1:A:284:SER:HB3	1:A:289:ASP:HB2	2.01	0.42
1:A:215:VAL:HG21	1:A:254:LEU:HB2	2.02	0.42
1:B:174:LEU:HD23	1:B:186:ILE:HD11	2.01	0.42
1:A:86:GLU:HG3	1:A:87:LYS:HG3	2.02	0.41
1:A:193:LEU:HD23	1:A:193:LEU:HA	1.87	0.41
1:B:126:HIS:HE1	1:B:181:GLN:HG2	1.85	0.41
1:B:149:ARG:HH21	2:B:401:7J8:C18	2.34	0.41
1:A:179:GLU:HB3	1:B:53:ILE:HD13	2.01	0.41
1:B:263:VAL:HA	1:B:264:PRO:HD3	1.93	0.41
1:A:22:ARG:HB3	1:A:54:ILE:HD13	2.02	0.40
1:A:147:SER:HB3	1:A:150:ARG:HD2	2.03	0.40
1:B:255:ARG:CZ	1:B:316:VAL:HG12	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/361 (88%)	311 (98%)	8 (2%)	0	100	100
1	B	316/361 (88%)	306 (97%)	10 (3%)	0	100	100
All	All	635/722 (88%)	617 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/299 (88%)	263 (100%)	0	100	100
1	B	259/299 (87%)	256 (99%)	3 (1%)	71	83
All	All	522/598 (87%)	519 (99%)	3 (1%)	86	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26	ARG
1	B	27	LYS
1	B	88	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	HIS
1	A	275	GLN
1	A	296	GLN
1	B	126	HIS
1	B	304	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FWL	B	402	-	14,17,17	2.59	3 (21%)	15,23,23	2.02	3 (20%)
2	7J8	A	401	1	56,63,63	1.63	8 (14%)	54,89,89	1.39	8 (14%)
2	7J8	B	401	1	56,63,63	1.62	7 (12%)	54,89,89	1.52	6 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FWL	B	402	-	-	0/9/11/11	0/1/1/1
2	7J8	A	401	1	-	7/42/48/48	0/4/4/4
2	7J8	B	401	1	-	6/42/48/48	0/4/4/4

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	FWL	C2A-C1A	7.30	1.50	1.39
2	A	401	7J8	C27-N	4.38	1.42	1.36
2	B	401	7J8	C27-N	4.31	1.42	1.36
3	B	402	FWL	C4A-I4A	4.20	2.15	2.11
3	B	402	FWL	C3A-C2A	3.96	1.48	1.39
2	B	401	7J8	C21-C22	3.80	1.54	1.51
2	A	401	7J8	C9-C12	3.67	1.47	1.39
2	B	401	7J8	C9-C12	3.54	1.47	1.39
2	A	401	7J8	C21-C22	3.50	1.54	1.51
2	A	401	7J8	C21-C20	3.28	1.54	1.51
2	B	401	7J8	C21-C20	3.16	1.53	1.51
2	B	401	7J8	C7-C8	2.88	1.53	1.51
2	A	401	7J8	C7-C8	2.79	1.53	1.51
2	A	401	7J8	C7-C6	2.62	1.53	1.51
2	B	401	7J8	C7-C6	2.43	1.53	1.51
2	A	401	7J8	C14-C13	2.18	1.53	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	7J8	C26-C23	2.05	1.43	1.40
2	A	401	7J8	C26-C23	2.00	1.43	1.40

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	7J8	C9-C10-C11	-5.22	107.97	114.21
3	B	402	FWL	C7A-C3A-C4A	4.64	130.56	127.30
3	B	402	FWL	C2A-C5A-C6A	-4.14	109.26	114.21
2	B	401	7J8	C31-C19-C20	-3.03	125.17	127.30
2	B	401	7J8	C16-C17-C18	-2.95	110.69	114.21
3	B	402	FWL	CHA-C1A-C2A	-2.94	124.38	130.00
2	A	401	7J8	C34-C35-C36	-2.82	105.84	113.76
2	A	401	7J8	C31-C19-C20	-2.63	125.45	127.30
2	A	401	7J8	C31-C32-C33	-2.40	107.03	113.76
2	B	401	7J8	C31-C32-C33	-2.40	107.04	113.76
2	A	401	7J8	C23-C24-C25	-2.38	111.37	114.21
2	B	401	7J8	C37-C38-C39	-2.34	107.21	113.76
2	A	401	7J8	C37-C38-C39	-2.15	107.74	113.76
2	B	401	7J8	O14-C39-C38	2.07	120.67	114.03
2	A	401	7J8	O14-C39-C38	2.03	120.56	114.03
2	A	401	7J8	O12-C36-C35	2.02	120.53	114.03
2	A	401	7J8	C37-C5-C6	-2.01	125.88	127.30

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	401	7J8	C9-C10-C11-O2
2	B	401	7J8	C9-C10-C11-O3
2	A	401	7J8	C9-C10-C11-O3
2	A	401	7J8	C9-C10-C11-O2
2	B	401	7J8	C28-C29-C30-O9
2	A	401	7J8	C28-C29-C30-O9
2	B	401	7J8	C28-C29-C30-O8
2	A	401	7J8	C2-C3-C4-O1
2	A	401	7J8	C23-C24-C25-O7
2	A	401	7J8	C28-C29-C30-O8
2	B	401	7J8	C31-C32-C33-O10
2	A	401	7J8	C23-C24-C25-O6
2	B	401	7J8	C31-C32-C33-O11


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	FWL	1	0
2	A	401	7J8	2	0
2	B	401	7J8	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)




Unable to reproduce the depositors R factor - this section is therefore empty.


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around 7J8 B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.