



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 09:11 PM EDT

PDB ID : 4CF3  
Title : Mutagenesis of a Rhodobacteraceae L-haloacid dehalogenase  
Authors : Novak, H.R.; Sayer, C.; Isupov, M.N.; Littlechild, J.A.  
Deposited on : 2013-11-13  
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	 83% 12% 5%
1	B	236	 84% 12% 5%
1	C	236	 2% 80% 15% 5%
1	D	236	 12% 74% 20% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	TRS	A	240	-	X	-	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called L-HALOACID DEHALOGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	225	Total 1749	C 1121	N 293	O 329	S 6	0	3	0
1	B	228	Total 1752	C 1121	N 294	O 331	S 6	0	0	0
1	C	225	Total 1728	C 1107	N 288	O 327	S 6	0	0	0
1	D	225	Total 1728	C 1107	N 288	O 327	S 6	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	183	ALA	HIS	engineered mutation	UNP M9P6K0
B	183	ALA	HIS	engineered mutation	UNP M9P6K0
C	183	ALA	HIS	engineered mutation	UNP M9P6K0
D	183	ALA	HIS	engineered mutation	UNP M9P6K0

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	8	4	1	3	0	0
2	B	1	8	4	1	3	0	0
2	C	1	8	4	1	3	0	0
2	D	1	8	4	1	3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	113	113	113	0	0
3	B	99	99	99	0	0
3	C	63	63	63	0	0
3	D	18	18	18	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-HALOACID DEHALOGENASE




- Molecule 1: L-HALOACID DEHALOGENASE




- Molecule 1: L-HALOACID DEHALOGENASE



- Molecule 1: L-HALOACID DEHALOGENASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.17Å 68.27Å 283.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.86 – 2.16 36.86 – 2.16	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.86-2.16) 99.9 (36.86-2.16)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.77 (at 2.16Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.182 , 0.254 0.182 , 0.252	Depositor DCC
$R_{free}$ test set	2330 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.0	Xtrriage
Anisotropy	0.467	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 49.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7282	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, ASB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/1783	0.71	0/2434
1	B	0.55	0/1784	0.69	0/2434
1	C	0.52	0/1759	0.69	0/2401
1	D	0.44	0/1759	0.64	1/2401 (0.0%)
All	All	0.52	0/7085	0.68	1/9670 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	52	LEU	CA-CB-CG	5.09	127.01	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1749	0	1742	20	0
1	B	1752	0	1739	20	0
1	C	1728	0	1713	25	0
1	D	1728	0	1713	30	0
2	A	8	0	12	0	0
2	B	8	0	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	8	0	12	0	0
2	D	8	0	12	0	0
3	A	113	0	0	5	0
3	B	99	0	0	5	0
3	C	63	0	0	0	0
3	D	18	0	0	1	0
All	All	7282	0	6955	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:ARG:HD2	3:A:2104:HOH:O	1.73	0.88
1:C:80:ASN:OD1	1:D:206:THR:HB	1.82	0.79
1:D:136:GLU:HA	1:D:141:ALA:HB2	1.67	0.77
1:D:184:ILE:HG13	1:D:207:PRO:HA	1.68	0.76
1:C:116:ALA:HB2	1:C:231:ILE:HD11	1.74	0.70
1:C:116:ALA:CB	1:C:231:ILE:CD1	2.70	0.69
1:C:116:ALA:HB2	1:C:231:ILE:CD1	2.27	0.65
1:D:73:VAL:O	1:D:77:VAL:HG23	1.97	0.65
1:C:116:ALA:CB	1:C:231:ILE:HD11	2.28	0.64
1:D:132:PRO:O	1:D:137:LYS:HE3	1.98	0.63
1:D:148:LEU:HD12	1:D:166:THR:HG23	1.79	0.63
1:D:30:SER:HB2	1:D:31:PRO:HD3	1.80	0.62
1:A:184[B]:ILE:HG13	1:A:207:PRO:HA	1.81	0.60
1:C:184:ILE:HG13	1:C:207:PRO:HA	1.83	0.59
1:B:211:ALA:O	1:B:212:GLU:HB2	2.02	0.59
1:D:21:GLU:HB3	1:D:182:CYS:SG	2.44	0.58
1:A:44:ARG:HD2	1:A:204:HIS:CD2	2.39	0.57
1:B:184:ILE:HG13	1:B:207:PRO:HA	1.85	0.57
1:C:80:ASN:OD1	1:D:206:THR:CB	2.53	0.56
1:A:115:ASP:HB3	3:A:2060:HOH:O	2.05	0.56
1:C:50:LEU:HD13	1:C:70:ALA:HA	1.88	0.55
1:C:20:ASN:O	1:C:21:GLU:HB2	2.05	0.55
1:D:14:ILE:HB	1:D:177:LEU:HD23	1.90	0.54
1:C:206:THR:H	1:D:80:ASN:HD21	1.56	0.53
1:C:211:ALA:O	1:D:64:ARG:HD3	2.08	0.53
1:D:136:GLU:CA	1:D:141:ALA:HB2	2.38	0.53
1:D:191:GLN:NE2	3:D:2013:HOH:O	2.40	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203:PRO:HG2	3:B:2008:HOH:O	2.10	0.52
1:B:184:ILE:O	1:B:188:ILE:HG13	2.10	0.51
1:B:12:SER:HB3	1:B:234:LEU:HD21	1.93	0.51
1:B:86:THR:OG1	1:B:88:ASP:HB2	2.10	0.51
1:A:94:LYS:HG3	3:A:2046:HOH:O	2.10	0.50
1:D:50:LEU:HD12	1:D:69:ILE:HG22	1.94	0.50
1:B:123:LEU:HG	1:B:157:LYS:HE3	1.93	0.50
1:D:219:ILE:HG22	1:D:220:GLY:N	2.27	0.49
1:A:206:THR:OG1	1:B:80:ASN:ND2	2.44	0.49
1:D:148:LEU:CD1	1:D:166:THR:HG23	2.42	0.49
1:D:37:PHE:HB3	1:D:42:VAL:HG21	1.94	0.49
1:D:110:LEU:HD22	1:D:120:LEU:HD13	1.94	0.49
1:A:48:PRO:HB2	1:B:48:PRO:HB2	1.95	0.49
1:A:175:GLU:O	1:A:196:ARG:NH1	2.46	0.49
1:A:20:ASN:O	1:A:21:GLU:HB2	2.15	0.47
1:C:64:ARG:NH1	1:C:68:GLU:OE1	2.48	0.46
1:D:132:PRO:HA	1:D:136:GLU:OE1	2.15	0.46
1:D:14:ILE:HD11	1:D:119:ARG:NH1	2.31	0.46
1:A:123:LEU:HG	1:A:157:LYS:HE3	1.98	0.46
1:B:41:LYS:N	1:B:41:LYS:HD3	2.30	0.46
1:A:204:HIS:ND1	3:A:2099:HOH:O	2.36	0.46
1:B:50:LEU:HD13	1:B:70:ALA:HA	1.98	0.45
1:D:199:PHE:CE2	1:D:207:PRO:HD3	2.52	0.45
1:A:21:GLU:HB3	1:A:182:CYS:SG	2.57	0.45
1:B:219:ILE:O	1:B:229:GLN:NE2	2.50	0.45
1:A:118:PHE:CZ	1:A:234:LEU:HD21	2.52	0.45
1:C:14:ILE:HB	1:C:177:LEU:HD23	1.99	0.45
1:C:206:THR:N	1:D:80:ASN:HD21	2.15	0.44
1:D:30:SER:HB2	1:D:31:PRO:CD	2.47	0.44
1:A:168:GLU:HB3	3:A:2087:HOH:O	2.17	0.44
1:D:202:ARG:HB3	1:D:203:PRO:HD2	1.99	0.44
1:D:219:ILE:CG2	1:D:220:GLY:N	2.81	0.43
1:D:44:ARG:HD2	1:D:204:HIS:CD2	2.53	0.43
1:A:206:THR:H	1:B:80:ASN:HD21	1.65	0.43
1:C:187:THR:O	1:C:191:GLN:HG3	2.17	0.43
1:C:97:LEU:HD23	1:C:97:LEU:HA	1.85	0.43
1:D:32:LEU:HD13	1:D:92:GLU:CD	2.39	0.42
1:A:112:ARG:NH1	1:A:228:ASP:OD1	2.44	0.42
1:B:84:LYS:HB2	3:B:2034:HOH:O	2.18	0.42
1:A:144:PHE:HB2	1:A:147:HIS:CE1	2.55	0.42
1:A:206:THR:N	1:B:80:ASN:HD21	2.18	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:201:ALA:O	1:C:202:ARG:HD3	2.20	0.41
1:C:75:GLU:O	1:C:78:ALA:HB3	2.19	0.41
1:B:174:PRO:HD3	3:B:2086:HOH:O	2.20	0.41
1:C:30:SER:HB2	1:C:31:PRO:HD3	2.02	0.41
1:C:199:PHE:CE2	1:C:207:PRO:HD3	2.56	0.41
1:B:11:ARG:HG3	3:B:2051:HOH:O	2.21	0.41
1:B:150:VAL:HB	1:B:155:ARG:O	2.20	0.41
1:C:150:VAL:HB	1:C:155:ARG:O	2.20	0.41
1:A:14:ILE:HB	1:A:177:LEU:HD23	2.03	0.41
1:B:129:SER:HB3	1:B:130:PRO:O	2.21	0.40
1:B:229:GLN:HG2	3:B:2099:HOH:O	2.19	0.40
1:C:39:ASP:OD1	1:C:41:LYS:HB2	2.22	0.40
1:C:26:LEU:HG	1:C:26:LEU:O	2.22	0.40
1:C:148:LEU:HD12	1:C:166:THR:HG23	2.03	0.40
1:D:105:ASP:OD2	1:D:222:ASP:HB2	2.21	0.40
1:D:173:LYS:HB2	1:D:174:PRO:CD	2.50	0.40
1:A:107:ALA:N	1:A:108:PRO:CD	2.85	0.40
1:C:202:ARG:O	1:C:203:PRO:C	2.60	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/236 (95%)	222 (99%)	3 (1%)	0	100	100
1	B	225/236 (95%)	219 (97%)	6 (3%)	0	100	100
1	C	222/236 (94%)	213 (96%)	9 (4%)	0	100	100
1	D	222/236 (94%)	208 (94%)	13 (6%)	1 (0%)	29	22
All	All	894/944 (95%)	862 (96%)	31 (4%)	1 (0%)	51	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	42	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/193 (96%)	183 (99%)	2 (1%)	73	78
1	B	186/193 (96%)	182 (98%)	4 (2%)	52	55
1	C	183/193 (95%)	180 (98%)	3 (2%)	62	67
1	D	183/193 (95%)	174 (95%)	9 (5%)	25	21
All	All	737/772 (96%)	719 (98%)	18 (2%)	49	51

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	LEU
1	A	94	LYS
1	B	41	LYS
1	B	64	ARG
1	B	94	LYS
1	B	208	LEU
1	C	64	ARG
1	C	129	SER
1	C	219	ILE
1	D	14	ILE
1	D	41	LYS
1	D	89	ASP
1	D	94	LYS
1	D	97	LEU
1	D	168	GLU
1	D	209	THR
1	D	233	SER
1	D	235	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	80	ASN
1	B	82	GLN
1	C	20	ASN
1	C	114	GLN
1	D	80	ASN
1	D	191	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	ASB	A	18	1	10,11,12	1.51	1 (10%)	9,13,15	2.43	3 (33%)
1	ASB	B	18	1	10,11,12	1.46	1 (10%)	9,13,15	2.10	3 (33%)
1	ASB	C	18	1	10,11,12	1.67	2 (20%)	9,13,15	2.56	4 (44%)
1	ASB	D	18	1	10,11,12	1.61	1 (10%)	9,13,15	2.62	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ASB	A	18	1	-	4/10/11/13	-
1	ASB	B	18	1	-	5/10/11/13	-
1	ASB	C	18	1	-	5/10/11/13	-
1	ASB	D	18	1	-	2/10/11/13	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	18	ASB	OD1-CG	4.21	1.45	1.33
1	A	18	ASB	OD1-CG	4.05	1.45	1.33
1	C	18	ASB	OD1-CG	3.99	1.45	1.33
1	B	18	ASB	OD1-CG	3.70	1.44	1.33
1	C	18	ASB	O1-C1	-2.02	1.24	1.30

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	18	ASB	OD1-CG-CB	6.17	122.55	111.17
1	D	18	ASB	OD1-CG-CB	5.97	122.18	111.17
1	A	18	ASB	OD1-CG-CB	5.10	120.57	111.17
1	B	18	ASB	OD1-CG-CB	4.25	119.00	111.17
1	B	18	ASB	OD1-C2-C1	-3.48	103.44	110.93
1	D	18	ASB	OD1-C2-C1	-3.39	103.64	110.93
1	A	18	ASB	OD1-C2-C1	-3.33	103.77	110.93
1	C	18	ASB	OD1-C2-C1	-3.13	104.19	110.93
1	D	18	ASB	OD2-CG-CB	-2.66	118.46	124.65
1	B	18	ASB	OD1-CG-OD2	-2.31	117.86	123.63
1	C	18	ASB	OD1-CG-OD2	-2.29	117.89	123.63
1	C	18	ASB	OD2-CG-CB	-2.19	119.56	124.65
1	A	18	ASB	O1-C1-C2	2.05	123.32	113.66

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	18	ASB	O1-C1-C2-OD1
1	A	18	ASB	O2-C1-C2-OD1
1	B	18	ASB	O2-C1-C2-OD1
1	C	18	ASB	O2-C1-C2-OD1
1	B	18	ASB	O1-C1-C2-OD1
1	C	18	ASB	O1-C1-C2-OD1
1	D	18	ASB	O1-C1-C2-OD1
1	D	18	ASB	O2-C1-C2-OD1
1	C	18	ASB	C1-C2-OD1-CG
1	A	18	ASB	CA-CB-CG-OD1
1	B	18	ASB	CA-CB-CG-OD1
1	C	18	ASB	CA-CB-CG-OD1
1	C	18	ASB	CA-CB-CG-OD2
1	B	18	ASB	C1-C2-OD1-CG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
1	A	18	ASB	CA-CB-CG-OD2
1	B	18	ASB	CA-CB-CG-OD2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TRS	A	240	-	7,7,7	0.37	0	9,9,9	5.14	6 (66%)
2	TRS	C	240	-	7,7,7	0.55	0	9,9,9	0.58	0
2	TRS	B	240	-	7,7,7	0.53	0	9,9,9	0.67	0
2	TRS	D	240	-	7,7,7	0.54	0	9,9,9	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	A	240	-	-	7/9/9/9	-
2	TRS	C	240	-	-	6/9/9/9	-
2	TRS	B	240	-	-	3/9/9/9	-
2	TRS	D	240	-	-	2/9/9/9	-



There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	240	TRS	C3-C-N	-10.76	80.74	108.17
2	A	240	TRS	C1-C-N	-7.14	89.97	108.17
2	A	240	TRS	C2-C-N	-5.83	93.30	108.17
2	A	240	TRS	C3-C-C1	3.74	120.63	110.66
2	A	240	TRS	C2-C-C1	3.35	119.57	110.66
2	A	240	TRS	C3-C-C2	3.30	119.43	110.66

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	240	TRS	C3-C-C2-O2
2	A	240	TRS	N-C-C3-O3
2	B	240	TRS	C1-C-C3-O3
2	B	240	TRS	N-C-C3-O3
2	C	240	TRS	C1-C-C3-O3
2	C	240	TRS	C2-C-C3-O3
2	C	240	TRS	N-C-C3-O3
2	A	240	TRS	C1-C-C3-O3
2	A	240	TRS	C3-C-C1-O1
2	A	240	TRS	N-C-C2-O2
2	B	240	TRS	C2-C-C3-O3
2	C	240	TRS	C3-C-C1-O1
2	A	240	TRS	C2-C-C1-O1
2	C	240	TRS	C2-C-C1-O1
2	D	240	TRS	C1-C-C3-O3
2	C	240	TRS	N-C-C1-O1
2	A	240	TRS	C2-C-C3-O3
2	D	240	TRS	N-C-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	224/236 (94%)	-0.23	2 (0%) 84 88	16, 31, 60, 106	0
1	B	227/236 (96%)	-0.33	3 (1%) 77 82	18, 30, 56, 93	0
1	C	224/236 (94%)	-0.16	5 (2%) 62 69	18, 36, 66, 123	0
1	D	224/236 (94%)	0.75	28 (12%) 3 5	24, 66, 91, 105	0
All	All	899/944 (95%)	0.01	38 (4%) 36 45	16, 37, 82, 123	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	236	ALA	8.9
1	A	235	THR	8.4
1	D	235	THR	8.3
1	C	236	ALA	8.1
1	A	236	ALA	7.7
1	D	138	ALA	5.1
1	D	129	SER	5.0
1	D	38	GLY	4.5
1	B	236	ALA	4.5
1	D	230	LEU	4.1
1	D	130	PRO	3.8
1	D	232	ALA	3.6
1	C	234	LEU	3.5
1	D	146	ALA	3.4
1	D	95	THR	3.2
1	D	231	ILE	3.2
1	D	107	ALA	3.0
1	D	131	ALA	3.0
1	D	234	LEU	3.0
1	B	9	PRO	3.0
1	C	235	THR	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	12	SER	3.0
1	D	13	GLY	2.9
1	D	83	ALA	2.9
1	D	118	PHE	2.8
1	D	37	PHE	2.7
1	D	40	ALA	2.7
1	D	132	PRO	2.7
1	B	10	SER	2.4
1	D	33	PHE	2.3
1	D	91	ALA	2.2
1	D	15	LEU	2.2
1	C	82	GLN	2.2
1	D	127	ALA	2.2
1	C	232	ALA	2.1
1	D	111	THR	2.1
1	D	144	PHE	2.1
1	D	106	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	ASB	D	18	12/13	0.92	0.12	37,44,50,57	0
1	ASB	C	18	12/13	0.96	0.10	17,25,36,37	0
1	ASB	A	18	12/13	0.98	0.08	16,24,31,34	0
1	ASB	B	18	12/13	0.98	0.08	17,22,26,27	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	TRS	C	240	8/8	0.80	0.35	27,38,45,49	8
2	TRS	D	240	8/8	0.84	0.15	38,45,55,62	0
2	TRS	A	240	8/8	0.93	0.11	30,37,47,47	0
2	TRS	B	240	8/8	0.93	0.12	29,37,40,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.