



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 5, 2023 – 07:21 pm BST

PDB ID : 8CGO  
Title : Structure of human butyrylcholinesterase in complex with N-{{2-(benzyloxy)-3-methoxyphenyl}methyl}-N-[3-(2-fluorophenyl)propyl]cyclobutanamine  
Authors : Brazzolotto, X.; Pidany, F.; Korabecny, J.; Cahlikova, L.; Nachon, F.  
Deposited on : 2023-02-06  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.33  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.33


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	529	 2% 81% 17%
2	B	2	 100%
2	C	2	 50% 50%
3	D	3	 33% 33% 33%
3	E	3	 67% 33%

## 2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 4588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cholinesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	527	4245	2740	713	776	16	0	6	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	17	GLN	ASN	engineered mutation	UNP P06276
A	455	GLN	ASN	engineered mutation	UNP P06276
A	481	GLN	ASN	engineered mutation	UNP P06276
A	486	GLN	ASN	engineered mutation	UNP P06276

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	24	14	1	9	0	0	0
2	C	2	24	14	1	9	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	3	38	22	2	14	0	0	0
3	E	3	38	22	2	14	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
5	A	1	12	6	1	4	1	0	0

- Molecule 6 is {N}-[3-(2-fluorophenyl)propyl]-{N}-[(3-methoxy-2-phenylmethoxy-phenyl)methyl]cyclobutanamine (three-letter code: XB8) (formula:  $C_{28}H_{32}FNO_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
6	A	1	32	28	1	1	2	0	0

- Molecule 7 is N-acetyl-alpha-neuraminic acid (three-letter code: SIA) (formula:  $C_{11}H_{19}NO_9$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
7	A	1	21	11	1	9	0	0

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	A	1	6	3	3	0	0

- Molecule 9 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	O	S	0	0
			5	4	1		
9	A	1	Total	O	S	0	0
			5	4	1		
9	A	1	Total	O	S	0	0
			5	4	1		
9	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	100	Total	O	0	0
			100	100		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cholinesterase




- Molecule 2: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  67% 33%

  
MAG1  
MAG2  
FUC3

## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.39Å 153.39Å 126.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.69 – 2.65 76.69 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.9 (76.69-2.65) 99.9 (76.69-2.65)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.34 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.178 , 0.215 0.183 , 0.217	Depositor DCC
$R_{free}$ test set	1107 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	60.6	Xtrriage
Anisotropy	0.455	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 59.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4588	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, XB8, GOL, FUC, MES, SIA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/4375	0.61	0/5938

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4245	0	4143	67	0
2	B	24	0	22	0	0
2	C	24	0	22	3	0
3	D	38	0	34	2	0
3	E	38	0	34	0	0
4	A	28	0	26	0	0
5	A	12	0	12	3	0
6	A	32	0	0	0	0
7	A	21	0	18	0	0
8	A	6	0	8	1	0
9	A	20	0	0	0	0
10	A	100	0	0	0	0
All	All	4588	0	4319	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:LYS:HE2	8:A:606:GOL:H12	1.74	0.68
1:A:501:LEU:HD11	1:A:508:THR:HG23	1.76	0.65
1:A:518:GLN:HE22	5:A:603:MES:H22	1.61	0.65
1:A:319:VAL:O	1:A:418:PHE:HA	1.98	0.63
1:A:191:SER:HB2	2:C:2:FUC:H4	1.81	0.63
1:A:522:TRP:O	1:A:527:PRO:HD3	2.02	0.58
1:A:308:GLU:OE1	1:A:408:LYS:HE2	2.03	0.58
1:A:77:HIS:O	1:A:81:MET:HG2	2.06	0.54
1:A:254:ARG:HB2	1:A:260:ILE:HG13	1.90	0.53
1:A:348:LYS:HD2	1:A:348:LYS:H	1.75	0.51
1:A:227:PHE:CE1	1:A:303:PRO:HB2	2.44	0.51
1:A:514:LEU:HD12	5:A:603:MES:H52	1.92	0.51
1:A:188:ASN:ND2	2:C:2:FUC:H3	2.26	0.51
1:A:26:THR:HB	1:A:99:ILE:HG12	1.91	0.51
1:A:40:ARG:HD3	1:A:40:ARG:H	1.76	0.50
1:A:378:ASP:O	1:A:384:ASN:ND2	2.38	0.49
1:A:451:GLU:OE1	1:A:453:ARG:HG3	2.13	0.49
1:A:449:PRO:HA	1:A:456:TYR:CD2	2.48	0.49
1:A:515:ARG:HG2	5:A:603:MES:H32	1.95	0.48
1:A:525:PHE:C	1:A:525:PHE:CD1	2.85	0.48
1:A:195:PHE:CB	1:A:221:ILE:HB	2.43	0.48
1:A:525:PHE:C	1:A:525:PHE:HD1	2.17	0.48
1:A:419:TYR:HB3	1:A:490:TRP:CZ2	2.49	0.47
1:A:18:LEU:O	1:A:24:THR:HA	2.15	0.47
1:A:348:LYS:HD2	1:A:348:LYS:N	2.29	0.47
1:A:205:SER:HB3	1:A:222:LEU:HD21	1.97	0.47
1:A:378:ASP:OD2	1:A:380:GLN:N	2.48	0.46
1:A:169:PHE:CZ	1:A:298:PHE:HB2	2.50	0.46
1:A:320:GLY:HA3	1:A:419:TYR:CD2	2.51	0.46
1:A:501:LEU:HD11	1:A:508:THR:CG2	2.44	0.46
1:A:69:ILE:HD12	1:A:84:PRO:HD2	1.98	0.46
1:A:161:GLU:HB3	1:A:261:ILE:HG21	1.98	0.44
1:A:320:GLY:HA3	1:A:419:TYR:CE2	2.53	0.44
1:A:376:TRP:HZ2	1:A:385:TYR:CE1	2.34	0.44
1:A:500:TYR:CZ	1:A:511[A]:MET:HB2	2.52	0.44
1:A:227:PHE:O	1:A:232:ALA:HB3	2.16	0.44
1:A:155:ALA:HB3	1:A:239:ALA:HB1	1.98	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:LEU:HD13	1:A:243:THR:HG21	1.98	0.44
1:A:278:PHE:O	3:D:3:FUC:H4	2.18	0.44
1:A:197:GLU:HG3	1:A:441:GLU:OE2	2.19	0.43
1:A:40:ARG:H	1:A:40:ARG:CD	2.31	0.43
1:A:249:LEU:HG	3:D:3:FUC:H62	2.00	0.42
1:A:374:THR:HG22	1:A:376:TRP:CZ2	2.53	0.42
1:A:156:LEU:HD12	1:A:156:LEU:HA	1.64	0.42
1:A:106:ASN:ND2	2:C:1:NAG:O7	2.52	0.42
1:A:307:LEU:HD11	1:A:408:LYS:HB2	2.01	0.42
1:A:488:THR:HB	1:A:508:THR:HG22	2.01	0.42
1:A:161:GLU:HG3	1:A:258:THR:HG23	2.01	0.42
1:A:280:VAL:HG22	1:A:281:PRO:HD2	2.01	0.42
1:A:304:ASP:OD1	1:A:304:ASP:N	2.51	0.42
1:A:393:VAL:O	1:A:397:ASN:HB2	2.19	0.42
1:A:419:TYR:HB3	1:A:490:TRP:CH2	2.55	0.42
1:A:195:PHE:HB2	1:A:221:ILE:HB	2.02	0.41
1:A:197:GLU:HA	1:A:223:GLN:O	2.20	0.41
1:A:506:GLU:H	1:A:506:GLU:HG3	1.62	0.41
1:A:222:LEU:HD13	1:A:319:VAL:HG22	2.02	0.41
1:A:133:LEU:HB2	1:A:141:VAL:HG21	2.02	0.41
1:A:220:ALA:HB3	1:A:317:ILE:HG22	2.02	0.41
1:A:205:SER:HB3	1:A:222:LEU:CD2	2.51	0.41
1:A:227:PHE:CD1	1:A:227:PHE:C	2.94	0.41
1:A:227:PHE:CD1	1:A:303:PRO:HB2	2.55	0.41
1:A:330:LEU:O	1:A:334:ALA:HB3	2.20	0.41
1:A:335:PRO:HG3	1:A:352:GLU:HB3	2.03	0.41
1:A:424:ARG:NH1	1:A:428:LEU:HD23	2.36	0.40
1:A:447:GLY:HA2	1:A:464:SER:OG	2.22	0.40
1:A:299:LEU:HD12	1:A:299:LEU:HA	1.95	0.40
1:A:131:LYS:HG3	1:A:132:PHE:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	531/529 (100%)	503 (95%)	27 (5%)	1 (0%)	47 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	506	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	458/454 (101%)	438 (96%)	20 (4%)	28 43

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	16	MET
1	A	40	ARG
1	A	48	SER
1	A	51	LYS
1	A	87	ASP
1	A	173	LEU
1	A	195	PHE
1	A	274	LEU
1	A	379	ASP
1	A	452	ARG
1	A	453	ARG
1	A	471	TRP
1	A	489	SER
1	A	495	SER
1	A	508	THR
1	A	511[A]	MET
1	A	511[B]	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	517	GLN
1	A	525	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.99	2 (14%)	17,19,21	0.61	0
2	FUC	B	2	2	10,10,11	1.07	1 (10%)	14,14,16	1.12	2 (14%)
2	NAG	C	1	1,2	14,14,15	0.78	0	17,19,21	0.51	0
2	FUC	C	2	2	10,10,11	2.40	6 (60%)	14,14,16	2.23	5 (35%)
3	NAG	D	1	3,1	14,14,15	0.25	0	17,19,21	0.60	0
3	NAG	D	2	3	14,14,15	0.65	0	17,19,21	0.63	1 (5%)
3	FUC	D	3	3	10,10,11	0.90	0	14,14,16	1.05	1 (7%)
3	NAG	E	1	3,1	14,14,15	0.66	0	17,19,21	0.47	0
3	NAG	E	2	3	14,14,15	0.40	0	17,19,21	0.55	0
3	FUC	E	3	3	10,10,11	1.38	2 (20%)	14,14,16	1.13	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	FUC	B	2	2	-	-	0/1/1/1
2	NAG	C	1	1,2	-	1/6/23/26	0/1/1/1
2	FUC	C	2	2	-	-	0/1/1/1
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	FUC	D	3	3	-	-	0/1/1/1
3	NAG	E	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	FUC	E	3	3	-	-	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	FUC	C4-C3	3.67	1.61	1.52
2	C	2	FUC	O5-C5	2.93	1.49	1.43
2	C	2	FUC	C4-C5	2.80	1.59	1.52
2	C	2	FUC	C1-C2	2.68	1.58	1.52
2	C	2	FUC	O5-C1	2.63	1.47	1.43
2	B	1	NAG	C1-C2	2.58	1.56	1.52
2	B	1	NAG	O5-C1	2.36	1.47	1.43
2	C	2	FUC	C2-C3	2.35	1.56	1.52
3	E	3	FUC	O5-C1	2.28	1.47	1.43
2	B	2	FUC	C1-C2	2.27	1.57	1.52
3	E	3	FUC	C1-C2	2.19	1.57	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	FUC	C1-C2-C3	4.11	114.72	109.67
2	C	2	FUC	O5-C5-C4	3.94	116.58	109.52
2	C	2	FUC	C1-O5-C5	3.47	120.65	112.78
2	C	2	FUC	C3-C4-C5	3.05	114.52	109.77
2	C	2	FUC	C2-C3-C4	3.00	116.09	110.89
3	E	3	FUC	C1-O5-C5	2.85	119.23	112.78
2	B	2	FUC	C1-O5-C5	2.60	118.66	112.78
3	D	3	FUC	O2-C2-C1	2.11	113.47	109.15
2	B	2	FUC	C1-C2-C3	2.02	112.14	109.67
3	D	2	NAG	C1-O5-C5	2.00	114.91	112.19



There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2	NAG	O5-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
3	E	1	NAG	C4-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
2	C	1	NAG	C3-C2-N2-C7



There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	3	FUC	2	0
2	C	2	FUC	2	0
2	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
9	SO4	A	609	-	4,4,4	0.26	0	6,6,6	0.19	0
9	SO4	A	610	-	4,4,4	0.51	0	6,6,6	0.47	0
7	SIA	A	605	-	21,21,21	1.88	4 (19%)	25,31,31	1.51	2 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	GOL	A	606	-	5,5,5	1.05	0	5,5,5	1.02	0
9	SO4	A	607	-	4,4,4	0.31	0	6,6,6	0.53	0
5	MES	A	603	-	12,12,12	2.12	1 (8%)	14,16,16	2.00	3 (21%)
4	NAG	A	602	1	14,14,15	0.62	0	17,19,21	0.63	0
6	XB8	A	604	-	35,35,35	2.10	10 (28%)	45,46,46	0.98	4 (8%)
9	SO4	A	608	-	4,4,4	0.17	0	6,6,6	0.21	0
4	NAG	A	601	1	14,14,15	0.35	0	17,19,21	0.71	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	SIA	A	605	-	-	5/20/38/38	0/1/1/1
8	GOL	A	606	-	-	4/4/4/4	-
5	MES	A	603	-	-	2/6/14/14	0/1/1/1
4	NAG	A	602	1	-	0/6/23/26	0/1/1/1
6	XB8	A	604	-	-	4/17/27/27	0/4/4/4
4	NAG	A	601	1	-	1/6/23/26	0/1/1/1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	603	MES	C8-S	-6.93	1.67	1.77
6	A	604	XB8	C08-C07	5.65	1.60	1.51
7	A	605	SIA	O6-C2	5.06	1.48	1.43
6	A	604	XB8	C08-N09	4.78	1.55	1.47
6	A	604	XB8	C13-C18	3.79	1.44	1.38
7	A	605	SIA	C2-C1	3.73	1.59	1.53
6	A	604	XB8	O02-C03	3.49	1.42	1.37
7	A	605	SIA	C3-C2	2.96	1.55	1.51
7	A	605	SIA	C7-C6	2.93	1.56	1.53
6	A	604	XB8	C24-C07	2.68	1.45	1.40
6	A	604	XB8	C06-C07	2.50	1.43	1.39
6	A	604	XB8	C16-C15	2.34	1.44	1.38
6	A	604	XB8	O25-C24	2.26	1.43	1.39
6	A	604	XB8	C17-C18	2.17	1.42	1.37
6	A	604	XB8	C15-C14	2.04	1.43	1.38

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	603	MES	C5-N4-C3	5.38	120.94	108.83
7	A	605	SIA	O1A-C1-C2	-3.98	117.56	123.59
5	A	603	MES	O3S-S-C8	2.60	109.97	105.77
5	A	603	MES	O1S-S-C8	2.29	109.67	106.92
6	A	604	XB8	C23-C20-N09	-2.23	113.62	120.29
4	A	601	NAG	C1-O5-C5	2.19	115.16	112.19
6	A	604	XB8	C01-O02-C03	-2.19	114.22	117.53
6	A	604	XB8	O02-C03-C04	-2.16	120.66	124.37
6	A	604	XB8	C21-C20-N09	-2.07	114.11	120.29
7	A	605	SIA	C4-C5-N5	2.05	114.44	110.38

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	605	SIA	O6-C6-C7-O7
8	A	606	GOL	C1-C2-C3-O3
6	A	604	XB8	N09-C10-C11-C12
8	A	606	GOL	O1-C1-C2-C3
8	A	606	GOL	O1-C1-C2-O2
8	A	606	GOL	O2-C2-C3-O3
6	A	604	XB8	C07-C08-N09-C20
4	A	601	NAG	O5-C5-C6-O6
6	A	604	XB8	C07-C08-N09-C10
5	A	603	MES	N4-C7-C8-S
5	A	603	MES	C8-C7-N4-C5
7	A	605	SIA	O1A-C1-C2-C3
7	A	605	SIA	C5-C6-C7-O7
7	A	605	SIA	O6-C6-C7-C8
6	A	604	XB8	C11-C10-N09-C20
7	A	605	SIA	O1A-C1-C2-O6


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	606	GOL	1	0
5	A	603	MES	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	527/529 (99%)	0.33	13 (2%) 57 53	50, 64, 99, 130	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	282	TYR	4.1
1	A	4	ILE	3.9
1	A	17	GLN	3.6
1	A	376	TRP	3.6
1	A	3	ASP	3.5
1	A	255	GLU	3.2
1	A	278	PHE	2.5
1	A	259	GLU	2.4
1	A	261	ILE	2.4
1	A	264	LEU	2.2
1	A	12	LYS	2.2
1	A	155	ALA	2.1
1	A	244	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.






Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	D	2	14/15	0.76	0.26	91,109,115,118	0
2	NAG	B	1	14/15	0.84	0.27	102,109,114,114	0
3	NAG	E	2	14/15	0.84	0.30	92,99,111,117	0
2	FUC	C	2	10/11	0.85	0.43	81,84,95,96	0
2	NAG	C	1	14/15	0.86	0.14	68,87,91,96	0
3	FUC	E	3	10/11	0.86	0.52	82,97,102,103	0
2	FUC	B	2	10/11	0.87	0.33	108,121,126,127	0
3	NAG	D	1	14/15	0.89	0.20	99,103,108,109	0
3	FUC	D	3	10/11	0.93	0.26	83,96,102,103	0
3	NAG	E	1	14/15	0.93	0.18	73,79,85,90	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain C:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	602	14/15	0.78	0.26	75,99,105,105	0
9	SO4	A	607	5/5	0.80	0.31	83,89,101,114	0
7	SIA	A	605	21/21	0.82	0.41	72,87,98,103	0
4	NAG	A	601	14/15	0.82	0.24	111,118,122,123	0
6	XB8	A	604	32/32	0.86	0.45	59,69,80,83	0
8	GOL	A	606	6/6	0.87	0.65	71,74,77,77	0
5	MES	A	603	12/12	0.87	0.34	70,87,107,115	0
9	SO4	A	610	5/5	0.88	0.17	101,108,119,123	0
9	SO4	A	608	5/5	0.94	0.13	75,89,101,101	0
9	SO4	A	609	5/5	0.95	0.22	85,88,101,102	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.