



Full wwPDB EM Validation Report ⓘ

Sep 3, 2024 – 12:29 PM EDT

PDB ID : 9CGX
EMDB ID : EMD-45588
Title : Alzheimer's Disease Seeded 0N3R Tau Fibrils
Authors : Duan, P.; Dregni, A.J.; Xu, H.; Changolkar, L.; Lee, V.M.-Y.; Hong, M.
Deposited on : 2024-07-01
Resolution : 2.97 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev112
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	19% 79%
1	B	352	21% 79%
1	C	352	19% 79%
1	D	352	20% 79%
1	E	352	20% 79%
1	F	352	21% 79%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6960 atoms, of which 3540 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform Fetal-tau of Microtubule-associated protein tau.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	A	74	1160	357	590	105	107	1	0	0
1	B	74	1160	357	590	105	107	1	0	0
1	C	74	1160	357	590	105	107	1	0	0
1	D	74	1160	357	590	105	107	1	0	0
1	E	74	1160	357	590	105	107	1	0	0
1	F	74	1160	357	590	105	107	1	0	0

◆ ◆ ◆
F378
R379
E380
ASN
ALA
LYS
ALA
LYS
THR
ASP
HIS
GLY
ALA
GLU
VAL
TYR
LYS
SER
PRO
VAL
VAL
SER
GLY
ASP
THR
SER
PRO
ARG
HIS
LEU
SER
ASN
VAL
SER
THR
GLY
SER
ILE
ASP
MET
VAL
ASP
PRO
GLN
LEU
ALA
THR
LEU
ALA
ASP
GLU
VAL
SER
ALA

LYS
GLN
GLY
LEU

● Molecule 1: Isoform Fetal-tau of Microtubule-associated protein tau

Chain F: 21% 79%

MET
ALA
GLU
PRO
ARG
GLN
GLU
PHE
MET
VAL
SER
MET
GLU
ASP
HIS
ALA
GLY
TYR
THR
GLY
LEU
SER
ASP
GLY
ASP
LYS
ARG
ALA
LYS
ASP
GLN
GLY
ALA
GLY
TYR
THR
MET
HIS
GLN
ALA
ASP
GLN
GLU
GLY
GLY
ASP
THR
ALA
ASP
PRO
PRO
GLY
LEU
LYS
ALA
ILE
GLY
THR
ARG
ASP
PRO
PRO
SER
LEU
GLU
ALA

ALA
GLY
HIS
VAL
THR
GLN
ALA
ARG
MET
VAL
SER
SER
LYS
LYS
ASP
THR
THR
SER
SER
ASP
LYS
LYS
ALA
GLY
ALA
ASP
GLY
LYS
THR
THR
ILE
ALA
THR
PRO
PRO
ARG
GLY
LYS
GLY
GLN
THR
THR
VAL
VAL
VAL
THR
ARG
ILE
THR
PRO
PRO
ALA
LYS
THR
PRO
PRO
SER

PRO
LYS
THR
PRO
PRO
SER
SER
GLY
PRO
PRO
LYS
SER
GLY
ASP
ARG
SER
SER
TYR
SER
SER
SER
PRO
GLY
THR
PRO
GLY
SER
SER
SER
SER
ARG
ARG
THR
PRO
SER
SER
LEU
PRO
PRO
THR
THR
PRO
SER
LEU
PRO
PRO
PRO
THR
ARG
GLU
PRO
LYS
VAL
VAL
VAL
VAL
ARG
THR
THR
PRO
PRO
LYS
SER
SER
SER

ALA
LYS
SER
ARG
LEU
GLN
THR
ALA
GLU
PRO
VAL
VAL
TYR
PRO
MET
PRO
ASP
LEU
LYS
ASN
SER
GLY
VAL
LYS
SER
SER
LYS
LEU
SER
ASN
VAL
VAL
SER
SER
HIS
GLN
PRO
GLY
GLY
GLY
LYS
VAL
VAL
ASP
SER
PRO
GLN
LEU
LEU
VAL
SER
SER
THR
GLY
SER
ILE
GLY
GLY
MET
LYS
VAL
ASP
SER
PRO
GLN
LEU
ALA
THR
LEU
LEU
ALA
THR
LEU
SER
SER
SER
LEU
LYS
L376
T377
F378
R379
E380
ASN
ALA
LYS

ALA
LYS
THR
ASP
HIS
GLY
ALA
GLU
ILE
TYR
LYS
SER
PRO
VAL
VAL
SER
SER
THR
SER
PRO
ARG
HIS
LEU
SER
ASN
VAL
VAL
SER
SER
THR
GLY
SER
ILE
GLY
MET
LYS
VAL
ASP
SER
PRO
GLN
LEU
ALA
THR
LEU
LEU
ALA
THR
LEU
SER
SER
SER
LEU
LYS
L376
T377
F378
R379
E380
ASN
ALA
LYS

4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=179.49°, rise=2.369 Å, axial sym=C1	Depositor
Number of segments used	30112	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{Å}^2$)	44.4	Depositor
Minimum defocus (nm)	200	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.082	Depositor
Minimum map value	-0.044	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	407.03998, 407.03998, 407.03998	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/579	0.96	0/774
1	B	0.53	0/579	0.95	0/774
1	C	0.53	0/579	0.97	0/774
1	D	0.53	0/579	0.95	0/774
1	E	0.53	0/579	1.05	2/774 (0.3%)
1	F	0.52	0/579	0.95	0/774
All	All	0.53	0/3474	0.97	2/4644 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	346	PHE	CB-CG-CD2	-10.99	113.11	120.80
1	E	346	PHE	CB-CG-CD1	6.54	125.37	120.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	346	PHE	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	570	590	589	1	0
1	B	570	590	589	0	0
1	C	570	590	589	1	0
1	D	570	590	589	0	0
1	E	570	590	589	0	0
1	F	570	590	589	0	0
All	All	3420	3540	3534	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:LEU:HD12	1:C:344:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	72/352 (20%)	65 (90%)	4 (6%)	3 (4%)	2 11
1	B	72/352 (20%)	67 (93%)	4 (6%)	1 (1%)	9 35
1	C	72/352 (20%)	66 (92%)	3 (4%)	3 (4%)	2 11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	72/352 (20%)	68 (94%)	2 (3%)	2 (3%)	4	19
1	E	72/352 (20%)	66 (92%)	4 (6%)	2 (3%)	4	19
1	F	72/352 (20%)	67 (93%)	4 (6%)	1 (1%)	9	35
All	All	432/2112 (20%)	399 (92%)	21 (5%)	12 (3%)	6	19

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	379	ARG
1	A	327	ASN
1	A	348	ASP
1	A	379	ARG
1	C	327	ASN
1	C	379	ARG
1	B	348	ASP
1	C	348	ASP
1	D	348	ASP
1	E	348	ASP
1	F	348	ASP
1	E	327	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/285 (23%)	63 (97%)	2 (3%)	35	66
1	B	65/285 (23%)	65 (100%)	0	100	100
1	C	65/285 (23%)	63 (97%)	2 (3%)	35	66
1	D	65/285 (23%)	65 (100%)	0	100	100
1	E	65/285 (23%)	63 (97%)	2 (3%)	35	66
1	F	65/285 (23%)	65 (100%)	0	100	100
All	All	390/1710 (23%)	384 (98%)	6 (2%)	60	82

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	346	PHE
1	A	372	GLU
1	C	346	PHE
1	C	372	GLU
1	E	346	PHE
1	E	372	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-45588. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map




6.1.2 Raw map




The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 202



Y Index: 201




Z Index: 178


6.3.2 Raw map



X Index: 203



Y Index: 199




Z Index: 205


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_45588_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 68 nm³; this corresponds to an approximate mass of 61 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.337 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.337 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.97	-	-
Author-provided FSC curve	2.97	3.34	3.01
Unmasked-calculated*	3.45	6.75	3.49

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.45 differs from the reported value 2.97 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-45588 and PDB model 9CGX. Per-residue inclusion information can be found in section [3](#) on page [4](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

9.4 Atom inclusion [i](#)



At the recommended contour level, 59% of all backbone atoms, 58% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5770	0.5410
A	0.5820	0.5460
B	0.5950	0.5480
C	0.6040	0.5550
D	0.6000	0.5520
E	0.5860	0.5270
F	0.5820	0.5160

