



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 22, 2023 – 04:13 AM EDT

PDB ID : 3CH2  
Title : Crystal Structure Analysis of SERA5E from plasmodium falciparum  
Authors : Smith, B.J.; Malby, R.L.; Colman, P.M.; Clarke, O.B.  
Deposited on : 2008-03-07  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	265	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2326 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine-repeat antigen protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	X	256	2095	1341	329	402	23	0	13	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	X	1	Total	Ca	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	X	230	Total	O	0	2
			230	230		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine-repeat antigen protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.59Å 103.59Å 72.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.46 – 1.80 33.46 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.46-1.80) 100.0 (33.46-1.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.73 (at 1.79Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.166 , 0.198 0.172 , 0.202	Depositor DCC
$R_{free}$ test set	1341 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.7	Xtrriage
Anisotropy	0.027	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 48.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2326	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	X	0.81	1/2192 (0.0%)	0.78	3/2970 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	610	CYS	CB-SG	-5.10	1.73	1.81

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	757[A]	ASP	CB-CG-OD1	6.69	124.32	118.30
1	X	757[B]	ASP	CB-CG-OD1	6.69	124.32	118.30
1	X	715	MET	CG-SD-CE	-5.00	92.19	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2095	0	1899	49	0
2	X	1	0	0	0	0
3	X	230	0	0	5	1
All	All	2326	0	1899	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:827:ASN:O	1:X:828:ASN:HB2	1.70	0.91
1:X:650:GLU:OE2	1:X:688:ILE:HD13	1.72	0.90
1:X:737[B]:LYS:HG3	1:X:739[B]:GLU:HG2	1.53	0.88
1:X:796[B]:GLU:HG3	3:X:967:HOH:O	1.75	0.86
1:X:612:LYS:HD2	1:X:826:MET:HG2	1.62	0.81
1:X:614:TYR:CE2	1:X:828:ASN:HB3	2.17	0.79
1:X:826:MET:O	1:X:826:MET:HE3	1.89	0.72
1:X:826:MET:O	1:X:826:MET:CE	2.38	0.71
1:X:781[A]:SER:OG	1:X:803[A]:MET:CE	2.40	0.69
1:X:689:LEU:CD1	1:X:823:ASP:HB3	2.22	0.69
1:X:781[A]:SER:OG	1:X:803[A]:MET:HE2	1.93	0.68
1:X:824:LEU:HD12	1:X:825:PRO:HD2	1.76	0.67
1:X:626:ASN:ND2	1:X:669:GLY:H	1.92	0.67
1:X:626:ASN:HD21	1:X:669:GLY:H	1.44	0.66
1:X:643:MET:N	1:X:643:MET:HE3	2.12	0.65
1:X:657:ALA:HB2	1:X:680[A]:MET:HG3	1.80	0.64
1:X:689:LEU:HD13	1:X:823:ASP:HB3	1.84	0.59
1:X:827:ASN:O	1:X:828:ASN:CB	2.47	0.57
1:X:614:TYR:CE2	1:X:828:ASN:CB	2.86	0.57
1:X:647:GLN:NE2	3:X:946:HOH:O	2.37	0.56
1:X:600:ALA:HA	1:X:765:ASN:HD21	1.71	0.55
1:X:689:LEU:HD11	1:X:823:ASP:HB3	1.87	0.55
1:X:737[B]:LYS:CG	1:X:739[B]:GLU:HG2	2.33	0.55
1:X:747:SER:O	1:X:752:GLN:NE2	2.40	0.54
1:X:660:ASN:HD21	1:X:680[A]:MET:H	1.55	0.53
1:X:660:ASN:HD21	1:X:680[B]:MET:H	1.55	0.53
1:X:781[A]:SER:OG	1:X:803[A]:MET:HE1	2.09	0.53
1:X:635:ARG:NH2	1:X:644:GLU:OE2	2.44	0.50
1:X:826:MET:O	1:X:827:ASN:CB	2.61	0.47
1:X:642:PRO:HB2	1:X:643:MET:HE3	1.96	0.47
1:X:650:GLU:OE2	1:X:688:ILE:CD1	2.56	0.46
1:X:826:MET:HB3	1:X:826:MET:HE2	1.42	0.46
1:X:781[A]:SER:HG	1:X:803[A]:MET:HE2	1.79	0.46
1:X:594[B]:ASP:OD2	1:X:639:GLY:N	2.49	0.45
1:X:609:ARG:NH1	3:X:964:HOH:O	2.50	0.45
1:X:683:TRP:CZ2	1:X:687:LYS:HA	2.52	0.45
1:X:727[A]:MET:HE3	3:X:987:HOH:O	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:577:ASP:C	1:X:577:ASP:OD1	2.54	0.44
1:X:641:SER:OG	1:X:643:MET:HG2	2.18	0.44
1:X:643:MET:N	1:X:643:MET:CE	2.79	0.43
1:X:614:TYR:CD2	1:X:828:ASN:HB3	2.51	0.43
1:X:609:ARG:NH1	1:X:682:LEU:HD22	2.34	0.43
1:X:657:ALA:HB2	1:X:680[A]:MET:CG	2.48	0.42
1:X:612:LYS:CD	1:X:826:MET:HG2	2.43	0.42
1:X:614:TYR:HE2	1:X:828:ASN:HB3	1.78	0.42
1:X:723:LYS:O	1:X:727[B]:MET:HG3	2.19	0.42
1:X:643:MET:HE3	1:X:643:MET:H	1.85	0.42
1:X:826:MET:O	1:X:826:MET:HE2	2.17	0.41
1:X:680[A]:MET:HE3	3:X:1054:HOH:O	2.19	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:X:957:HOH:O	3:X:995:HOH:O[6_555]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	265/265 (100%)	253 (96%)	11 (4%)	1 (0%)	34 21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	827	ASN



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	222/235 (94%)	214 (96%)	8 (4%)	35 20

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	565[A]	MET
1	X	565[B]	MET
1	X	635	ARG
1	X	643	MET
1	X	684	ASP
1	X	689	LEU
1	X	823	ASP
1	X	826	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	626	ASN
1	X	660	ASN
1	X	740	ASN
1	X	765	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	X	256/265 (96%)	0.61	25 (9%) <b>7</b> <b>5</b>	21, 31, 49, 54	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	652	TYR	6.4
1	X	566	PHE	6.2
1	X	685	ASN	5.2
1	X	823	ASP	5.0
1	X	826	MET	4.9
1	X	632	HIS	4.7
1	X	684	ASP	4.5
1	X	564	ASN	4.5
1	X	828	ASN	4.3
1	X	565[A]	MET	3.9
1	X	827	ASN	3.6
1	X	654	PHE	3.1
1	X	690	HIS	2.8
1	X	744	TYR	2.7
1	X	579	ASN	2.6
1	X	614	TYR	2.4
1	X	733	ILE	2.4
1	X	680[A]	MET	2.3
1	X	593	CYS	2.3
1	X	825	PRO	2.2
1	X	613	GLY	2.1
1	X	677	ASP	2.1
1	X	631	GLU	2.0
1	X	672	CYS	2.0
1	X	675	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	X	1	1/1	0.98	0.24	24,24,24,24	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.