



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 06:56 am BST

PDB ID : 1CKI  
Title : RECOMBINANT CASEIN KINASE I DELTA TRUNCATION MUTANT  
CONTAINING RESIDUES 1-317  
Authors : Longenecker, K.L.; Roach, P.J.; Hurley, T.D.  
Deposited on : 1995-08-25  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	
1	B	317	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4823 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CASEIN KINASE I DELTA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	292	2386	1531	416	424	15	0	0	0
1	B	286	2340	1502	406	417	15	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	56	Total	O	0	0
			56	56		
2	B	41	Total	O	0	0
			41	41		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CASEIN KINASE I DELTA



- Molecule 1: CASEIN KINASE I DELTA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.84Å 115.47Å 60.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.30 24.44 – 2.20	Depositor EDS
% Data completeness (in resolution range)	78.0 (8.00-2.30) 73.7 (24.44-2.20)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 2.19Å)	Xtrriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.193 , 0.279 0.180 , 0.260	Depositor DCC
$R_{free}$ test set	1975 reflections (6.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.8	Xtrriage
Anisotropy	0.417	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 85.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4823	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/2441	0.68	0/3278
1	B	0.41	0/2394	0.66	0/3217
All	All	0.43	0/4835	0.67	0/6495

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2386	0	2383	56	0
1	B	2340	0	2335	44	0
2	A	56	0	0	4	0
2	B	41	0	0	0	0
All	All	4823	0	4718	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:VAL:HG13	1:B:46:LYS:HD3	1.55	0.86

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:ALA:HB1	1:A:304:ARG:HG2	1.61	0.81
1:A:23:PHE:CZ	1:A:51:GLN:HB3	2.25	0.71
1:B:139:MET:CE	1:B:150:VAL:HG22	2.24	0.68
2:A:351:HOH:O	1:B:290:ASP:HB3	1.93	0.68
1:A:53:HIS:O	1:A:57:LYS:HG2	1.96	0.66
1:A:42:LEU:HD23	1:A:82:VAL:HG12	1.81	0.61
1:A:23:PHE:CE2	1:A:51:GLN:HB3	2.35	0.60
1:B:231:GLU:O	1:B:235:GLU:HG3	2.02	0.60
1:B:102:LYS:HD2	1:B:102:LYS:O	2.01	0.59
1:A:181:THR:HG22	1:A:183:ARG:H	1.67	0.59
1:B:139:MET:HE1	1:B:150:VAL:HG22	1.84	0.59
1:B:63:GLN:OE1	1:B:70:THR:HG22	2.02	0.58
1:B:73:TRP:CH2	1:B:75:GLY:HA3	2.38	0.57
1:A:16:ARG:HB3	2:A:362:HOH:O	2.04	0.57
1:B:230:TYR:HA	1:B:233:ILE:HD12	1.86	0.57
1:A:257:LEU:O	1:A:261:ARG:HG2	2.05	0.57
1:B:69:PRO:HG3	1:B:153:ILE:HG22	1.88	0.56
1:A:283:HIS:HA	2:A:337:HOH:O	2.05	0.56
1:B:11:ARG:O	1:B:32:ILE:HG13	2.06	0.56
1:A:16:ARG:HG3	1:A:17:LYS:N	2.19	0.56
1:B:281:LEU:O	1:B:285:GLN:HG2	2.06	0.55
1:A:217:PRO:HG2	1:A:218:TRP:CE3	2.40	0.55
1:A:206:TYR:OH	1:A:261:ARG:NH1	2.40	0.55
1:A:44:CYS:HA	1:A:79:ASP:O	2.06	0.55
1:A:302:ALA:HB1	1:A:304:ARG:CG	2.35	0.55
1:B:161:TYR:CZ	1:B:197:ARG:HD3	2.41	0.55
1:B:42:LEU:CD1	1:B:82:VAL:HG22	2.38	0.54
1:A:26:ILE:HD12	1:A:26:ILE:N	2.22	0.54
1:A:61:MET:SD	1:A:127:PHE:HE1	2.31	0.54
1:A:14:LEU:HD12	1:A:40:ILE:HD12	1.90	0.53
1:A:63:GLN:OE1	1:A:70:THR:HA	2.08	0.53
1:B:182:ALA:HB2	1:B:230:TYR:CZ	2.44	0.53
1:A:178:LEU:HD12	1:A:178:LEU:H	1.73	0.53
1:A:91:SER:HA	1:A:138:LEU:HD23	1.91	0.52
1:B:148:ASN:HB3	1:B:291:TYR:CD1	2.45	0.52
1:A:20:SER:HA	1:A:25:ASP:HA	1.91	0.52
1:B:139:MET:HE2	1:B:150:VAL:HG22	1.91	0.52
1:B:105:LEU:O	1:B:109:LEU:HG	2.10	0.52
1:B:218:TRP:HB2	1:B:233:ILE:HG23	1.92	0.52
1:A:71:ILE:HD13	1:A:85:MET:HB2	1.92	0.51
1:A:131:ASP:HB2	1:A:157:LEU:HD12	1.93	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:THR:HG22	1:B:49:HIS:H	1.75	0.50
1:A:42:LEU:CD2	1:A:82:VAL:HG12	2.41	0.50
1:B:183:ARG:HD2	1:B:184:TYR:CZ	2.45	0.50
1:A:56:SER:O	1:A:60:LYS:HG3	2.12	0.48
1:B:112:ALA:O	1:B:116:ILE:HG13	2.13	0.48
1:A:47:THR:O	1:A:47:THR:HG22	2.12	0.48
1:B:45:VAL:CG1	1:B:46:LYS:HD3	2.36	0.48
1:B:71:ILE:HD13	1:B:85:MET:HB3	1.96	0.48
1:A:23:PHE:O	1:A:41:LYS:HE2	2.13	0.48
1:A:105:LEU:O	1:A:109:LEU:HG	2.14	0.48
1:A:46:LYS:O	1:A:46:LYS:HG3	2.13	0.47
1:B:42:LEU:HD11	1:B:82:VAL:HG22	1.95	0.47
1:A:122:ILE:CD1	1:A:152:ILE:HD13	2.44	0.47
1:B:88:LEU:HB2	1:B:138:LEU:HD23	1.96	0.47
1:A:163:ASP:O	1:A:167:HIS:HA	2.15	0.47
1:B:264:ARG:O	1:B:267:ASP:HB2	2.13	0.47
1:A:174:GLU:HG2	1:B:64:GLY:O	2.15	0.46
1:B:110:LEU:HD13	1:B:282:PHE:CE2	2.50	0.46
1:A:23:PHE:CZ	1:A:156:GLY:HA2	2.51	0.46
1:B:92:LEU:HD21	1:B:139:MET:CE	2.46	0.46
1:A:146:LYS:HA	2:A:356:HOH:O	2.15	0.45
1:A:281:LEU:O	1:A:285:GLN:HG2	2.15	0.45
1:B:23:PHE:CD2	1:B:51:GLN:HB2	2.51	0.45
1:B:69:PRO:HB3	1:B:88:LEU:HD11	1.99	0.45
1:A:23:PHE:CE2	1:A:55:GLU:HB2	2.52	0.44
1:A:61:MET:SD	1:A:127:PHE:CE1	3.10	0.44
1:B:22:SER:OG	1:B:156:GLY:HA3	2.16	0.44
1:B:99:CYS:SG	1:B:298:LEU:HD11	2.58	0.44
1:A:47:THR:HA	1:A:50:PRO:HD3	1.98	0.44
1:A:65:GLY:HA3	1:A:121:TYR:CZ	2.52	0.44
1:A:259:PHE:CD2	1:A:274:LEU:HD21	2.52	0.43
1:A:134:PRO:HG3	1:A:207:VAL:HG13	1.99	0.43
1:B:45:VAL:HG13	1:B:46:LYS:CD	2.39	0.43
1:A:137:PHE:HA	1:A:151:TYR:O	2.18	0.43
1:B:148:ASN:HB3	1:B:291:TYR:CE1	2.54	0.43
1:B:259:PHE:CD1	1:B:274:LEU:HD21	2.54	0.43
1:A:218:TRP:CE2	1:A:237:LYS:HE2	2.53	0.43
1:B:92:LEU:HA	1:B:92:LEU:HD23	1.89	0.43
1:A:145:LYS:HD2	1:A:145:LYS:H	1.84	0.42
1:A:98:PHE:CE2	1:A:298:LEU:HD23	2.53	0.42
1:B:8:VAL:HG11	1:B:82:VAL:HG11	2.02	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:ASP:OD2	1:A:34:ALA:HB3	2.19	0.42
1:A:110:LEU:HD12	1:A:110:LEU:HA	1.93	0.41
1:B:179:THR:O	1:B:179:THR:HG23	2.20	0.41
1:A:45:VAL:HG12	1:A:45:VAL:O	2.19	0.41
1:A:152:ILE:HG23	1:A:152:ILE:O	2.21	0.41
1:A:122:ILE:HD11	1:A:152:ILE:HG21	2.02	0.41
1:A:46:LYS:C	1:A:48:LYS:H	2.24	0.41
1:A:209:MET:O	1:A:213:LEU:HB2	2.21	0.41
1:B:120:GLU:HG3	1:B:271:TYR:CD2	2.56	0.41
1:A:101:ARG:NH2	1:A:210:TYR:OH	2.54	0.40
1:A:44:CYS:SG	1:A:46:LYS:HG2	2.61	0.40
1:A:53:HIS:CE1	1:A:54:ILE:HG13	2.57	0.40
1:B:128:ILE:HG23	1:B:161:TYR:CD2	2.56	0.40
1:A:298:LEU:HB2	1:A:303:SER:HB3	2.04	0.40
1:B:133:LYS:HB2	1:B:134:PRO:HD2	2.03	0.40
1:B:163:ASP:O	1:B:167:HIS:HA	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/317 (91%)	263 (91%)	23 (8%)	2 (1%)	22	26
1	B	282/317 (89%)	258 (92%)	24 (8%)	0	100	100
All	All	570/634 (90%)	521 (91%)	47 (8%)	2 (0%)	34	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	47	THR
1	A	45	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/276 (93%)	240 (94%)	16 (6%)	18	24
1	B	252/276 (91%)	244 (97%)	8 (3%)	39	54
All	All	508/552 (92%)	484 (95%)	24 (5%)	26	37

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ARG
1	A	11	ARG
1	A	28	LEU
1	A	63	GLN
1	A	77	GLU
1	A	81	ASN
1	A	85	MET
1	A	110	LEU
1	A	159	LYS
1	A	165	ARG
1	A	177	ASN
1	A	213	LEU
1	A	229	LYS
1	A	272	SER
1	A	276	GLN
1	A	299	LYS
1	B	42	LEU
1	B	79	ASP
1	B	102	LYS
1	B	110	LEU
1	B	243	GLU
1	B	245	LEU
1	B	258	ASN
1	B	291	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	GLN
1	A	81	ASN
1	A	276	GLN
1	B	51	GLN
1	B	53	HIS
1	B	175	ASN
1	B	177	ASN
1	B	258	ASN
1	B	280	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/317 (92%)	0.09	25 (8%) 10 14	9, 30, 78, 98	1 (0%)
1	B	286/317 (90%)	0.01	12 (4%) 36 43	14, 36, 69, 91	0
All	All	578/634 (91%)	0.05	37 (6%) 19 25	9, 35, 72, 98	1 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	303	SER	16.9
1	A	23	PHE	5.6
1	A	10	ASN	5.4
1	B	230	TYR	5.2
1	A	49	HIS	5.1
1	A	11	ARG	4.8
1	A	304	ARG	4.7
1	A	47	THR	4.6
1	B	167	HIS	4.2
1	A	178	LEU	3.9
1	A	302	ALA	3.7
1	B	165	ARG	3.7
1	A	165	ARG	3.6
1	B	291	TYR	3.5
1	B	232	ARG	3.5
1	A	44	CYS	3.5
1	A	164	ALA	3.4
1	B	175	ASN	3.4
1	B	179	THR	3.2
1	B	164	ALA	3.2
1	A	79	ASP	3.1
1	A	80	TYR	3.1
1	A	33	ALA	3.0
1	A	177	ASN	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	176	LYS	2.8
1	B	49	HIS	2.7
1	A	167	HIS	2.6
1	A	73	TRP	2.5
1	A	45	VAL	2.4
1	B	264	ARG	2.4
1	A	50	PRO	2.4
1	B	23	PHE	2.4
1	A	32	ILE	2.2
1	B	244	VAL	2.2
1	A	34	ALA	2.2
1	A	1	MET	2.1
1	A	77	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.