



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2023 – 10:40 PM EST

PDB ID : 1CL1
Title : CYSTATHIONINE BETA-LYASE (CBL) FROM ESCHERICHIA COLI
Authors : Clausen, T.; Huber, R.; Messerschmidt, A.
Deposited on : 1997-09-02
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	395	
1	B	395	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9180 atoms, of which 2558 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYSTATHIONINE BETA-LYASE.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	S			
1	A	391	3651	1905	652	518	561	1	14	0	0	0
1	B	392	3644	1899	654	520	556	1	14	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	210	LLP	LYS	modified residue	UNP P06721
B	210	LLP	LYS	modified residue	UNP P06721

- Molecule 2 is BICARBONATE ION (three-letter code: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	5	1	1	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	B	1	5	1	1	3	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
3	A	307	921	614	307	0	0
3	B	318	954	636	318	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: CYSTATHIONINE BETA-LYASE

Chain A:  86% 13% ..



- Molecule 1: CYSTATHIONINE BETA-LYASE

Chain B:  88% 11% ..



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	60.90Å 154.70Å 152.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.83	Depositor
% Data completeness (in resolution range)	92.6 (8.00-1.83)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.151 , 0.221	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	9180	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, BCT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/3037	0.63	1/4122 (0.0%)
1	B	0.39	0/3028	0.63	0/4113
All	All	0.39	0/6065	0.63	1/8235 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	207	ALA	N-CA-C	-5.16	97.06	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2999	652	2943	34	1
1	B	2990	654	2926	28	3
2	A	4	1	1	0	0
2	B	4	1	1	0	0
3	A	307	614	0	3	5
3	B	318	636	0	6	4
All	All	6622	2558	5871	62	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225:CYS:SG	1:B:233:LEU:HD23	2.18	0.84
1:B:18:LYS:HA	1:B:18:LYS:HE2	1.63	0.80
1:A:225:CYS:SG	1:A:233:LEU:HD23	2.25	0.75
1:B:57:GLY:HA2	3:B:508:HOH:O	1.89	0.71
1:A:64:HIS:HE1	1:A:244:VAL:O	1.76	0.69
1:B:21:LEU:HD12	1:B:27:VAL:HG11	1.77	0.67
1:A:391:PHE:O	1:A:395:VAL:HG23	1.95	0.66
1:A:5:LYS:HB3	3:A:609:HOH:O	1.98	0.63
1:A:61:THR:H	1:A:64:HIS:HD2	1.45	0.63
1:B:344:GLU:HB2	3:B:640:HOH:O	1.99	0.62
1:B:288:HIS:HD2	1:B:290:ALA:H	1.47	0.62
1:B:61:THR:H	1:B:64:HIS:CD2	2.17	0.62
1:A:317:LYS:HD2	1:A:319:LEU:HD23	1.82	0.62
1:A:61:THR:H	1:A:64:HIS:CD2	2.18	0.62
1:A:288:HIS:HD2	1:A:290:ALA:H	1.49	0.60
1:A:289:PRO:O	1:A:297:HIS:HD2	1.84	0.59
1:B:64:HIS:HE1	1:B:244:VAL:O	1.86	0.58
1:B:208:ALA:HB3	1:B:220:ILE:HG23	1.86	0.58
1:A:282:GLN:NE2	1:A:395:VAL:HG22	2.19	0.58
1:B:61:THR:H	1:B:64:HIS:HD2	1.50	0.57
1:B:34:LEU:HD11	3:B:508:HOH:O	2.05	0.57
1:A:285:ARG:HE	1:A:287:ASN:HD21	1.55	0.54
1:B:280:HIS:CD2	1:B:282:GLN:H	2.26	0.54
1:A:211:TYR:HB2	3:A:457:HOH:O	2.08	0.53
1:A:280:HIS:HD2	1:A:282:GLN:H	1.55	0.53
1:B:17:LYS:HB3	1:B:17:LYS:NZ	2.25	0.52
1:B:280:HIS:HD2	1:B:282:GLN:H	1.59	0.51
1:A:15:ARG:HD2	1:A:24:VAL:O	2.12	0.50
1:B:210:LLP:HE3	1:B:339:SER:HA	1.94	0.50
1:B:189:ALA:HB1	1:B:193:LEU:HB2	1.95	0.49
1:A:298:GLU:CD	1:A:298:GLU:H	2.16	0.49
1:A:355:ILE:HG22	1:A:364:ILE:HD11	1.94	0.49
1:A:208:ALA:HB3	1:A:220:ILE:HG23	1.94	0.48
1:B:130:SER:HB2	3:B:490:HOH:O	2.15	0.47
1:A:112:GLU:HB3	1:A:113:PRO:HD3	1.96	0.47
1:A:136:ILE:HD13	1:A:140:ILE:HA	1.95	0.47
1:A:395:VAL:HG21	3:A:661:HOH:O	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:61:THR:HG22	3:B:508:HOH:O	2.15	0.47
1:B:108:ASN:HB2	1:B:360:PRO:HD2	1.97	0.46
1:A:95:LEU:HD12	1:A:98:ILE:HD11	1.96	0.46
1:B:112:GLU:HB3	1:B:113:PRO:HD3	1.99	0.45
1:A:48:ASN:HB2	1:A:53:GLU:HG3	1.99	0.45
1:A:251:ILE:HD13	1:A:251:ILE:HA	1.88	0.44
1:A:282:GLN:NE2	1:A:391:PHE:HB3	2.32	0.44
1:A:20:THR:HB	1:A:23:ALA:O	2.17	0.43
1:B:280:HIS:HA	1:B:281:PRO:HD3	1.85	0.43
1:B:392:ALA:HA	1:B:395:VAL:HG22	1.99	0.43
1:B:211:TYR:CE2	1:B:308:SER:HB2	2.54	0.43
1:A:392:ALA:HA	1:A:395:VAL:HG23	2.02	0.42
1:A:131:TRP:CD1	1:A:131:TRP:N	2.87	0.42
1:A:131:TRP:CG	1:A:360:PRO:HB2	2.55	0.42
1:A:65:PHE:HA	1:A:68:GLN:HE21	1.84	0.41
1:A:351:GLN:HA	1:A:351:GLN:OE1	2.20	0.41
1:A:136:ILE:HG12	1:A:139:ASP:HB2	2.01	0.41
1:B:24:VAL:HG22	3:B:709:HOH:O	2.20	0.41
1:B:338:TYR:O	1:B:339:SER:CB	2.68	0.41
1:A:121:ILE:O	1:A:124:LYS:HG2	2.21	0.41
1:A:131:TRP:CD1	1:A:360:PRO:HB2	2.56	0.41
1:B:18:LYS:HA	1:B:18:LYS:CE	2.42	0.41
1:A:359:ARG:HG3	1:A:364:ILE:CD1	2.52	0.40
1:B:330:ASN:HB2	1:B:393:ARG:NE	2.37	0.40
1:B:70:ALA:O	1:B:74:LEU:HG	2.22	0.40

All (11) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:592:HOH:H1	3:A:592:HOH:H2[3_555]	0.02	1.58
3:A:592:HOH:O	3:A:592:HOH:O[3_555]	0.86	1.34
3:B:624:HOH:O	3:B:624:HOH:O[3_555]	1.00	1.20
1:B:58:ARG:HH22	1:B:111:TYR:HH[3_555]	1.22	0.38
1:B:124:LYS:HZ3	3:B:561:HOH:H2[3_555]	1.24	0.36
1:B:15:ARG:HH12	3:A:627:HOH:H2[3_555]	1.27	0.33
1:A:100:GLN:HE22	1:A:100:GLN:HE22[3_555]	1.30	0.30
3:A:484:HOH:O	3:A:592:HOH:H1[3_555]	1.45	0.15
3:B:550:HOH:O	3:B:567:HOH:H1[3_555]	1.45	0.15
3:B:550:HOH:O	3:B:567:HOH:H2[3_555]	1.45	0.15
3:A:443:HOH:O	3:A:462:HOH:H2[3_555]	1.58	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/395 (98%)	375 (97%)	13 (3%)	0	100	100
1	B	389/395 (98%)	380 (98%)	9 (2%)	0	100	100
All	All	777/790 (98%)	755 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/321 (97%)	306 (98%)	5 (2%)	62	49
1	B	308/321 (96%)	303 (98%)	5 (2%)	62	49
All	All	619/642 (96%)	609 (98%)	10 (2%)	62	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	186	ASN
1	A	267	HIS
1	A	279	GLU
1	A	317	LYS
1	A	364	ILE
1	B	18	LYS
1	B	186	ASN
1	B	239	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	267	HIS
1	B	395	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	A	68	GLN
1	A	143	HIS
1	A	186	ASN
1	A	265	GLN
1	A	280	HIS
1	A	282	GLN
1	A	287	ASN
1	A	288	HIS
1	A	297	HIS
1	B	51	ASN
1	B	64	HIS
1	B	68	GLN
1	B	69	GLN
1	B	100	GLN
1	B	186	ASN
1	B	280	HIS
1	B	288	HIS
1	B	297	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	210	1	23,24,25	1.61	4 (17%)	25,32,34	2.03	7 (28%)
1	LLP	B	210	1	23,24,25	1.60	7 (30%)	25,32,34	2.04	7 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	210	1	-	4/16/17/19	0/1/1/1
1	LLP	B	210	1	-	2/16/17/19	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	210	LLP	C4-C5	4.65	1.47	1.42
1	B	210	LLP	C4-C5	3.55	1.46	1.42
1	B	210	LLP	C4'-NZ	2.96	1.37	1.27
1	B	210	LLP	C3-C2	2.81	1.43	1.40
1	B	210	LLP	C2-N1	2.35	1.38	1.33
1	A	210	LLP	C3-C2	2.34	1.43	1.40
1	B	210	LLP	C4-C3	-2.30	1.37	1.40
1	A	210	LLP	P-OP3	-2.25	1.46	1.54
1	B	210	LLP	C2'-C2	2.25	1.54	1.50
1	A	210	LLP	C4'-NZ	2.18	1.34	1.27
1	B	210	LLP	P-OP3	-2.04	1.47	1.54

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	210	LLP	C2'-C2-C3	5.77	128.01	120.89
1	B	210	LLP	C2'-C2-C3	5.44	127.60	120.89
1	A	210	LLP	C6-N1-C2	3.61	125.85	119.17
1	B	210	LLP	C5-C6-N1	-3.37	118.20	123.82
1	B	210	LLP	C6-N1-C2	3.32	125.31	119.17
1	A	210	LLP	C3-C2-N1	-3.26	116.56	120.77
1	B	210	LLP	C3-C2-N1	-3.15	116.70	120.77
1	A	210	LLP	C5-C6-N1	-3.13	118.61	123.82
1	B	210	LLP	O3-C3-C4	2.72	126.90	119.60
1	A	210	LLP	O3-C3-C4	2.67	126.78	119.60
1	A	210	LLP	CE-NZ-C4'	2.56	126.77	118.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	210	LLP	O3-C3-C2	-2.38	112.30	117.49
1	B	210	LLP	O3-C3-C2	-2.28	112.52	117.49
1	B	210	LLP	CE-NZ-C4'	2.23	125.74	118.90

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	210	LLP	C-CA-CB-CG
1	A	210	LLP	O-C-CA-CB
1	B	210	LLP	O-C-CA-CB
1	B	210	LLP	C4-C4'-NZ-CE
1	A	210	LLP	CA-CB-CG-CD
1	A	210	LLP	C4-C4'-NZ-CE

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	210	LLP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BCT	A	402	-	2,3,3	0.77	0	2,3,3	0.54	0
2	BCT	B	403	-	2,3,3	0.89	0	2,3,3	0.60	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.