



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2023 – 03:11 pm GMT

PDB ID : 8CLC
Title : Tubulin (T2R-TTL) complex
Authors : Wranik, M.; Bertrand, Q.; Kepa, M.W.; Weinert, T.; Steinmetz, M.; Standfuss, J.
Deposited on : 2023-02-16
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	2808 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	4% (poor fit), 99% (0-1 outliers), 1% (2-3 outliers), 0% (4+ outliers)
1	C	440	0% (poor fit), 98% (0-1 outliers), 2% (2-3 outliers), 0% (4+ outliers)
2	B	430	4% (poor fit), 98% (0-1 outliers), 2% (2-3 outliers), 0% (4+ outliers)
2	D	430	6% (poor fit), 97% (0-1 outliers), 3% (2-3 outliers), 0% (4+ outliers)
3	E	123	11% (poor fit), 97% (0-1 outliers), 2% (2-3 outliers), 0% (4+ outliers)
4	F	332	17% (poor fit), 95% (0-1 outliers), 5% (2-3 outliers), 0% (4+ outliers)

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 17569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	439	Total	C	N	O	S	0	1	0
			3436	2173	584	656	23			
1	C	440	Total	C	N	O	S	0	1	0
			3440	2177	584	656	23			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	425	Total	C	N	O	S	2	2	0
			3360	2111	574	649	26			
2	D	427	Total	C	N	O	S	2	1	0
			3358	2107	574	651	26			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	123	Total	C	N	O	S	0	0	0
			1014	625	183	201	5			

- Molecule 4 is a protein called Tubulin-Tyrosine Ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	F	332	Total	C	N	O	S	0	1	0
			2714	1740	464	495	15			

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		
7	C	1	Total	Ca	0	0
			1	1		

- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			
8	B	1	Total	28	10	5	11	2	0	0
8	D	1	Total	28	10	5	11	2	0	0

- Molecule 9 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			
9	F	1	Total	31	11	5	12	3	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	15	Total O 15 15	0	0
10	B	21	Total O 21 21	0	0
10	C	38	Total O 38 38	0	0
10	D	3	Total O 3 3	0	0
10	E	2	Total O 2 2	0	0
10	F	12	Total O 12 12	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain




- Molecule 2: Tubulin beta-2B chain



- Molecule 3: Stathmin-4



- Molecule 4: Tubulin-Tyrosine Ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.53Å 160.48Å 181.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.20 – 2.70 15.20 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (15.20-2.70) 99.9 (15.20-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.20_4487	Depositor
R, R_{free}	0.186 , 0.234 0.190 , 0.231	Depositor DCC
R_{free} test set	627 reflections (0.74%)	wwPDB-VP
Wilson B-factor (Å ²)	66.6	Xtrriage
Anisotropy	0.008	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 79.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17569	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG, ACP, GTP, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3514	0.50	0/4770
1	C	0.26	0/3521	0.47	0/4780
2	B	0.28	0/3437	0.48	0/4655
2	D	0.27	0/3431	0.47	0/4645
3	E	0.23	0/1022	0.41	0/1356
4	F	0.27	0/2777	0.49	0/3750
All	All	0.27	0/17702	0.48	0/23956

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	284	ARG	Sidechain
1	C	221	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3436	0	3343	0	0
1	C	3440	0	3353	0	0
2	B	3360	0	3238	0	0
2	D	3358	0	3232	0	0
3	E	1014	0	1029	0	0
4	F	2714	0	2682	0	0
5	A	32	0	12	0	0
5	C	32	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0
8	B	28	0	12	0	0
8	D	28	0	12	0	0
9	F	31	0	14	0	0
10	A	15	0	0	0	0
10	B	21	0	0	0	0
10	C	38	0	0	0	0
10	D	3	0	0	0	0
10	E	2	0	0	0	0
10	F	12	0	0	0	0
All	All	17569	0	16939	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). Clashscore could not be calculated for this entry.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/440 (100%)	423 (97%)	15 (3%)	0	100	100
1	C	439/440 (100%)	425 (97%)	14 (3%)	0	100	100
2	B	423/430 (98%)	409 (97%)	14 (3%)	0	100	100
2	D	423/430 (98%)	406 (96%)	17 (4%)	0	100	100
3	E	119/123 (97%)	117 (98%)	2 (2%)	0	100	100
4	F	321/332 (97%)	302 (94%)	19 (6%)	0	100	100
All	All	2163/2195 (98%)	2082 (96%)	81 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/371 (100%)	366 (99%)	5 (1%)	69	87
1	C	372/371 (100%)	365 (98%)	7 (2%)	57	82
2	B	369/371 (100%)	364 (99%)	5 (1%)	67	86
2	D	367/371 (99%)	359 (98%)	8 (2%)	52	79
3	E	110/110 (100%)	106 (96%)	4 (4%)	35	64
4	F	298/297 (100%)	282 (95%)	16 (5%)	22	47
All	All	1887/1891 (100%)	1842 (98%)	45 (2%)	49	77

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	GLU
1	A	124	LYS
1	A	221	ARG
1	A	262	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	367	ASP
2	B	139	HIS
2	B	241	CYS
2	B	284	ARG
2	B	341	SER
2	B	346	TRP
1	C	48	SER
1	C	71	GLU
1	C	120	ASP
1	C	123	ARG
1	C	170	SER
1	C	276	ILE
1	C	339	ARG
2	D	116	ASP
2	D	120	ASP
2	D	139	HIS
2	D	207	GLU
2	D	215	ARG
2	D	369	ARG
2	D	400	ARG
2	D	423	SER
3	E	6	MET
3	E	107	SER
3	E	122	ARG
3	E	126	LYS
4	F	19	ARG
4	F	31	ARG
4	F	87	LEU
4	F	88	SER
4	F	91	CYS
4	F	131	PHE
4	F	139	ARG
4	F	145	ASN
4	F	163	SER
4	F	164	SER
4	F	197	ARG
4	F	211	TYR
4	F	220	VAL
4	F	242	ASN
4	F	343	TYR
4	F	361	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	HIS
1	A	285	GLN
1	A	301	GLN
2	B	424	ASN
1	C	133	GLN
1	C	256	GLN
2	D	50	ASN
2	D	101	ASN
3	E	51	GLN
3	E	91	ASN
4	F	348	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	GDP	B	501	6	24,30,30	1.00	2 (8%)	30,47,47	0.65	1 (3%)
5	GTP	C	501	6	26,34,34	1.01	2 (7%)	32,54,54	0.68	1 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	GDP	D	501	-	24,30,30	0.98	2 (8%)	30,47,47	0.68	1 (3%)
9	ACP	F	401	-	27,33,33	0.91	1 (3%)	32,52,52	0.82	2 (6%)
5	GTP	A	501	6	26,34,34	1.01	2 (7%)	32,54,54	0.69	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GDP	B	501	6	-	3/12/32/32	0/3/3/3
5	GTP	C	501	6	-	6/18/38/38	0/3/3/3
8	GDP	D	501	-	-	2/12/32/32	0/3/3/3
9	ACP	F	401	-	-	4/15/38/38	0/3/3/3
5	GTP	A	501	6	-	5/18/38/38	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	501	GTP	C5-C6	-2.72	1.41	1.47
8	D	501	GDP	C5-C6	-2.71	1.41	1.47
8	B	501	GDP	C5-C6	-2.68	1.42	1.47
5	C	501	GTP	C5-C6	-2.67	1.42	1.47
9	F	401	ACP	PB-O2B	-2.46	1.50	1.56
5	C	501	GTP	C8-N7	-2.16	1.31	1.35
8	B	501	GDP	C8-N7	-2.14	1.31	1.35
5	A	501	GTP	C8-N7	-2.10	1.31	1.35
8	D	501	GDP	C8-N7	-2.08	1.31	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	F	401	ACP	C5-C6-N6	2.27	123.81	120.35
9	F	401	ACP	O1G-PG-C3B	-2.11	106.69	111.24
5	C	501	GTP	O6-C6-C5	2.07	128.41	124.37
5	A	501	GTP	O6-C6-C5	2.07	128.41	124.37
8	D	501	GDP	O6-C6-C5	2.06	128.39	124.37
8	B	501	GDP	O6-C6-C5	2.03	128.33	124.37

There are no chirality outliers.


All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	501	GTP	C5'-O5'-PA-O1A
5	C	501	GTP	PB-O3B-PG-O3G
5	C	501	GTP	C5'-O5'-PA-O1A
8	B	501	GDP	C5'-O5'-PA-O1A
8	B	501	GDP	C5'-O5'-PA-O2A
8	D	501	GDP	C5'-O5'-PA-O3A
8	D	501	GDP	C5'-O5'-PA-O2A
9	F	401	ACP	PB-C3B-PG-O1G
5	A	501	GTP	PB-O3B-PG-O2G
5	A	501	GTP	PB-O3B-PG-O3G
5	A	501	GTP	C5'-O5'-PA-O3A
5	C	501	GTP	C5'-O5'-PA-O3A
9	F	401	ACP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O2A
5	C	501	GTP	C5'-O5'-PA-O2A
9	F	401	ACP	PB-C3B-PG-O2G
8	B	501	GDP	C5'-O5'-PA-O3A
5	C	501	GTP	PB-O3A-PA-O1A
5	C	501	GTP	PB-O3A-PA-O2A
9	F	401	ACP	C5'-O5'-PA-O1A


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
4	F	5
3	E	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	E	28:SER	C	44:ASP	N	32.44
1	F	362:ALA	C	372:THR	N	16.72
1	F	155:ALA	C	162:ILE	N	13.81
1	F	102:PRO	C	125:THR	N	12.29
1	F	246:GLN	C	256:TYR	N	11.47

Continued on next page...

Continued from previous page...

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	F	231:ALA	C	235:ASP	N	6.17

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/440 (99%)	-0.06	18 (4%) 37 36	39, 68, 116, 239	0
1	C	440/440 (100%)	-0.21	6 (1%) 75 77	38, 55, 96, 133	0
2	B	425/430 (98%)	-0.00	17 (4%) 38 37	40, 63, 111, 163	4 (0%)
2	D	427/430 (99%)	0.20	26 (6%) 21 20	41, 80, 131, 181	3 (0%)
3	E	123/123 (100%)	0.69	14 (11%) 5 4	53, 80, 144, 173	0
4	F	332/332 (100%)	0.79	58 (17%) 1 1	57, 95, 172, 218	0
All	All	2186/2195 (99%)	0.14	139 (6%) 19 18	38, 72, 135, 239	7 (0%)

All (139) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	285	ALA	11.1
4	F	372	THR	10.0
4	F	154	GLY	8.9
4	F	155	ALA	8.8
3	E	143	ALA	8.3
4	F	137	ARG	8.3
4	F	142	ARG	7.6
1	A	439	SER	7.5
4	F	244	CYS	6.7
4	F	133	ALA	6.7
1	A	438	ASP	6.6
4	F	153	ALA	6.2
3	E	142	GLU	6.1
4	F	236	LYS	6.0
1	A	282	TYR	5.8
2	D	179	ASP	5.8
4	F	132	LEU	5.6
4	F	242	ASN	5.4
3	E	139	LEU	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	F	143	GLU	5.0
4	F	140	GLU	4.9
4	F	152	SER	4.8
4	F	125	THR	4.7
2	B	80	SER	4.7
3	E	48	GLU	4.4
4	F	176	GLN	4.4
2	D	416	MET	4.3
4	F	171	ASP	4.3
4	F	381	HIS	4.1
4	F	175	GLU	4.1
3	E	28	SER	4.1
2	D	280	SER	4.0
3	E	138	GLU	4.0
1	C	440	VAL	3.8
4	F	245	ILE	3.7
2	D	278	ARG	3.7
3	E	141	GLU	3.6
2	D	441	ASP	3.6
4	F	131	PHE	3.6
2	D	295	MET	3.6
4	F	141	GLY	3.6
2	D	277	SER	3.6
3	E	128	LYS	3.6
2	D	215	ARG	3.5
2	B	59	ASN	3.5
4	F	168	GLU	3.5
4	F	229	ASN	3.5
2	D	281	GLN	3.5
4	F	135	TYR	3.5
1	A	280	LYS	3.4
2	D	401	ARG	3.3
2	B	370	GLY	3.3
2	D	276	THR	3.3
4	F	139	ARG	3.2
1	A	281	ALA	3.2
4	F	21	LEU	3.2
4	F	90	SER	3.2
1	A	163	LYS	3.2
2	B	321	GLY	3.2
2	D	400	ARG	3.1
2	B	57	THR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	F	243	HIS	3.1
2	B	56	ALA	3.1
2	B	438	ALA	3.1
4	F	174	ASP	3.1
3	E	140	LYS	3.1
2	B	39	ASP	3.0
4	F	89	GLU	3.0
1	A	262	TYR	3.0
2	D	82	PRO	3.0
2	D	406	HIS	3.0
2	D	76	ASP	3.0
4	F	130	VAL	3.0
1	A	283	HIS	2.9
2	B	369	ARG	2.9
4	F	256	TYR	2.9
2	B	85	GLN	2.9
1	A	437	VAL	2.9
4	F	259	GLY	2.9
4	F	333	ASN	2.9
4	F	177	GLY	2.9
4	F	167	SER	2.8
1	A	365	GLY	2.8
2	D	85	GLN	2.8
2	D	116	ASP	2.8
2	D	80	SER	2.7
4	F	46	ARG	2.7
1	A	77	GLU	2.7
4	F	136	ASN	2.7
2	D	96	GLN	2.7
4	F	194	PRO	2.7
2	B	282	GLN	2.7
4	F	180	HIS	2.7
4	F	163	SER	2.7
4	F	165	GLU	2.6
4	F	91	CYS	2.6
2	B	416	MET	2.6
3	E	44	ASP	2.6
4	F	178	GLN	2.6
2	D	37	HIS	2.6
1	C	249	ASN	2.6
1	C	349	THR	2.5
4	F	379	HIS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	439	THR	2.5
3	E	125	GLU	2.5
4	F	258	GLU	2.5
1	A	123	ARG	2.5
2	B	284	ARG	2.5
1	A	279	GLU	2.5
1	A	130	THR	2.5
2	D	2[A]	ARG	2.4
2	D	38	GLY	2.4
4	F	126	ASP	2.4
4	F	235	ASP	2.4
4	F	362	ALA	2.4
1	C	357	TYR	2.3
1	A	215	ARG	2.3
1	A	337	THR	2.3
1	A	43	GLY	2.3
3	E	63	TYR	2.3
4	F	151	SER	2.3
2	D	372	LYS	2.3
4	F	241	THR	2.3
2	B	209	LEU	2.3
2	B	130	ASP	2.3
3	E	122	ARG	2.2
4	F	129	GLU	2.2
4	F	323	GLU	2.1
2	B	440	ALA	2.1
2	D	221	THR	2.1
1	C	285	GLN	2.1
2	D	56	ALA	2.1
3	E	106	GLU	2.1
4	F	257	GLU	2.0
1	A	85	GLN	2.0
4	F	340	GLN	2.0
4	F	273	ASP	2.0
1	C	337	THR	2.0
4	F	196	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	ACP	F	401	31/31	0.92	0.13	92,115,145,162	0
8	GDP	D	501	28/28	0.94	0.16	66,76,101,113	0
7	CA	A	503	1/1	0.94	0.06	89,89,89,89	0
6	MG	A	502	1/1	0.96	0.08	44,44,44,44	0
7	CA	C	503	1/1	0.96	0.04	72,72,72,72	0
8	GDP	B	501	28/28	0.97	0.15	34,50,64,71	0
5	GTP	A	501	32/32	0.97	0.15	37,50,68,79	0
6	MG	B	502	1/1	0.97	0.24	57,57,57,57	0
6	MG	C	502	1/1	0.98	0.09	45,45,45,45	0
5	GTP	C	501	32/32	0.98	0.12	36,46,62,72	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP F 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GDP D 501:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.