

# Full wwPDB X-ray Structure Validation Report (i)

### Aug 17, 2020 – 12:54 PM BST

PDB ID	:	6CMG
Title	:	Crystal Structure of the Hendra Virus Attachment G Glycoprotein Bound to
		a Potent Cross-Reactive Neutralizing Human Monoclonal Antibody m102.3
Authors	:	Xu, K.; Nikolov, D.
Deposited on	:	2018-03-05
Resolution	:	2.70  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
$\operatorname{Refmac}$	:	5.8.0158
$\operatorname{CCP4}$	:	7.0.044  (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	Similar resolution (#Entries, resolution range(Å))				
R <sub>free</sub>	130704	2808 (2.70-2.70)				
Clashscore	141614	3122(2.70-2.70)				
Ramachandran outliers	138981	3069(2.70-2.70)				
Sidechain outliers	138945	3069(2.70-2.70)				
RSRZ outliers	127900	2737 (2.70-2.70)				

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	428	% 	17% •
2	В	214	15%	18% • •
3	С	235	9%	25% • •
4	D	2	50%	50%
4	Е	2	100%	
4	F	2	50%	50%



Mol	Chain	Length		Quality of chain
5	G	6	17%	83%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	F	2	-	-	-	Х



# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 6891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glycoprotein G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	428	Total 3368	C 2145	N 566	O 638	S 19	0	0	0

• Molecule 2 is a protein called m102.3 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	206	Total 1585	C 991	N 273	O 316	${f S}5$	0	0	0

• Molecule 3 is a protein called IgG Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	С	226	Total 1704	C 1080	N 284	O 333	S 7	0	0	0

• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	D	2	Total         C         N         O           28         16         2         10	0	0	0
4	Е	2	Total         C         N         O           28         16         2         10	0	0	0
4	F	2	Total         C         N         O           28         16         2         10	0	0	0

• Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha



-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	G	6	Total 70	C 40	N 2	O 28	0	0	0

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
6	А	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	60	Total         O           60         60	0	0
7	В	2	Total O 2 2	0	0
7	С	4	Total O 4 4	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Glycoprotein G



#### D227 K228 K229 V230 P232 P232 SER CYS

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:	50%	50%

#### NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

100%

Chain E:

### NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

Chain F:	50%	50%

#### NAG1 NAG2

 $\label{eq:mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose]$ 

Chain G:	17%	83%
NAG1 NAG2 BMA3 MAN4 FUC5 FUC5		



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants	72.92Å 72.92Å 653.81Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Bosolution (Å)	45.42 - 2.70	Depositor
	45.42 - 2.70	EDS
% Data completeness	$100.0 \ (45.42 - 2.70)$	Depositor
(in resolution range)	99.1 (45.42 - 2.70)	EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.14 (at $2.69$ Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
B B.	0.229 , $0.275$	Depositor
$n, n_{free}$	0.232 , $0.274$	DCC
$R_{free}$ test set	1530 reflections $(5.07\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	47.3	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , $43.4$	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6891	wwPDB-VP
Average B, all atoms $(Å^2)$	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



 $<sup>^1 {\</sup>rm Intensities}$  estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
	Cham	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.29	0/3449	0.56	1/4698~(0.0%)
2	В	0.28	0/1620	0.55	0/2197
3	С	0.30	0/1747	0.54	0/2381
All	All	0.29	0/6816	0.55	1/9276~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1
3	С	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	212	ARG	NE-CZ-NH1	-7.59	116.51	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	287	HIS	Peptide
3	С	29	PHE	Peptide



## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3368	0	3306	47	0
2	В	1585	0	1543	31	0
3	С	1704	0	1657	38	0
4	D	28	0	25	1	0
4	Е	28	0	25	0	0
4	F	28	0	25	1	0
5	G	70	0	61	0	0
6	А	14	0	13	0	0
7	А	60	0	0	2	0
7	В	2	0	0	0	0
7	C	4	0	0	0	0
All	All	6891	0	6655	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:470:ASP:HB2	1:A:478:GLN:OE1	1.21	1.24
3:C:55:LEU:HB3	3:C:57:ILE:HD13	1.52	0.91
1:A:470:ASP:CB	1:A:478:GLN:OE1	2.17	0.83
1:A:404:LYS:HE2	1:A:404:LYS:HA	1.65	0.78
2:B:108:ARG:NH1	2:B:170:ASP:O	2.17	0.77
2:B:118:PHE:HB2	2:B:133:VAL:HG22	1.66	0.76
3:C:89:GLU:N	3:C:89:GLU:OE1	2.21	0.73
3:C:182:VAL:HG22	3:C:201:VAL:HG12	1.69	0.72
2:B:143:GLU:OE1	2:B:143:GLU:N	2.22	0.71
3:C:18:VAL:HG23	3:C:86:LEU:HD11	1.79	0.64
3:C:91:THR:HG23	3:C:129:THR:HA	1.80	0.64
3:C:19:LYS:HG3	3:C:82:GLU:HB2	1.79	0.63
1:A:561:THR:HA	1:A:577:LEU:O	2.01	0.61
3:C:166:PRO:O	3:C:219:HIS:NE2	2.33	0.61
2:B:145:LYS:HZ3	2:B:195:GLU:HG2	1.65	0.61
3:C:2:VAL:HG13	3:C:26:GLY:H	1.65	0.60



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:310:TRP:O	1:A:347:LYS:NZ	2.32	0.60
1:A:294:LEU:HD11	1:A:426:LEU:HD11	1.84	0.60
2:B:179:LEU:HD22	2:B:181:LEU:HD23	1.83	0.60
3:C:157:LEU:HG	3:C:201:VAL:HG23	1.84	0.60
2:B:21:LEU:HD22	2:B:102:THR:HG21	1.83	0.59
3:C:229:LYS:HD3	3:C:229:LYS:H	1.66	0.59
2:B:49:ILE:HD11	2:B:74:LEU:HD13	1.86	0.58
2:B:133:VAL:HG21	3:C:143:LEU:HD13	1.85	0.58
1:A:484:VAL:HG23	1:A:485:ILE:HG12	1.86	0.57
1:A:327:SER:HB2	1:A:330:TYR:HB3	1.86	0.57
1:A:311:THR:O	7:A:802:HOH:O	2.17	0.57
2:B:145:LYS:NZ	2:B:195:GLU:HG2	2.20	0.57
1:A:313:SER:OG	7:A:801:HOH:O	2.12	0.56
1:A:587:VAL:HG12	1:A:589:ARG:HB3	1.88	0.55
2:B:190:LYS:HA	2:B:211:ARG:HH22	1.71	0.55
3:C:16:SER:OG	3:C:17:SER:N	2.40	0.55
2:B:149:LYS:HG3	2:B:154:LEU:HD23	1.89	0.54
2:B:49:ILE:HG23	2:B:54:SER:H	1.73	0.54
3:C:138:PRO:HB3	3:C:164:TYR:HB3	1.90	0.54
3:C:51:ILE:HD13	3:C:72:THR:HG23	1.90	0.54
2:B:150:VAL:HG12	2:B:151:ASP:H	1.72	0.54
2:B:50:TYR:O	2:B:54:SER:OG	2.25	0.53
1:A:258:ARG:NH2	1:A:260:ASP:OD1	2.42	0.53
2:B:36:TRP:CD1	2:B:49:ILE:HD12	2.44	0.53
1:A:580:ILE:HD11	1:A:591:LYS:HG3	1.91	0.53
3:C:6:GLN:H	3:C:124:GLN:HE22	1.56	0.52
2:B:133:VAL:HG12	2:B:178:THR:HG23	1.90	0.52
1:A:413:LEU:HD22	1:A:475:LEU:HD22	1.92	0.52
2:B:92:TYR:CD1	3:C:117:TYR:HA	2.45	0.52
1:A:218:THR:OG1	1:A:219:ASP:N	2.44	0.51
2:B:34:LEU:HD22	2:B:72:PHE:CD1	2.46	0.50
3:C:214:ILE:HD12	3:C:228:LYS:O	2.12	0.50
1:A:329:ASP:OD2	1:A:333:LYS:NZ	2.36	0.50
1:A:264:SER:HB3	1:A:266:PHE:CE1	2.47	0.49
4:F:2:NAG:H3	4:F:2:NAG:H83	1.93	0.49
1:A:450:GLN:HB2	1:A:516:ARG:NH2	2.27	0.49
3:C:103:GLU:O	3:C:104:GLN:NE2	2.43	0.49
3:C:157:LEU:HG	3:C:201:VAL:CG2	2.42	0.49
2:B:145:LYS:HG3	2:B:197:THR:HB	1.94	0.49
1:A:324:LYS:HE3	1:A:327:SER:HB3	1.95	0.48
1:A:541:LYS:HD3	$1:\overline{A:546:LEU:HD21}$	1.94	0.48



	A h a	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
3:C:30:SER:OG	3:C:31:ASN:N	2.46	0.48
1:A:372:ARG:NH1	1:A:405:SER:OG	2.47	0.48
2:B:136:LEU:HB2	2:B:175:LEU:HB3	1.96	0.48
2:B:163:VAL:HG22	2:B:175:LEU:HD12	1.95	0.47
1:A:453:PHE:O	1:A:466:LEU:HA	2.14	0.47
3:C:13:LYS:NZ	3:C:14:ARG:O	2.36	0.46
3:C:138:PRO:HD2	3:C:224:THR:HG21	1.97	0.46
3:C:166:PRO:HG2	3:C:221:PRO:HB2	1.97	0.46
3:C:54:ILE:HG23	3:C:55:LEU:HD13	1.97	0.46
2:B:49:ILE:HA	2:B:54:SER:O	2.16	0.46
3:C:56:GLY:C	3:C:57:ILE:HD12	2.36	0.45
1:A:555:ASP:N	1:A:555:ASP:OD1	2.32	0.45
3:C:219:HIS:HB3	3:C:224:THR:OG1	2.17	0.45
1:A:573:TRP:CH2	1:A:596:LYS:HB2	2.52	0.44
1:A:191:GLN:HB2	1:A:601:CYS:SG	2.57	0.44
3:C:18:VAL:O	3:C:82:GLU:HA	2.18	0.44
4:D:1:NAG:H61	4:D:2:NAG:C7	2.47	0.44
1:A:468:ASP:CG	1:A:519:TRP:HZ2	2.21	0.44
1:A:554:ASP:N	1:A:554:ASP:OD1	2.51	0.44
3:C:67:ARG:NH2	3:C:90:ASP:OD2	2.49	0.44
2:B:38:GLN:HB2	2:B:87:TYR:CE2	2.52	0.44
3:C:173:TRP:CZ3	3:C:215:CYS:HB3	2.53	0.44
2:B:61:ASP:N	2:B:61:ASP:OD2	2.51	0.43
1:A:555:ASP:HB3	2:B:30:ARG:NH1	2.34	0.43
1:A:420:LEU:HA	1:A:420:LEU:HD23	1.83	0.43
2:B:74:LEU:HD12	2:B:75:THR:H	1.83	0.43
1:A:207:LEU:HD11	1:A:265:MET:HE2	1.99	0.43
2:B:136:LEU:O	2:B:174:SER:HA	2.19	0.43
1:A:355:GLY:HA2	1:A:444:ILE:HG23	2.01	0.43
3:C:68:VAL:HG22	3:C:83:LEU:HD13	2.00	0.43
2:B:176:SER:HB3	3:C:185:PHE:CE1	2.54	0.42
1:A:210:ASN:CB	1:A:212:ARG:HH12	2.33	0.42
1:A:210:ASN:HB2	1:A:212:ARG:HH12	1.85	0.42
3:C:135:THR:HG23	3:C:166:PRO:HG3	2.00	0.42
3:C:69:THR:HG22	3:C:71:THR:HG23	2.01	0.42
1:A:283:SER:O	1:A:293:THR:HA	2.20	0.42
1:A:284:SER:OG	1:A:293:THR:HB	2.20	0.41
1:A:460:TRP:CD2	1:A:500:PRO:HA	2.55	0.41
1:A:468:ASP:OD1	1:A:519:TRP:HZ2	2.03	0.41
1:A:181:LEU:HD21	1:A:514:ILE:HA	2.00	0.41
1:A:398:SER:OG	1:A:502:VAL:HA	2.21	0.41



Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (A)	overlap (Å)
1:A:566:PHE:HE2	1:A:575:ILE:HG13	1.85	0.41
2:B:48:LEU:HD11	2:B:87:TYR:CE2	2.55	0.41
1:A:232:SER:HA	1:A:249:ILE:O	2.21	0.41
3:C:145:PRO:HD3	3:C:157:LEU:HB3	2.03	0.41
3:C:12:LYS:HG3	3:C:18:VAL:HG22	2.03	0.41
2:B:38:GLN:HB2	2:B:87:TYR:HE2	1.85	0.41
3:C:169:VAL:HG23	3:C:219:HIS:HB2	2.02	0.40
1:A:490:GLN:HB2	1:A:505:GLU:CD	2.42	0.40
1:A:580:ILE:CD1	1:A:591:LYS:HG3	2.51	0.40
1:A:385:ILE:HD13	1:A:385:ILE:HA	1.95	0.40
1:A:466:LEU:O	1:A:480:ARG:HG2	2.22	0.40
1:A:285:THR:HG23	1:A:364:PHE:CE1	2.56	0.40
1:A:211:THR:HG21	1:A:589:ARG:NH1	2.36	0.40
2:B:176:SER:HB3	3:C:185:PHE:CD1	2.57	0.40
3:C:36:TRP:CE2	3:C:81:MET:HB2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	426/428~(100%)	405~(95%)	21~(5%)	0	100 100
2	В	202/214~(94%)	191~(95%)	11 (5%)	0	100 100
3	С	222/235~(94%)	216 (97%)	6 (3%)	0	100 100
All	All	850/877~(97%)	812 (96%)	38 (4%)	0	100 100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	382/383~(100%)	374~(98%)	8 (2%)	53 80
2	В	177/185~(96%)	170~(96%)	7 (4%)	31 60
3	С	187/197~(95%)	176 (94%)	11 (6%)	19 43
All	All	746/765~(98%)	720 (96%)	26 (4%)	36 65

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	293	THR
1	А	342	ARG
1	А	351	TYR
1	А	413	LEU
1	А	478	GLN
1	А	487	ARG
1	А	519	TRP
1	А	554	ASP
2	В	14	SER
2	В	55	ARG
2	В	61	ASP
2	В	66	SER
2	В	145	LYS
2	В	185	ASP
2	В	190	LYS
3	С	3	GLN
3	С	16	SER
3	С	63	LYS
3	С	81	MET
3	С	88	SER
3	С	131	SER
3	С	157	LEU
3	С	180	SER
3	С	189	LEU
3	С	216	ASN
3	С	229	LYS



Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	Dec	Tink	Bo	ond leng	ths	B	ond ang	gles
	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
4	NAG	D	1	1,4	14,14,15	0.26	0	17,19,21	0.65	1 (5%)
4	NAG	D	2	4	14,14,15	0.29	0	17,19,21	0.50	0
4	NAG	E	1	1,4	14, 14, 15	0.37	0	17,19,21	0.45	0
4	NAG	E	2	4	$14,\!14,\!15$	0.26	0	17,19,21	0.52	0
4	NAG	F	1	1,4	14, 14, 15	0.39	0	17,19,21	0.61	0
4	NAG	F	2	4	14,14,15	0.39	0	17,19,21	1.30	1 (5%)
5	NAG	G	1	1,5	14,14,15	0.69	1 (7%)	17,19,21	0.58	0
5	NAG	G	2	5	14,14,15	0.28	0	17,19,21	0.42	0
5	BMA	G	3	5	11,11,12	1.11	1 (9%)	15,15,17	1.03	1 (6%)
5	MAN	G	4	5	11,11,12	1.17	2 (18%)	15,15,17	1.12	1 (6%)
5	FUC	G	5	5	10, 10, 11	0.66	0	14,14,16	1.03	1 (7%)
5	FUC	G	6	5	10, 10, 11	0.89	0	14,14,16	0.98	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	NAG	Е	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Ε	2	4	-	2/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	5/6/23/26	0/1/1/1
5	NAG	G	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	G	2	5	-	2/6/23/26	0/1/1/1
5	BMA	G	3	5	-	0/2/19/22	0/1/1/1
5	MAN	G	4	5	-	0/2/19/22	0/1/1/1
5	FUC	G	5	5	-	_	0/1/1/1
5	FUC	G	6	5	_	-	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
5	G	3	BMA	O5-C1	-2.60	1.39	1.43
5	G	1	NAG	O5-C1	-2.49	1.39	1.43
5	G	4	MAN	O5-C1	-2.09	1.40	1.43
5	G	4	MAN	C2-C3	2.02	1.55	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	F	2	NAG	C2-N2-C7	4.47	129.28	122.90
5	G	4	MAN	O2-C2-C3	-2.50	105.13	110.14
5	G	6	FUC	C1-C2-C3	2.34	112.54	109.67
4	D	1	NAG	C1-O5-C5	2.21	115.18	112.19
5	G	3	BMA	C1-C2-C3	2.20	112.37	109.67
5	G	5	FUC	C1-O5-C5	2.12	117.58	112.78

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Е	2	NAG	O5-C5-C6-O6
4	Е	2	NAG	C4-C5-C6-O6
4	F	2	NAG	C8-C7-N2-C2
4	F	2	NAG	O7-C7-N2-C2
5	G	2	NAG	C8-C7-N2-C2
5	G	2	NAG	O7-C7-N2-C2



Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	F	2	NAG	C4-C5-C6-O6
4	F	2	NAG	C3-C2-N2-C7
4	F	2	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	2	NAG	1	0
4	D	2	NAG	1	0
4	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.















# 5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Chain	Dog	Tink	Bo	ond leng	$_{\rm ths}$	B	ond ang	gles
			nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
6	NAG	А	703	1	14,14,15	0.39	0	17,19,21	0.73	1 (5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	А	703	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
6	А	703	NAG	C1-O5-C5	2.43	115.48	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	А	703	NAG	O5-C5-C6-O6
6	А	703	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	428/428~(100%)	-0.03	6 (1%) 75 77	13, 26, 58, 104	0
2	В	206/214~(96%)	0.70	32~(15%) 2 1	33, 64, 99, 111	0
3	С	226/235~(96%)	0.48	21 (9%) 8 6	23, 63, 90, 101	0
All	All	860/877~(98%)	0.28	59 (6%) 16 15	13, 45, 90, 111	0

All (59) RSRZ outliers are listed below:

Mol	Chain	hain Res		RSRZ
2	В	184	ALA	5.8
2	В	152	ASN	5.1
3	С	27	GLY	4.7
2	В	192	TYR	4.3
2	В	188	LYS	4.1
3	С	141	PHE	3.9
1	А	205	TYR	3.8
2	В	181	LEU	3.6
1	А	327	SER	3.5
2	В	154	LEU	3.4
2	В	129	THR	3.4
3	С	142	PRO	3.3
3	С	28	THR	3.3
2	В	133	VAL	3.2
2	В	191	LEU	3.2
1	А	211	THR	3.1
2	В	132	VAL	3.1
2	В	194	CYS	3.1
2	В	130	ALA	3.0
2	В	186	TYR	2.9
2	В	134	CYS	2.9
2	В	182	SER	2.9
3	С	196	SER	2.9



$6 \mathrm{CMG}$

Mol	Chain	Res	Type	RSRZ
3	C	26	GLY	2.8
3	С	76	THR	2.8
2	В	189	HIS	2.7
2	В	209	PHE	2.7
3	С	1	GLU	2.7
3	С	217	VAL	2.6
2	В	118	PHE	2.6
2	В	131	SER	2.6
3	С	3	GLN	2.6
3	С	2	VAL	2.6
3	С	29	PHE	2.5
2	В	179	LEU	2.5
2	В	190	LYS	2.5
2	В	178	THR	2.5
3	С	231	GLU	2.4
2	В	153	ALA	2.4
2	В	76	ILE	2.4
2	В	187	GLU	2.4
2	В	58	GLY	2.3
1	А	206	THR	2.3
2	В	211	ARG	2.3
2	В	156	SER	2.3
3	С	32	TYR	2.3
1	А	380	SER	2.2
3	С	226	VAL	2.2
2	В	180	THR	2.2
3	С	165	PHE	2.2
3	С	14	ARG	2.1
2	В	177	SER	2.1
3	С	191	SER	2.1
3	С	200	VAL	2.1
3	С	25	SER	2.1
2	В	78	ARG	2.0
3	С	138	PRO	2.0
1	А	602	SER	2.0
2	В	128	GLY	2.0

Continued from previous page...

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B}$ -factors(Å <sup>2</sup> )	Q<0.9
5	MAN	G	4	11/12	0.72	0.21	$70,\!78,\!85,\!87$	0
4	NAG	F	2	14/15	0.75	0.52	$79,\!93,\!97,\!97$	0
4	NAG	Е	2	14/15	0.77	0.27	$57,\!63,\!79,\!84$	0
5	BMA	G	3	11/12	0.80	0.16	$61,\!68,\!76,\!80$	0
4	NAG	D	2	14/15	0.87	0.19	49,65,81,82	0
5	NAG	G	2	14/15	0.88	0.12	$46,\!54,\!60,\!64$	0
4	NAG	F	1	14/15	0.88	0.26	$30,\!55,\!75,\!83$	0
5	FUC	G	6	10/11	0.89	0.13	$39,\!43,\!47,\!48$	0
4	NAG	Е	1	14/15	0.90	0.14	$18,\!40,\!47,\!58$	0
5	NAG	G	1	14/15	0.91	0.12	$32,\!38,\!49,\!49$	0
4	NAG	D	1	14/15	0.92	0.14	$35,\!40,\!47,\!53$	0
5	FUC	G	5	10/11	0.93	0.15	$50,\!53,\!58,\!66$	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

















# 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{\AA}^2)$	Q<0.9
6	NAG	А	703	14/15	0.67	0.39	$83,\!94,\!99,\!99$	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

