



# Full wwPDB NMR Structure Validation Report ⓘ

May 7, 2024 – 12:52 pm BST

PDB ID : 2CNJ  
Title : NMR studies on the interaction of Insulin-Growth Factor II (IGF-II) with IGF2R domain 11  
Authors : Williams, C.; Prince, S.; Zaccheo, O.; Hassan, A.B.; Crosby, J.; Crump, M.  
Deposited on : 2006-05-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	D	151	

## 2 Ensemble composition and analysis i

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	D:1515-D:1520, D:1526-D:1542, D:1546-D:1597, D:1605-D:1615, D:1622-D:1647 (112)	0.47	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 7, 11, 18
2	4, 5, 17
3	14, 15, 16
4	8, 20
5	2, 6
6	3, 10
Single-model clusters	9; 12; 13; 19

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2151 atoms, of which 1064 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	D	144	2151	677	1064	189	211	10	1


## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR

Chain D: 




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR


Chain D: 



#### 4.2.2 Score per residue for model 2

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR


Chain D: 



### 4.2.3 Score per residue for model 3

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR


Chain D: 49% 23% 21% 5%



### 4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR


Chain D: 48% 25% 21% 5%



### 4.2.5 Score per residue for model 5


- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR

Chain D: 50% 22% 21% 5%




### 4.2.6 Score per residue for model 6

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR




### 4.2.7 Score per residue for model 7

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR




### 4.2.8 Score per residue for model 8


- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR



### 4.2.9 Score per residue for model 9

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR






#### 4.2.10 Score per residue for model 10

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR


Chain D: 50% 23% 21% 5%



#### 4.2.11 Score per residue for model 11

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR

Chain D: 49% 23% 21% 5%



#### 4.2.12 Score per residue for model 12

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR

Chain D: 49% 23% 21% 5%




#### 4.2.13 Score per residue for model 13

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR

Chain D: 46% 25% 21% 5%








#### 4.2.14 Score per residue for model 14

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR


Chain D: 52% 20% 21% 5%



#### 4.2.15 Score per residue for model 15

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR


Chain D: 50% 23% 21% 5%



#### 4.2.16 Score per residue for model 16


- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR

Chain D: 52% 22% 21% 5%




#### 4.2.17 Score per residue for model 17

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR




#### 4.2.18 Score per residue for model 18

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR




#### 4.2.19 Score per residue for model 19

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR



#### 4.2.20 Score per residue for model 20

- Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR



GI595	C1598	P1599	S1600	K1601	S1602	G1603	L1604	K1607	I1610	S1611	F1612	V1613	C1614	R1615	P1616	E1617	A1618	G1619	P1620	T1621	M1622	R1623	C1634	T1635	L1644	A1645	C1646	E1647	Q1648	A1649	T1650	K1651	E1652	H1653	HIS	HIS	HIS	HIS	HIS
-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-----	-----	-----	-----	-----

## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *WATER REFINED IN CNS*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *LOWEST ENERGY, LEAST VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
Sparky	structure solution	
NMRDraw	structure solution	
NMRPipe	structure solution	
CNS	structure solution	

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	D	1.14±0.03	0±0/882 ( 0.0± 0.1%)	0.82±0.02	0±0/1197 ( 0.0± 0.0%)
All	All	1.14	3/17640 ( 0.0%)	0.82	0/23940 ( 0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	D	0.0±0.0	0.1±0.2
All	All	0	1

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	D	1634	CYS	C-N	-6.20	1.19	1.34	14	1
1	D	1551	SER	CA-CB	-5.75	1.44	1.52	11	1
1	D	1550	MET	C-N	-5.08	1.22	1.34	11	1

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	D	1623	ARG	Sidechain	1

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	D	863	856	855	18±2
All	All	17260	17120	17100	351

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:D:1643:PRO:HA	1:D:1646:CYS:SG	0.81	2.16	19	11
1:D:1530:SER:HA	1:D:1533:SER:OG	0.65	1.90	4	17
1:D:1516:CYS:SG	1:D:1579:LYS:HB2	0.65	2.32	3	8
1:D:1589:GLN:HA	1:D:1610:ILE:O	0.62	1.95	12	20
1:D:1586:GLN:HB3	1:D:1614:CYS:SG	0.62	2.34	14	5
1:D:1559:CYS:SG	1:D:1566:CYS:HB2	0.59	2.38	15	14
1:D:1520:ASN:HB2	1:D:1583:TYR:CE2	0.58	2.33	19	1
1:D:1552:ILE:HA	1:D:1592:TYR:OH	0.58	1.97	8	16
1:D:1606:TYR:HA	1:D:1634:CYS:HB2	0.57	1.75	14	3
1:D:1516:CYS:HB3	1:D:1553:CYS:SG	0.57	2.40	19	3
1:D:1584:VAL:O	1:D:1587:VAL:HB	0.56	2.00	12	15
1:D:1535:ARG:HA	1:D:1555:GLU:O	0.55	2.01	1	11
1:D:1517:GLN:HA	1:D:1527:PHE:O	0.55	2.01	6	11
1:D:1614:CYS:HB3	1:D:1645:ALA:O	0.54	2.03	6	18
1:D:1518:VAL:HG21	1:D:1583:TYR:N	0.53	2.19	4	7
1:D:1552:ILE:HA	1:D:1592:TYR:HH	0.52	1.65	16	6
1:D:1520:ASN:HB2	1:D:1583:TYR:HE2	0.52	1.63	19	1
1:D:1556:ASN:O	1:D:1559:CYS:HB2	0.52	2.04	15	13
1:D:1546:GLY:HA2	1:D:1569:GLN:NE2	0.52	2.19	3	4
1:D:1553:CYS:HA	1:D:1578:ASN:O	0.51	2.05	14	6
1:D:1517:GLN:HB3	1:D:1526:LEU:HD21	0.51	1.81	14	1
1:D:1518:VAL:HG21	1:D:1582:ARG:HA	0.50	1.84	1	4
1:D:1607:LYS:HE2	1:D:1635:THR:OG1	0.50	2.07	5	1
1:D:1558:ASN:OD1	1:D:1571:ARG:HD2	0.50	2.07	11	1
1:D:1607:LYS:HD3	1:D:1635:THR:OG1	0.50	2.07	2	4
1:D:1529:LEU:HD13	1:D:1612:PHE:CZ	0.49	2.42	17	9
1:D:1535:ARG:HG2	1:D:1557:GLU:N	0.49	2.21	20	6

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:D:1611:SER:O	1:D:1639:SER:HA	0.49	2.06	1	16
1:D:1581:LEU:HD13	1:D:1590:LEU:HD23	0.49	1.84	13	1
1:D:1581:LEU:HD11	1:D:1588:LEU:HB3	0.49	1.85	20	6
1:D:1537:GLY:HA3	1:D:1551:SER:HA	0.49	1.83	9	5
1:D:1591:VAL:HA	1:D:1608:SER:O	0.49	2.08	19	6
1:D:1556:ASN:HB3	1:D:1559:CYS:SG	0.49	2.48	14	7
1:D:1539:THR:O	1:D:1623:ARG:HD2	0.48	2.08	19	3
1:D:1582:ARG:O	1:D:1588:LEU:HA	0.48	2.07	17	2
1:D:1515:ASP:O	1:D:1517:GLN:HG2	0.48	2.08	12	1
1:D:1542:TYR:OH	1:D:1629:LEU:HD22	0.48	2.08	18	1
1:D:1533:SER:HA	1:D:1552:ILE:O	0.48	2.08	7	1
1:D:1567:PHE:O	1:D:1571:ARG:HD3	0.48	2.08	18	1
1:D:1546:GLY:HA3	1:D:1569:GLN:NE2	0.48	2.24	17	1
1:D:1607:LYS:HE3	1:D:1635:THR:OG1	0.48	2.08	4	1
1:D:1516:CYS:SG	1:D:1530:SER:HB3	0.47	2.49	16	1
1:D:1640:TRP:CD1	1:D:1642:THR:HB	0.47	2.45	10	4
1:D:1515:ASP:CB	1:D:1579:LYS:HD3	0.47	2.39	8	1
1:D:1615:ARG:NE	1:D:1615:ARG:HA	0.47	2.24	20	1
1:D:1537:GLY:CA	1:D:1551:SER:HA	0.46	2.41	10	4
1:D:1559:CYS:HB3	1:D:1563:VAL:HB	0.46	1.87	15	19
1:D:1614:CYS:O	1:D:1615:ARG:HD2	0.46	2.09	7	1
1:D:1592:TYR:O	1:D:1607:LYS:HA	0.46	2.11	10	6
1:D:1520:ASN:HA	1:D:1583:TYR:CE2	0.46	2.45	15	1
1:D:1566:CYS:SG	1:D:1571:ARG:HA	0.45	2.52	6	2
1:D:1581:LEU:HA	1:D:1590:LEU:CD2	0.45	2.42	12	1
1:D:1553:CYS:SG	1:D:1579:LYS:HE2	0.45	2.51	16	1
1:D:1535:ARG:HG3	1:D:1555:GLU:O	0.44	2.11	9	1
1:D:1591:VAL:HG13	1:D:1608:SER:O	0.44	2.12	13	3
1:D:1647:GLU:OE1	1:D:1647:GLU:HA	0.44	2.13	9	1
1:D:1592:TYR:HB2	1:D:1608:SER:HB2	0.44	1.89	3	5
1:D:1593:LYS:HA	1:D:1607:LYS:HB2	0.44	1.90	7	1
1:D:1518:VAL:HB	1:D:1581:LEU:O	0.43	2.13	6	1
1:D:1534:GLY:O	1:D:1551:SER:HB2	0.43	2.14	8	1
1:D:1529:LEU:HG	1:D:1581:LEU:HD22	0.43	1.90	9	2
1:D:1591:VAL:HG12	1:D:1593:LYS:HG3	0.43	1.91	16	1
1:D:1553:CYS:HB2	1:D:1579:LYS:CB	0.42	2.44	11	2
1:D:1614:CYS:HB2	1:D:1646:CYS:HB3	0.42	1.91	17	1
1:D:1519:THR:HG22	1:D:1526:LEU:N	0.42	2.30	19	2
1:D:1552:ILE:HG22	1:D:1553:CYS:SG	0.42	2.54	3	3
1:D:1516:CYS:HB3	1:D:1553:CYS:HG	0.42	1.74	19	1
1:D:1546:GLY:HA3	1:D:1569:GLN:HE21	0.42	1.75	16	1

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:D:1606:TYR:HA	1:D:1634:CYS:CB	0.41	2.46	3	1
1:D:1646:CYS:O	1:D:1647:GLU:HB2	0.41	2.15	1	1
1:D:1515:ASP:HB3	1:D:1579:LYS:O	0.41	2.15	8	2
1:D:1627:ILE:HD11	1:D:1639:SER:OG	0.41	2.16	1	1
1:D:1535:ARG:HB3	1:D:1557:GLU:HB2	0.41	1.91	4	1
1:D:1593:LYS:HA	1:D:1607:LYS:CB	0.41	2.46	9	2
1:D:1587:VAL:HG11	1:D:1611:SER:HB3	0.41	1.92	12	1
1:D:1528:ASP:O	1:D:1644:LEU:HD11	0.41	2.15	8	1
1:D:1538:PHE:N	1:D:1550:MET:O	0.41	2.53	18	1
1:D:1592:TYR:CE2	1:D:1610:ILE:HD11	0.40	2.51	8	1
1:D:1580:ARG:HD3	1:D:1580:ARG:O	0.40	2.16	7	1
1:D:1576:LYS:HE3	1:D:1594:ASP:O	0.40	2.16	1	1
1:D:1549:TYR:CE2	1:D:1568:GLY:HA2	0.40	2.51	10	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	112/151 (74%)	102±1 (92±1%)	8±1 (7±1%)	2±1 (2±1%)	13	56
All	All	2240/3020 (74%)	2050 (92%)	151 (7%)	39 (2%)	13	56

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	D	1647	GLU	20
1	D	1595	GLY	8
1	D	1622	ASN	7
1	D	1634	CYS	2
1	D	1561	PRO	2



### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	97/131 (74%)	90±2 (93±2%)	7±2 (7±2%)	17	65
All	All	1940/2620 (74%)	1795 (93%)	145 (7%)	17	65

All 24 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	D	1555	GLU	20
1	D	1582	ARG	16
1	D	1607	LYS	14
1	D	1571	ARG	13
1	D	1519	THR	12
1	D	1579	LYS	11
1	D	1644	LEU	9
1	D	1570	THR	7
1	D	1583	TYR	6
1	D	1611	SER	5
1	D	1647	GLU	5
1	D	1547	LEU	5
1	D	1528	ASP	4
1	D	1633	THR	2
1	D	1557	GLU	2
1	D	1580	ARG	2
1	D	1615	ARG	2
1	D	1630	ASP	2
1	D	1590	LEU	2
1	D	1593	LYS	2
1	D	1520	ASN	1
1	D	1622	ASN	1
1	D	1631	LYS	1
1	D	1526	LEU	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	14-D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
14	D	1634:CYS	C	1635:THR	N	1.19

## 7 Chemical shift validation

No chemical shift data were provided