



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 4, 2024 – 01:40 am GMT

PDB ID : 8CN6  
Title : CD59 in complex with CP-06 peptide  
Authors : Bickel, J.; Ahmed, A.; Morgan, M.; Horrell, S.; El Omari, K.; Couves, E.C.;  
Bubeck, D.; Tate, E.  
Deposited on : 2023-02-22  
Resolution : 2.43 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2124 (2.46-2.42)
Clashscore	180529	2259 (2.46-2.42)
Ramachandran outliers	177936	2244 (2.46-2.42)
Sidechain outliers	177891	2244 (2.46-2.42)
RSRZ outliers	164620	2124 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	77	 66% 27% 6%
1	B	77	 83% 14% ..
1	C	77	 78% 17% . .
1	D	77	 82% 17% .
1	E	77	 74% 18% 5% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	77	
2	G	16	
2	H	16	
2	J	16	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DAR	J	13	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7804 atoms, of which 3741 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CD59 glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	77	1198	385	577	104	121	11	14	0	0
1	B	76	1183	380	571	103	118	11	14	0	0
1	C	76	1183	380	571	103	118	11	14	0	0
1	D	77	1198	385	577	104	121	11	14	0	0
1	E	76	1183	380	571	103	118	11	14	0	0
1	F	77	1172	379	559	102	121	11	18	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P13987
B	0	MET	-	initiating methionine	UNP P13987
C	0	MET	-	initiating methionine	UNP P13987
D	0	MET	-	initiating methionine	UNP P13987
E	0	MET	-	initiating methionine	UNP P13987
F	0	MET	-	initiating methionine	UNP P13987

- Molecule 2 is a protein called Cyclic peptide CP-06.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	J	16	225	74	105	24	21	1	5	0	1
2	G	16	225	74	105	24	21	1	5	0	1
2	H	16	225	74	105	24	21	1	5	0	1

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total 1	Zn 1	0	0
3	F	1	Total 1	Zn 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	O 1	0	0
4	B	3	Total 3	O 3	0	0
4	C	3	Total 3	O 3	0	0
4	E	1	Total 1	O 1	0	0
4	F	1	Total 1	O 1	0	0
4	H	1	Total 1	O 1	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CD59 glycoprotein

Chain A:  66% 27% 6%





- Molecule 1: CD59 glycoprotein

Chain B:  83% 14% ..





- Molecule 1: CD59 glycoprotein

Chain C:  78% 17% ..



- Molecule 1: CD59 glycoprotein

Chain D:  82% 17% .



- Molecule 1: CD59 glycoprotein

Chain E:  74% 18% 5% ..




- Molecule 1: CD59 glycoprotein


Chain F:  64% 27% 8% .




- Molecule 2: Cyclic peptide CP-06



- Molecule 2: Cyclic peptide CP-06



- Molecule 2: Cyclic peptide CP-06



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	30.94Å 85.88Å 96.57Å 90.00° 90.03° 90.00°	Depositor
Resolution (Å)	48.33 – 2.43 48.33 – 2.43	Depositor EDS
% Data completeness (in resolution range)	98.9 (48.33-2.43) 98.8 (48.33-2.43)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 2.42Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.235 , 0.298 0.234 , 0.298	Depositor DCC
$R_{free}$ test set	969 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	66.4	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 52.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.408 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7804	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DSN, DCY, DAR, ACE, NH2, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	1/633 (0.2%)	1.21	6/856 (0.7%)
1	B	0.71	1/624 (0.2%)	1.24	5/844 (0.6%)
1	C	0.62	1/624 (0.2%)	1.21	5/844 (0.6%)
1	D	0.61	1/633 (0.2%)	1.17	4/856 (0.5%)
1	E	0.77	2/624 (0.3%)	1.28	7/844 (0.8%)
1	F	0.65	1/625 (0.2%)	1.34	6/848 (0.7%)
2	G	0.67	0/79	1.93	3/106 (2.8%)
2	H	0.65	0/79	2.56	8/106 (7.5%)
2	J	1.08	0/79	1.90	1/106 (0.9%)
All	All	0.68	7/4000 (0.2%)	1.31	45/5410 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	G	0	2
2	H	0	3
2	J	0	4
All	All	0	9

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	44	HIS	CB-CG	7.88	1.64	1.50
1	B	44	HIS	CB-CG	6.98	1.62	1.50
1	E	43	GLU	CD-OE2	6.25	1.32	1.25
1	C	44	HIS	CB-CG	6.19	1.61	1.50
1	F	44	HIS	CB-CG	6.15	1.61	1.50
1	D	44	HIS	CB-CG	5.51	1.59	1.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	44	HIS	CB-CG	5.16	1.59	1.50

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	4	TRP	CA-CB-CG	10.56	133.76	113.70
2	G	8	ASN	CB-CA-C	-10.48	89.45	110.40
1	E	44	HIS	CA-CB-CG	9.81	130.28	113.60
1	B	44	HIS	CB-CA-C	9.28	128.96	110.40
1	E	44	HIS	CB-CA-C	9.13	128.67	110.40
2	H	8	ASN	CB-CA-C	-8.97	92.46	110.40
1	F	44	HIS	CB-CA-C	8.77	127.94	110.40
1	B	44	HIS	CA-CB-CG	8.60	128.22	113.60
1	F	44	HIS	CA-CB-CG	8.49	128.04	113.60
1	C	44	HIS	CB-CA-C	8.47	127.35	110.40
1	D	44	HIS	CB-CA-C	8.15	126.71	110.40
2	H	4	TRP	N-CA-CB	-8.02	96.16	110.60
2	H	2	TYR	CB-CA-C	8.01	126.43	110.40
1	A	44	HIS	CB-CA-C	8.01	126.42	110.40
1	C	44	HIS	CA-CB-CG	7.71	126.70	113.60
1	A	44	HIS	CA-CB-CG	7.54	126.41	113.60
1	D	44	HIS	CA-CB-CG	7.43	126.24	113.60
2	G	8	ASN	N-CA-C	7.08	130.11	111.00
1	F	45	CYS	CB-CA-C	-6.62	97.17	110.40
1	F	64	CYS	CA-CB-SG	-6.59	102.14	114.00
1	E	45	CYS	CB-CA-C	-6.50	97.41	110.40
1	A	45	CYS	CB-CA-C	-6.37	97.67	110.40
1	C	45	CYS	CB-CA-C	-6.32	97.75	110.40
1	A	1	LEU	CB-CG-CD2	6.32	121.74	111.00
1	D	0	MET	CG-SD-CE	-6.14	90.38	100.20
2	J	4	TRP	CA-CB-CG	6.10	125.30	113.70
1	F	58	GLU	CB-CG-CD	5.90	130.12	114.20
1	E	44	HIS	ND1-CG-CD2	-5.87	97.78	106.00
1	D	45	CYS	CB-CA-C	-5.82	98.76	110.40
2	H	4	TRP	N-CA-C	-5.74	95.52	111.00
2	H	8	ASN	CA-CB-CG	-5.74	100.78	113.40
2	H	6	TRP	N-CA-C	-5.48	96.22	111.00
1	C	0	MET	CG-SD-CE	-5.34	91.66	100.20
1	A	63	CYS	CB-CA-C	-5.31	99.78	110.40
1	B	0	MET	CG-SD-CE	5.23	108.57	100.20
1	A	49	ASP	CB-CA-C	5.20	120.80	110.40
1	E	29	THR	OG1-CB-CG2	-5.13	98.20	110.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	ACE	C-N-CA	5.13	134.52	121.70
1	E	44	HIS	CB-CG-ND1	5.11	135.97	123.20
1	E	49	ASP	CB-CA-C	5.09	120.58	110.40
1	F	69	CYS	CA-CB-SG	-5.07	104.88	114.00
1	B	1	LEU	N-CA-CB	-5.05	100.30	110.40
2	H	4	TRP	CB-CG-CD2	5.03	133.13	126.60
1	C	49	ASP	CB-CA-C	5.02	120.45	110.40
1	B	45	CYS	CB-CA-C	-5.01	100.38	110.40

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	G	2	TYR	Peptide
2	G	8	ASN	Peptide
2	H	2	TYR	Peptide
2	H	8	ASN	Peptide
2	H	9	DAR	Peptide
2	J	10	DSN	Peptide
2	J	11	DSN	Peptide
2	J	4	TRP	Peptide
2	J	8	ASN	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	621	577	576	21	0
1	B	612	571	570	5	0
1	C	612	571	570	9	0
1	D	621	577	576	9	0
1	E	612	571	570	12	0
1	F	613	559	554	18	0
2	G	120	105	99	9	0
2	H	120	105	101	10	0
2	J	120	105	101	15	0
3	C	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	1	0	0	0	0
4	A	1	0	0	0	0
4	B	3	0	0	1	0
4	C	3	0	0	1	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	H	1	0	0	0	0
All	All	4063	3741	3717	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:66:LYS:O	1:E:67:ASP:HB2	1.62	0.98
1:F:66:LYS:O	1:F:67:ASP:HB2	1.67	0.94
2:J:9:DAR:C	2:J:10:DSN:OG	2.14	0.92
1:F:0:MET:O	1:F:0:MET:SD	2.36	0.83
1:A:27:LEU:HD11	1:A:29:THR:CG2	2.09	0.83
2:J:6:TRP:O	2:J:12:VAL:HG12	1.78	0.82
1:E:27:LEU:HD11	1:E:29:THR:CG2	2.11	0.81
2:J:4:TRP:HB3	2:J:13:DAR:HH12	1.54	0.71
1:F:60:THR:HG22	2:H:8:ASN:OD1	1.92	0.70
1:D:62:TYR:HD1	2:H:13:DAR:HB2	1.60	0.66
1:A:22:ASP:OD2	1:A:23:PHE:CD2	2.49	0.66
2:J:6:TRP:O	2:J:12:VAL:CG1	2.44	0.64
1:F:64:CYS:SG	1:F:65:LYS:N	2.70	0.64
1:A:61:TYR:CZ	2:G:14:GLY:HA3	2.34	0.62
2:G:8:ASN:HB3	2:G:9:DAR:HA	1.81	0.62
1:D:55:ARG:HG3	1:D:55:ARG:HH11	1.65	0.62
1:A:22:ASP:OD2	1:A:23:PHE:CE2	2.52	0.62
1:A:55:ARG:HH11	1:A:55:ARG:HG3	1.64	0.61
2:G:8:ASN:O	2:G:9:DAR:HB2	2.01	0.61
1:A:30:LYS:HG2	1:A:35:VAL:HG22	1.83	0.60
1:A:1:LEU:HD11	1:A:26:CYS:SG	2.41	0.60
2:H:3:SER:HA	2:H:15:DCY:SG	2.41	0.60
2:G:8:ASN:O	2:G:9:DAR:CB	2.49	0.59
2:J:4:TRP:HB3	2:J:13:DAR:NH1	2.15	0.59
1:A:1:LEU:CD1	1:A:3:CYS:SG	2.91	0.58
1:C:43:GLU:OE1	1:C:44:HIS:NE2	2.38	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:60:THR:HG22	2:G:8:ASN:OD1	2.04	0.56
2:G:8:ASN:HB2	2:G:10:DSN:HA	1.86	0.56
1:D:43:GLU:OE2	1:D:44:HIS:NE2	2.39	0.56
1:B:43:GLU:OE2	1:B:44:HIS:NE2	2.39	0.55
1:A:0:MET:N	1:F:34:GLN:OE1	2.28	0.55
2:H:4:TRP:CH2	2:H:13:DAR:NH1	2.74	0.55
1:A:1:LEU:HD13	1:A:3:CYS:SG	2.47	0.55
1:A:43:GLU:OE2	1:A:44:HIS:NE2	2.40	0.55
1:A:22:ASP:OD1	1:A:41:LYS:HE3	2.07	0.55
1:F:43:GLU:OE2	1:F:44:HIS:NE2	2.40	0.55
1:E:28:ILE:CD1	1:E:72:ASN:HA	2.38	0.54
1:E:43:GLU:OE2	1:E:44:HIS:NE2	2.41	0.53
1:E:66:LYS:O	1:E:67:ASP:CB	2.44	0.53
1:E:58:GLU:HG2	2:J:9:DAR:NH2	2.24	0.53
2:H:1:ACE:H3	2:H:15:DCY:C	2.39	0.52
1:B:28:ILE:CD1	1:B:72:ASN:HA	2.40	0.52
1:E:27:LEU:HD11	1:E:29:THR:HG22	1.91	0.52
1:E:64:CYS:CB	2:J:3:SER:HB3	2.40	0.52
2:J:4:TRP:CB	2:J:13:DAR:HH12	2.21	0.52
1:F:31:ALA:O	1:F:34:GLN:NE2	2.44	0.51
2:H:8:ASN:HB2	2:H:11:DSN:HA	1.92	0.51
2:J:4:TRP:CD1	2:J:13:DAR:NH1	2.77	0.51
2:J:4:TRP:HD1	2:J:13:DAR:NH1	2.09	0.51
1:E:30:LYS:HE3	1:E:75:LEU:HD12	1.93	0.50
1:C:69:CYS:O	1:C:69:CYS:SG	2.68	0.50
1:F:28:ILE:CD1	1:F:72:ASN:HA	2.42	0.50
1:A:27:LEU:HD11	1:A:29:THR:HG22	1.90	0.49
1:C:75:LEU:C	4:C:203:HOH:O	2.51	0.49
1:F:75:LEU:O	1:F:76:GLU:C	2.51	0.48
2:G:8:ASN:HB2	2:G:11:DSN:HA	1.95	0.48
1:A:59:LEU:C	1:A:59:LEU:HD12	2.34	0.48
1:C:59:LEU:HD12	1:C:59:LEU:C	2.33	0.48
1:F:59:LEU:C	1:F:59:LEU:HD12	2.34	0.48
1:E:59:LEU:C	1:E:59:LEU:HD12	2.34	0.48
2:J:4:TRP:HD1	2:J:13:DAR:HH12	1.58	0.48
1:F:69:CYS:SG	1:F:70:ASN:N	2.87	0.48
1:C:28:ILE:CD1	1:C:72:ASN:HA	2.44	0.47
1:F:61:TYR:O	2:H:6:TRP:HA	2.14	0.47
1:A:28:ILE:CD1	1:A:72:ASN:HA	2.45	0.47
1:B:47:PHE:HB2	4:B:103:HOH:O	2.14	0.47
1:D:28:ILE:CD1	1:D:72:ASN:HA	2.45	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:29:THR:HG22	1:E:61:TYR:CB	2.46	0.47
1:A:12:ASP:O	1:A:14:LYS:HD2	2.15	0.46
1:B:59:LEU:C	1:B:59:LEU:HD12	2.36	0.46
1:D:59:LEU:C	1:D:59:LEU:HD12	2.35	0.46
1:E:28:ILE:HD11	1:E:72:ASN:HA	1.98	0.46
1:C:65:LYS:HE3	1:C:65:LYS:HB2	1.71	0.45
1:A:67:ASP:OD2	1:F:32:GLY:HA3	2.16	0.45
1:F:66:LYS:O	1:F:67:ASP:CB	2.48	0.45
1:C:59:LEU:HD12	1:C:59:LEU:O	2.17	0.44
1:D:62:TYR:CD1	2:H:13:DAR:HB2	2.47	0.44
2:J:6:TRP:O	2:J:12:VAL:HA	2.17	0.44
1:C:1:LEU:CD1	1:C:65:LYS:O	2.66	0.44
1:A:29:THR:HG22	1:A:61:TYR:CB	2.48	0.43
1:A:61:TYR:O	2:G:13:DAR:HB3	2.18	0.43
1:F:63:CYS:O	2:H:4:TRP:HA	2.17	0.43
2:G:8:ASN:CB	2:G:10:DSN:HA	2.49	0.43
2:H:4:TRP:CZ2	2:H:13:DAR:NH1	2.86	0.43
2:J:7:GLY:HA2	2:J:11:DSN:OG	2.19	0.43
1:D:51:THR:HB	1:D:56:GLU:O	2.19	0.42
1:F:59:LEU:HD12	1:F:59:LEU:O	2.19	0.42
1:A:51:THR:HB	1:A:56:GLU:O	2.20	0.42
2:J:7:GLY:CA	2:J:11:DSN:OG	2.68	0.41
1:D:59:LEU:HD12	1:D:59:LEU:O	2.20	0.41
1:A:59:LEU:HD12	1:A:59:LEU:O	2.21	0.41
1:F:28:ILE:HD11	1:F:72:ASN:HA	2.03	0.41
2:J:8:ASN:O	2:J:11:DSN:HB2	2.21	0.41
1:B:28:ILE:HD11	1:B:72:ASN:HA	2.03	0.41
1:D:43:GLU:OE2	1:D:44:HIS:CD2	2.73	0.40
1:F:51:THR:HB	1:F:56:GLU:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	75/77 (97%)	73 (97%)	2 (3%)	0	100	100
1	B	74/77 (96%)	74 (100%)	0	0	100	100
1	C	74/77 (96%)	74 (100%)	0	0	100	100
1	D	75/77 (97%)	73 (97%)	1 (1%)	1 (1%)	10	9
1	E	74/77 (96%)	73 (99%)	0	1 (1%)	9	8
1	F	75/77 (97%)	74 (99%)	0	1 (1%)	10	9
2	G	9/16 (56%)	8 (89%)	0	1 (11%)	0	0
2	H	9/16 (56%)	7 (78%)	1 (11%)	1 (11%)	0	0
2	J	9/16 (56%)	6 (67%)	1 (11%)	2 (22%)	0	0
All	All	474/510 (93%)	462 (98%)	5 (1%)	7 (2%)	8	7

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	67	ASP
1	F	67	ASP
2	J	3	SER
2	J	4	TRP
1	D	13	CYS
2	G	3	SER
2	H	4	TRP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	72/72 (100%)	69 (96%)	3 (4%)	25	35
1	B	71/72 (99%)	68 (96%)	3 (4%)	25	35
1	C	71/72 (99%)	66 (93%)	5 (7%)	12	15
1	D	72/72 (100%)	70 (97%)	2 (3%)	38	51
1	E	71/72 (99%)	66 (93%)	5 (7%)	12	15
1	F	70/72 (97%)	62 (89%)	8 (11%)	4	4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	G	7/7 (100%)	3 (43%)	4 (57%)	0	0
2	H	7/7 (100%)	4 (57%)	3 (43%)	0	0
2	J	7/7 (100%)	5 (71%)	2 (29%)	0	0
All	All	448/453 (99%)	413 (92%)	35 (8%)	10	12

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	ASP
1	A	14	LYS
1	A	22	ASP
1	B	29	THR
1	B	33	LEU
1	B	34	GLN
1	C	22	ASP
1	C	29	THR
1	C	34	GLN
1	C	65	LYS
1	C	69	CYS
1	D	22	ASP
1	D	29	THR
1	E	0	MET
1	E	28	ILE
1	E	33	LEU
1	E	34	GLN
1	E	67	ASP
1	F	14	LYS
1	F	22	ASP
1	F	28	ILE
1	F	29	THR
1	F	34	GLN
1	F	57	ASN
1	F	64	CYS
1	F	76	GLU
2	J	8	ASN
2	J	12	VAL
2	G	2	TYR
2	G	3	SER
2	G	5	THR
2	G	12	VAL
2	H	3	SER

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
2	H	4	TRP
2	H	5	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	34	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

15 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DCY	H	15	2	4,5,6	0.95	0	1,5,7	2.36	1 (100%)
2	DCY	J	15	2	4,5,6	0.60	0	1,5,7	1.49	0
2	DCY	G	15	2	4,5,6	0.68	0	1,5,7	1.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DCY	H	15	2	-	1/1/4/6	-
2	DCY	J	15	2	-	0/1/4/6	-
2	DCY	G	15	2	-	1/1/4/6	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	15	DCY	CA-CB-SG	-2.36	109.35	114.44

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	G	15	DCY	N-CA-CB-SG
2	H	15	DCY	N-CA-CB-SG

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	15	DCY	2	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	77/77 (100%)	-0.65	0 100 100	56, 74, 104, 121	0
1	B	76/77 (98%)	-0.62	0 100 100	46, 71, 98, 114	0
1	C	76/77 (98%)	-0.49	0 100 100	47, 84, 123, 130	0
1	D	77/77 (100%)	-0.58	0 100 100	51, 72, 102, 105	0
1	E	76/77 (98%)	-0.57	0 100 100	48, 72, 99, 121	0
1	F	77/77 (100%)	-0.50	0 100 100	47, 83, 121, 135	0
2	G	9/16 (56%)	-0.92	0 100 100	53, 64, 86, 109	0
2	H	9/16 (56%)	-0.79	0 100 100	54, 63, 93, 127	0
2	J	9/16 (56%)	-0.52	0 100 100	51, 65, 111, 125	0
All	All	486/510 (95%)	-0.58	0 100 100	46, 75, 112, 135	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DSN	J	11	6/7	0.94	0.08	30,67,81,132	1
2	DAR	H	9	11/12	0.96	0.08	76,85,92,95	0
2	DAR	H	13	11/12	0.96	0.08	51,82,113,116	0
2	DCY	G	15	6/7	0.96	0.06	83,94,104,111	0
2	DCY	H	15	6/7	0.96	0.08	77,91,100,105	0
2	DAR	G	13	11/12	0.97	0.10	61,88,144,147	0
2	DSN	H	10	6/7	0.97	0.06	30,74,82,85	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	DSN	G	11	6/7	0.98	0.06	30,62,69,69	1
2	DSN	H	11	6/7	0.98	0.06	30,56,62,64	1
2	DAR	J	13	11/12	0.98	0.06	56,69,79,82	0
2	DSN	J	10	6/7	0.98	0.06	30,86,102,104	1
2	DSN	G	10	6/7	0.98	0.06	30,67,76,77	1
2	DAR	G	9	11/12	0.98	0.06	59,66,96,101	0
2	DAR	J	9	11/12	0.98	0.07	87,98,114,118	0
2	DCY	J	15	6/7	0.99	0.04	70,75,85,89	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	C	101	1/1	0.99	0.03	93,93,93,93	0
3	ZN	F	101	1/1	1.00	0.02	90,90,90,90	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.