



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2023 – 10:23 am BST

PDB ID : 8CNH
Title : Crystal structure of human soluble adenylyl cyclase (sAC) in complex with inhibitor TDI-10512
Authors : Steegborn, C.
Deposited on : 2023-02-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.32.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	513	-	-	-	X

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7466 atoms, of which 3657 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Adenylate cyclase type 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	453	7153	2333	3554	586	649	31	0	4	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	470	HIS	-	expression tag	UNP Q96PN6
A	471	HIS	-	expression tag	UNP Q96PN6
A	472	HIS	-	expression tag	UNP Q96PN6
A	473	HIS	-	expression tag	UNP Q96PN6
A	474	HIS	-	expression tag	UNP Q96PN6
A	475	HIS	-	expression tag	UNP Q96PN6

- Molecule 2 is methyl 2-[[3-(2-azanyl-6-chloranyl-pyrimidin-4-yl)-1-methyl-pyrazol-4-yl]methyl]benzoate (three-letter code: V6U) (formula: C₁₇H₁₆ClN₅O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Cl	H	N			O
2	A	1	41	17	1	16	5	2	0	0

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	S		
3	A	1	10	2	6	1	1	0	0
3	A	1	10	2	6	1	1	0	0
3	A	1	10	2	6	1	1	0	0
3	A	1	10	2	6	1	1	0	0
3	A	1	10	2	6	1	1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	A	1	7	2	3	2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	125	125	125	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenylate cyclase type 10



4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	99.14Å 99.14Å 99.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.38 – 2.00 44.38 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.5 (44.38-2.00) 95.5 (44.38-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.189 , 0.237 0.189 , 0.237	Depositor DCC
R_{free} test set	1798 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	37.8	Xtrriage
Anisotropy	0.072	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.055 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7466	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, V6U, EDO, DMS, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	4/3688 (0.1%)	0.84	6/4993 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	GLU	CB-CG	-7.17	1.38	1.52
1	A	456	SER	CA-CB	-6.13	1.43	1.52
1	A	373	CYS	CB-SG	-5.72	1.72	1.81
1	A	240	TYR	CE1-CZ	-5.18	1.31	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	22	ASP	CB-CG-OD2	-7.44	111.60	118.30
1	A	278	ASP	CB-CG-OD1	-6.18	112.73	118.30
1	A	418	MET	CG-SD-CE	-5.77	90.97	100.20
1	A	84	GLU	CA-CB-CG	5.60	125.72	113.40
1	A	22	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	367	MET	CG-SD-CE	5.32	108.71	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3599	3554	3538	41	0
2	A	25	16	0	0	0
3	A	20	30	30	1	0
4	A	36	54	54	2	0
5	A	4	3	3	1	0
6	A	125	0	0	5	0
All	All	3809	3657	3625	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:PHE:HE2	1:A:79:ILE:HD11	1.58	0.68
1:A:207:VAL:HG23	1:A:210:GLN:HB2	1.76	0.67
1:A:324:VAL:HG12	1:A:365:CYS:SG	2.35	0.66
1:A:416:ARG:NE	6:A:601:HOH:O	2.29	0.61
1:A:51:PHE:CE2	1:A:79:ILE:HD11	2.34	0.61
1:A:52:THR:HG22	6:A:718:HOH:O	2.03	0.59
1:A:84:GLU:O	1:A:88:ILE:HG23	2.06	0.55
1:A:350:PHE:HB3	1:A:351:PRO:HD2	1.89	0.55
1:A:90:GLY:O	4:A:516:EDO:H12	2.08	0.54
1:A:347:VAL:HB	1:A:350:PHE:CE2	2.44	0.53
1:A:309:GLU:OE2	1:A:378:LYS:HE2	2.09	0.53
1:A:355:VAL:HG12	1:A:356:PRO:HD2	1.91	0.51
1:A:57:LYS:O	1:A:60:SER:HB3	2.13	0.49
1:A:355:VAL:HG12	1:A:356:PRO:CD	2.43	0.48
1:A:23:LEU:HA	5:A:513:ACT:H2	1.93	0.48
1:A:107:VAL:HG11	1:A:115:ILE:HG13	1.96	0.47
1:A:350:PHE:HB3	1:A:351:PRO:CD	2.45	0.47
1:A:331:GLN:HB2	1:A:350:PHE:CE2	2.50	0.46
1:A:297:VAL:O	1:A:343:SER:HA	2.16	0.46
1:A:29:PHE:CD2	1:A:29:PHE:N	2.81	0.45
1:A:189:LEU:HG	1:A:215:VAL:HG23	1.98	0.45
1:A:141:ILE:HG23	1:A:141:ILE:O	2.15	0.45
1:A:207:VAL:HG22	1:A:212:ALA:O	2.17	0.44
1:A:207:VAL:HG23	1:A:210:GLN:CB	2.45	0.44
1:A:376:VAL:HB	1:A:379:ILE:HD12	1.99	0.44
1:A:328:PHE:CD2	1:A:365:CYS:HB2	2.54	0.43
1:A:416:ARG:NH1	1:A:451:LYS:O	2.52	0.43
1:A:386:VAL:O	1:A:427:CYS:HA	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:PHE:CD2	1:A:179:GLN:HG3	2.54	0.43
1:A:351:PRO:HG3	6:A:614:HOH:O	2.19	0.42
1:A:466:THR:HG23	1:A:468:LYS:O	2.20	0.42
1:A:377:HIS:O	1:A:378:LYS:HB2	2.20	0.42
1:A:141:ILE:O	1:A:142:ARG:HG3	2.19	0.42
1:A:276:GLN:HG3	1:A:281:GLN:HE21	1.84	0.41
1:A:276:GLN:HG3	1:A:281:GLN:NE2	2.35	0.41
1:A:28:HIS:HE1	6:A:603:HOH:O	2.04	0.41
1:A:29:PHE:H	3:A:503:DMS:C2	2.33	0.41
1:A:419:MET:SD	6:A:601:HOH:O	2.62	0.41
1:A:336:PHE:CE2	4:A:511:EDO:C1	3.04	0.41
1:A:148:ALA:HA	1:A:190:SER:HB3	2.02	0.40
1:A:417:MET:HE1	1:A:427:CYS:CA	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	452/475 (95%)	439 (97%)	11 (2%)	2 (0%)	34 30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	456	SER
1	A	455	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	392/418 (94%)	383 (98%)	9 (2%)	50 53

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	LYS
1	A	201	MET
1	A	209	ASP
1	A	238	HIS
1	A	296	PHE
1	A	303	ASP
1	A	323	SER
1	A	350	PHE
1	A	402	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	GLN
1	A	281	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	253	1	8,9,10	1.04	0	5,9,11	1.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	253	1	-	3/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	253	CME	CA-CB-SG-SD
1	A	253	CME	SD-CE-CZ-OH
1	A	253	CME	CZ-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	514	-	3,3,3	0.49	0	2,2,2	0.21	0
3	DMS	A	512	-	3,3,3	0.62	0	3,3,3	0.61	0
3	DMS	A	504	-	3,3,3	0.72	0	3,3,3	0.89	0
3	DMS	A	502	-	3,3,3	0.62	0	3,3,3	0.61	0
3	DMS	A	506	-	3,3,3	0.68	0	3,3,3	0.44	0
4	EDO	A	509	-	3,3,3	0.84	0	2,2,2	0.77	0
4	EDO	A	511	-	3,3,3	0.47	0	2,2,2	0.91	0
4	EDO	A	515	-	3,3,3	0.35	0	2,2,2	0.41	0
4	EDO	A	510	-	3,3,3	0.36	0	2,2,2	0.73	0
5	ACT	A	513	-	3,3,3	1.23	0	3,3,3	1.19	0
3	DMS	A	503	-	3,3,3	0.71	0	3,3,3	1.07	0
4	EDO	A	507	-	3,3,3	0.51	0	2,2,2	0.25	0
4	EDO	A	516	-	3,3,3	0.53	0	2,2,2	0.49	0
2	V6U	A	501	-	24,27,27	3.40	14 (58%)	29,38,38	2.44	9 (31%)
4	EDO	A	508	-	3,3,3	0.70	0	2,2,2	0.25	0
4	EDO	A	505	-	3,3,3	0.48	0	2,2,2	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	514	-	-	0/1/1/1	-
4	EDO	A	509	-	-	1/1/1/1	-
4	EDO	A	511	-	-	1/1/1/1	-
4	EDO	A	515	-	-	1/1/1/1	-
4	EDO	A	510	-	-	1/1/1/1	-
4	EDO	A	507	-	-	0/1/1/1	-
4	EDO	A	516	-	-	0/1/1/1	-
2	V6U	A	501	-	-	0/12/14/14	0/3/3/3
4	EDO	A	508	-	-	1/1/1/1	-
4	EDO	A	505	-	-	0/1/1/1	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	V6U	C12-C13	6.82	1.49	1.40
2	A	501	V6U	C17-C12	6.47	1.50	1.39
2	A	501	V6U	C14-C13	5.52	1.48	1.39
2	A	501	V6U	C15-C14	5.27	1.50	1.38
2	A	501	V6U	C09-N08	-4.53	1.31	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	V6U	C06-N05	3.89	1.42	1.35
2	A	501	V6U	O23-C22	3.84	1.41	1.33
2	A	501	V6U	C16-C15	3.77	1.48	1.38
2	A	501	V6U	C06-N20	3.66	1.41	1.33
2	A	501	V6U	C16-C17	3.61	1.46	1.38
2	A	501	V6U	C06-N01	2.83	1.40	1.35
2	A	501	V6U	C04-N05	2.82	1.39	1.34
2	A	501	V6U	O24-C22	-2.81	1.15	1.22
2	A	501	V6U	C02-N01	2.05	1.36	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	V6U	C04-N05-C06	8.71	121.30	116.34
2	A	501	V6U	C09-C04-N05	4.73	122.23	116.61
2	A	501	V6U	N20-C06-N05	3.97	123.42	117.25
2	A	501	V6U	N05-C06-N01	-3.31	120.23	125.42
2	A	501	V6U	C03-C04-N05	-2.96	118.83	122.35
2	A	501	V6U	C18-N07-N08	2.66	123.63	120.50
2	A	501	V6U	O23-C22-C12	2.50	116.44	112.30
2	A	501	V6U	C17-C12-C13	2.38	121.99	119.43
2	A	501	V6U	C03-C02-CL19	-2.25	116.07	118.88

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	508	EDO	O1-C1-C2-O2
4	A	510	EDO	O1-C1-C2-O2
4	A	509	EDO	O1-C1-C2-O2
4	A	515	EDO	O1-C1-C2-O2
4	A	511	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	511	EDO	1	0
5	A	513	ACT	1	0
3	A	503	DMS	1	0
4	A	516	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/475 (95%)	0.18	27 (5%) 21 20	25, 45, 94, 135	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	350	PHE	6.6
1	A	351	PRO	6.5
1	A	51	PHE	5.9
1	A	356	PRO	5.2
1	A	449	VAL	4.2
1	A	352	GLY	4.2
1	A	209	ASP	4.0
1	A	208	PRO	3.8
1	A	454	ALA	3.5
1	A	451	LYS	3.4
1	A	455	ASP	3.0
1	A	453	VAL	2.9
1	A	466	THR	2.7
1	A	306	LYS	2.6
1	A	468	LYS	2.5
1	A	50	GLY	2.5
1	A	52	THR	2.5
1	A	447	LYS	2.4
1	A	353	GLU	2.4
1	A	450	MET	2.4
1	A	185	ASN	2.3
1	A	457	GLY	2.3
1	A	211	ARG	2.2
1	A	49	SER	2.2
1	A	456	SER	2.1
1	A	8	PHE	2.1
1	A	142	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	253	10/11	0.93	0.12	32,57,91,92	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACT	A	513	4/4	0.42	0.43	83,84,99,99	0
4	EDO	A	508	4/4	0.62	0.21	69,83,84,86	0
4	EDO	A	510	4/4	0.85	0.19	60,72,77,80	0
4	EDO	A	511	4/4	0.85	0.17	58,70,73,75	0
4	EDO	A	505	4/4	0.85	0.19	70,84,85,86	0
4	EDO	A	509	4/4	0.86	0.19	48,58,61,61	0
4	EDO	A	514	4/4	0.88	0.16	64,77,82,83	0
3	DMS	A	512	4/4	0.90	0.16	85,102,104,104	0
4	EDO	A	515	4/4	0.90	0.37	98,118,119,119	0
4	EDO	A	507	4/4	0.90	0.21	59,71,77,78	0
4	EDO	A	516	4/4	0.94	0.15	43,52,56,56	0
3	DMS	A	503	4/4	0.95	0.23	46,55,59,59	0
3	DMS	A	506	4/4	0.96	0.22	78,94,95,95	0
3	DMS	A	502	4/4	0.96	0.15	77,93,93,93	0
2	V6U	A	501	25/25	0.98	0.12	25,33,44,53	0
3	DMS	A	504	4/4	0.98	0.15	59,75,75,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.