



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 10:46 PM EDT

PDB ID : 7CP6  
Title : Crystal structure of FqzB  
Authors : Hara, K.; Hashimoto, H.; Matsushita, T.; Kishimoto, S.; Watanabe, K.  
Deposited on : 2020-08-06  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	459	
1	B	459	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6880 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MAK1-like monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	420	3276	2079	595	592	10	0	0	0
1	B	423	3301	2094	603	594	10	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	119	ILE	LEU	engineered mutation	UNP A0A0J5T0B0
A	140	THR	ALA	engineered mutation	UNP A0A0J5T0B0
A	453	GLY	-	expression tag	UNP A0A0J5T0B0
A	454	HIS	-	expression tag	UNP A0A0J5T0B0
A	455	HIS	-	expression tag	UNP A0A0J5T0B0
A	456	HIS	-	expression tag	UNP A0A0J5T0B0
A	457	HIS	-	expression tag	UNP A0A0J5T0B0
A	458	HIS	-	expression tag	UNP A0A0J5T0B0
A	459	HIS	-	expression tag	UNP A0A0J5T0B0
B	119	ILE	LEU	engineered mutation	UNP A0A0J5T0B0
B	140	THR	ALA	engineered mutation	UNP A0A0J5T0B0
B	453	GLY	-	expression tag	UNP A0A0J5T0B0
B	454	HIS	-	expression tag	UNP A0A0J5T0B0
B	455	HIS	-	expression tag	UNP A0A0J5T0B0
B	456	HIS	-	expression tag	UNP A0A0J5T0B0
B	457	HIS	-	expression tag	UNP A0A0J5T0B0
B	458	HIS	-	expression tag	UNP A0A0J5T0B0
B	459	HIS	-	expression tag	UNP A0A0J5T0B0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	I	0	0
			3	3		
3	B	6	Total	I	0	0
			6	6		

- Molecule 4 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Hg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	99	Total	O	0	0
			99	99		
5	B	88	Total	O	0	0
			88	88		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MAK1-like monooxygenase



- Molecule 1: MAK1-like monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.89Å 120.18Å 68.05Å 90.00° 94.11° 90.00°	Depositor
Resolution (Å)	41.64 – 2.20 44.99 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (41.64-2.20) 99.8 (44.99-2.20)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.28 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.213 , 0.238 0.213 , 0.238	Depositor DCC
$R_{free}$ test set	2257 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.7	Xtrriage
Anisotropy	0.143	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6880	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, HG, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/3354	0.42	0/4558
1	B	0.23	0/3384	0.42	0/4603
All	All	0.23	0/6738	0.42	0/9161

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3276	0	3247	12	0
1	B	3301	0	3265	19	0
2	A	53	0	30	1	0
2	B	53	0	30	1	0
3	A	3	0	0	1	0
3	B	6	0	0	2	0
4	A	1	0	0	0	0
5	A	99	0	0	1	0
5	B	88	0	0	0	0
All	All	6880	0	6572	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:170:VAL:O	1:B:175:ARG:NH2	2.28	0.67
1:B:179:THR:HG23	1:B:181:GLN:H	1.63	0.64
1:A:438:PRO:HG2	1:A:441:GLY:HA3	1.81	0.62
1:A:170:VAL:O	1:A:175:ARG:NH2	2.35	0.60
1:A:405:TRP:CD2	1:A:438:PRO:HD3	2.42	0.55
1:B:301:ARG:O	1:B:373:GLN:NE2	2.40	0.55
1:B:249:THR:OG1	1:B:250:ASP:N	2.40	0.54
1:B:326:PRO:HA	3:B:505:IOD:I	2.79	0.53
1:A:93:ASP:HB3	1:A:99:ILE:HD11	1.92	0.52
1:A:326:PRO:HA	3:A:502:IOD:I	2.80	0.51
1:A:96:ASP:OD2	1:A:280:ARG:HD3	2.12	0.49
1:B:96:ASP:OD2	1:B:280:ARG:HD3	2.14	0.48
1:B:227:ASP:HA	1:B:385:TRP:CZ3	2.49	0.47
1:B:105:ALA:HA	1:B:106:GLY:HA3	1.60	0.47
1:B:367:PRO:C	1:B:437:ILE:HD11	2.35	0.47
1:B:93:ASP:HB3	1:B:99:ILE:HD11	1.97	0.46
1:B:142:TYR:CZ	1:B:174:ALA:HB2	2.49	0.46
1:B:88:HIS:NE2	1:B:103:GLU:HG2	2.31	0.46
1:B:338:ASP:OD1	1:B:365:ARG:NE	2.40	0.45
1:A:138:ARG:NH1	5:A:608:HOH:O	2.42	0.45
1:B:229:ILE:HB	1:B:275:TRP:CZ2	2.52	0.44
1:B:139:VAL:HG13	1:B:151:VAL:HB	2.01	0.43
1:B:411:CYS:HB2	3:B:507:IOD:I	2.89	0.43
1:B:140:THR:HG22	1:B:141:LYS:H	1.84	0.42
2:B:501:FAD:H9	2:B:501:FAD:H1'1	1.83	0.42
1:A:62:PRO:HD2	1:A:407:LEU:HD22	2.00	0.42
1:B:113:THR:OG1	1:B:114:ASN:N	2.52	0.41
1:A:168:ASP:OD1	1:A:168:ASP:N	2.51	0.41
1:B:106:GLY:HA2	1:B:107:ARG:HA	1.57	0.41
1:A:140:THR:OG1	1:A:141:LYS:N	2.54	0.41
1:A:113:THR:OG1	1:A:114:ASN:N	2.51	0.40
2:A:501:FAD:H1'1	2:A:501:FAD:H9	1.84	0.40
1:A:142:TYR:CZ	1:A:174:ALA:HB2	2.56	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/459 (89%)	401 (98%)	9 (2%)	0	100	100
1	B	417/459 (91%)	404 (97%)	13 (3%)	0	100	100
All	All	827/918 (90%)	805 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/371 (91%)	338 (100%)	0	100	100
1	B	340/371 (92%)	340 (100%)	0	100	100
All	All	678/742 (91%)	678 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	B	501	1	53,58,58	2.44	24 (45%)	68,89,89	1.49	11 (16%)
2	FAD	A	501	-	53,58,58	2.26	22 (41%)	68,89,89	1.27	8 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	501	1	-	3/30/50/50	0/6/6/6
2	FAD	A	501	-	-	8/30/50/50	0/6/6/6

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	FAD	C4-N3	-5.21	1.29	1.38
2	B	501	FAD	C5X-N5	-4.76	1.30	1.39
2	A	501	FAD	P-O2P	-4.73	1.33	1.55
2	B	501	FAD	PA-O1A	-4.38	1.35	1.50
2	A	501	FAD	P-O1P	-4.23	1.35	1.50
2	A	501	FAD	C4-N3	-4.21	1.31	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	FAD	PA-O2A	-4.10	1.36	1.55
2	B	501	FAD	C9-C8	-3.97	1.33	1.39
2	B	501	FAD	P-O2P	-3.97	1.36	1.55
2	B	501	FAD	C4A-N3A	-3.86	1.30	1.35
2	A	501	FAD	C4A-N3A	-3.80	1.30	1.35
2	B	501	FAD	P-O1P	-3.74	1.37	1.50
2	A	501	FAD	PA-O2A	-3.73	1.37	1.55
2	A	501	FAD	PA-O1A	-3.39	1.38	1.50
2	A	501	FAD	C9A-C5X	3.36	1.46	1.41
2	A	501	FAD	C9-C8	-3.36	1.34	1.39
2	B	501	FAD	C2B-C1B	-3.31	1.48	1.53
2	A	501	FAD	C5X-N5	-3.29	1.33	1.39
2	A	501	FAD	C2-N3	-3.29	1.31	1.39
2	A	501	FAD	O2'-C2'	-3.28	1.36	1.43
2	B	501	FAD	O4-C4	-3.19	1.17	1.23
2	B	501	FAD	C2-N1	-3.15	1.29	1.36
2	B	501	FAD	C9-C9A	-3.10	1.34	1.39
2	A	501	FAD	C2A-N1A	-3.07	1.28	1.33
2	B	501	FAD	O4B-C4B	-2.97	1.38	1.45
2	B	501	FAD	O2-C2	-2.93	1.18	1.24
2	B	501	FAD	C2A-N1A	-2.88	1.28	1.33
2	B	501	FAD	C9A-N10	-2.87	1.36	1.41
2	B	501	FAD	C6-C7	-2.83	1.35	1.39
2	B	501	FAD	O4'-C4'	-2.81	1.37	1.43
2	A	501	FAD	C5A-N7A	-2.70	1.29	1.39
2	A	501	FAD	C6-C7	-2.65	1.35	1.39
2	B	501	FAD	C2-N3	-2.64	1.32	1.39
2	A	501	FAD	C8A-N7A	-2.60	1.30	1.34
2	A	501	FAD	O4B-C4B	-2.52	1.39	1.45
2	B	501	FAD	C5A-N7A	-2.46	1.30	1.39
2	A	501	FAD	O3'-C3'	-2.46	1.37	1.43
2	A	501	FAD	C2B-C1B	-2.45	1.50	1.53
2	A	501	FAD	C6-C5X	-2.33	1.36	1.40
2	B	501	FAD	O3'-C3'	-2.23	1.37	1.43
2	A	501	FAD	C1'-N10	-2.19	1.42	1.48
2	B	501	FAD	C8A-N7A	-2.14	1.30	1.34
2	A	501	FAD	P-O5'	-2.12	1.50	1.59
2	B	501	FAD	C9A-C5X	2.09	1.44	1.41
2	A	501	FAD	O4-C4	-2.09	1.19	1.23
2	B	501	FAD	O2B-C2B	-2.02	1.38	1.43

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	C4A-C5A-N7A	-4.25	104.97	109.40
2	B	501	FAD	C4A-C5A-N7A	-4.01	105.22	109.40
2	B	501	FAD	O2-C2-N1	-3.57	115.90	121.83
2	B	501	FAD	N3A-C2A-N1A	-3.04	123.92	128.68
2	A	501	FAD	C4-C4X-N5	2.92	122.38	118.23
2	B	501	FAD	O5'-C5'-C4'	2.89	117.07	109.36
2	A	501	FAD	O4-C4-C4X	-2.81	119.13	126.60
2	B	501	FAD	C4-C4X-N5	2.67	122.04	118.23
2	B	501	FAD	C4-N3-C2	-2.67	120.70	125.64
2	B	501	FAD	O4-C4-C4X	-2.62	119.64	126.60
2	B	501	FAD	N3-C2-N1	2.48	124.26	119.38
2	A	501	FAD	N3A-C2A-N1A	-2.48	124.80	128.68
2	B	501	FAD	O4B-C1B-C2B	-2.40	103.42	106.93
2	A	501	FAD	C10-N1-C2	2.39	121.69	116.90
2	B	501	FAD	C4X-C10-N1	-2.34	119.30	124.73
2	A	501	FAD	O2'-C2'-C1'	-2.27	104.30	109.80
2	B	501	FAD	C4X-C4-N3	2.25	118.91	113.19
2	A	501	FAD	C4X-C4-N3	2.21	118.80	113.19
2	A	501	FAD	C4X-C10-N1	-2.17	119.70	124.73

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	FAD	O2'-C2'-C3'-C4'
2	A	501	FAD	O2'-C2'-C3'-O3'
2	A	501	FAD	PA-O3P-P-O5'
2	B	501	FAD	PA-O3P-P-O5'
2	A	501	FAD	O4B-C4B-C5B-O5B
2	A	501	FAD	P-O3P-PA-O1A
2	A	501	FAD	C3B-C4B-C5B-O5B
2	A	501	FAD	P-O3P-PA-O2A
2	B	501	FAD	O4B-C4B-C5B-O5B
2	A	501	FAD	C1'-C2'-C3'-O3'
2	B	501	FAD	C1'-C2'-C3'-O3'


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	FAD	1	0
2	A	501	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	420/459 (91%)	0.05	8 (1%) 66 65	22, 36, 62, 88	0
1	B	423/459 (92%)	0.30	26 (6%) 21 20	23, 42, 70, 89	0
All	All	843/918 (91%)	0.17	34 (4%) 38 36	22, 39, 69, 89	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	422	VAL	4.6
1	B	426	VAL	4.6
1	A	441	GLY	4.3
1	B	437	ILE	4.1
1	A	440	ASP	3.9
1	B	107	ARG	3.8
1	B	7	LEU	3.8
1	B	402	ASN	3.3
1	B	190	TRP	3.2
1	B	106	GLY	3.1
1	A	104	LEU	3.1
1	B	396	SER	2.9
1	A	442	ALA	2.8
1	B	432	TYR	2.8
1	B	429	GLU	2.6
1	B	140	THR	2.5
1	B	418	VAL	2.5
1	A	443	TYR	2.5
1	B	431	GLU	2.5
1	A	396	SER	2.5
1	B	105	ALA	2.4
1	A	102	HIS	2.3
1	B	397	VAL	2.2
1	B	380	GLN	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	419	PHE	2.2
1	B	146	ASP	2.2
1	B	400	PHE	2.2
1	B	325	PHE	2.1
1	B	256	GLU	2.1
1	B	369	ALA	2.1
1	B	295	ASP	2.1
1	B	383	LEU	2.0
1	A	429	GLU	2.0
1	B	6	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	HG	A	505	1/1	-0.57	0.23	302,302,302,302	0
3	IOD	A	503	1/1	0.43	0.16	182,182,182,182	0
3	IOD	B	503	1/1	0.89	0.08	107,107,107,107	0
3	IOD	B	506	1/1	0.96	0.09	53,53,53,53	0
3	IOD	B	507	1/1	0.96	0.05	53,53,53,53	0
2	FAD	B	501	53/53	0.96	0.10	27,33,37,39	0
3	IOD	B	504	1/1	0.97	0.08	68,68,68,68	0
2	FAD	A	501	53/53	0.97	0.10	21,30,39,40	0
3	IOD	B	505	1/1	0.98	0.10	46,46,46,46	0
3	IOD	A	502	1/1	0.99	0.11	41,41,41,41	0
3	IOD	B	502	1/1	0.99	0.06	66,66,66,66	0
3	IOD	A	504	1/1	1.00	0.08	43,43,43,43	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.