

# wwPDB X-ray Structure Validation Summary Report (i)

Sep 18, 2023 – 07:19 pm BST

PDB ID : 8CQL

Title: pVHL:EloB:EloC in complex with (2S,4R)-N-((S)-1-(5-Fluoro-2-methoxy-4-(4

-methylthiazol-5-yl)phenyl)ethyl)-1-((S)-2-(1-fluorocyclopropane-1-carboxami do)-3,3-dimethylbutanoyl)-4-hydroxypyrrolidine-2-carboxamide (Compound

33)

Authors: Casement, R.; Phuong Vu, L.; Ciulli, A.; Gutschow, M.

 $Deposited \ on \quad : \quad 2023\text{-}03\text{-}06$ 

Resolution : 2.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$ 

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

 $\begin{tabular}{lll} CCP4 & : & 7.0.044 & (Gargrove) \end{tabular}$ 

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1




## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned}  ext{Similar resolution} \ (\# ext{Entries, resolution range}(\mathring{ ext{A}})) \end{aligned}$		
$R_{free}$	130704	5509 (2.40-2.36)		
Clashscore	141614	6082 (2.40-2.36)		
Ramachandran outliers	138981	5973 (2.40-2.36)		
Sidechain outliers	138945	5975 (2.40-2.36)		
RSRZ outliers	127900	5397 (2.40-2.36)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	104	83%	17%					
1	D	104	89%	11%					
1	G	104	85%	15%					
1	J	104	83%	16%					
2	В	97	73% 18%	9%					

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain		
2	Е	97	80%	11%	8%
2	Н	97	77%	11%	11%
2	K	97	79%	9%	11%
3	С	162	72%	15% •	12%
3	F	162	% • 78%	10%	12%
3	I	162	72%	15%	12%
3	L	162	77%	12%	12%



## 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10737 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Elongin-B.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
1	Λ	104	Total	As	С	N	О	S	0	0	0
1	A	104	803	2	510	133	154	4	0	U	
1	D	104	Total	As	С	N	О	S	0	0	0
1	ש	104	813	2	515	137	154	5			
1	G	104	Total	As	С	N	О	S	0	0	0
1	G	104	764	2	488	127	143	4			U
1	1 J	J 104	Total	As	С	N	О	S	0	0	0
1			797	2	507	135	149	4	U	U	U

• Molecule 2 is a protein called Elongin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	В 88	Total	С	N	О	S	0	0	0
2	Б		682	443	108	124	7	0	0	
2	E	89	Total	С	N	О	S	0	0	0
2	<u> 1</u> 2	09	697	450	113	128	6			U
2	Н	86	Total	С	N	О	S	0	0	0
2	11	80	670	435	106	124	5			
9	2 K	86	Total	С	N	О	S	0	0	0
			666	432	105	124	5	U		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	16	MET	-	initiating methionine	<u> </u>
E	16	MET	-	initiating methionine	UNP Q15369
Н	16	MET	-	initiating methionine	UNP Q15369
K	16	MET	-	initiating methionine	UNP Q15369

• Molecule 3 is a protein called von Hippel-Lindau disease tumor suppressor.




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	143	Total	As	С	N	О	S	0	0	0
3		143	1128	1	719	207	199	2		0	0
3	F	143	Total	As	С	N	О	S	0	0	0
3	I.	140	1134	1	723	205	203	2	0		
3	Т	142	Total	As	С	N	О	S	0	0	0
3	1	142	1086	1	694	190	199	2			0
2	3 L	143	Total	As	С	N	О	S	0	0	0
3			1105	1	704	197	201	2	0		U

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	52	GLY	-	expression tag	UNP P40337
С	53	SER	-	expression tag	UNP P40337
F	52	GLY	-	expression tag	UNP P40337
F	53	SER	-	expression tag	UNP P40337
I	52	GLY	-	expression tag	UNP P40337
I	53	SER	-	expression tag	UNP P40337
L	52	GLY	-	expression tag	UNP P40337
L	53	SER	-	expression tag	UNP P40337

• Molecule 4 is (2 {S},4 {R})-1-[(2 {S})-2-[(1-fluoranylcyclopropyl)carbonylamino]-3,3-d imethyl-butanoyl]- {N}-[(1 {S})-1-[5-fluoranyl-2-methoxy-4-(4-methyl-1,3-thiazol-5-yl )phenyl]ethyl]-4-oxidanyl-pyrrolidine-2-carboxamide (three-letter code: VH3) (formula:  $C_{28}H_{36}F_2N_4O_5S$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
1	С	1	Total	С	F	N	О	S	0	0	
4	4	1	40	28	2	4	5	1			
1	F	1	Total C F N O S		0	0					
4	4 Г	1	40	28	2	4	5	1			
1	Т	1	Total	С	F	N	О	S	0	0	
4	$4 \mid 1$	1	40	28	2	4	5	1	0	0	
1	Т	1	Total	С	F	N	О	S	0	0	
4	ь	1	40	28	2	4	5	1		U	


#### • Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	17	Total O	0	0
	Λ	11	17 17	U	U
5	В	14	Total O	0	0
	Б	11	14 14	Ů	
5	С	31	Total O	0	0
		91	31 31		Ŭ
5	D	14	Total O	0	0
	_		14 14		Ū.
5	E	23	Total O	0	0
		_	23 23	_	_
5	F	24	Total O	0	0
			24 24		
5	G	10	Total O	0	0
			10 10		
5	Н	11	Total O	0	0
			11 11 Total O		
5	I	31	31 31	0	0
			Total O		
5	J	25	25 $25$	0	0
			Total O		
5	K	12	10 tai 0 12 12	0	0
			Total O		
5	L	20	20 20	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.















## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	92.69Å 92.69Å 359.62Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.53 - 2.38	Depositor
resolution (A)	89.91 - 2.38	EDS
% Data completeness	99.5 (64.53-2.38)	Depositor
(in resolution range)	99.6 (89.91-2.38)	EDS
$R_{merge}$	0.26	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.61 (at 2.37Å)	Xtriage
Refinement program	PHENIX 1.18.2	Depositor
P.P.	0.226 , $0.275$	Depositor
$R, R_{free}$	0.230 , $0.269$	DCC
$R_{free}$ test set	3241  reflections  (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.7	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33, 43.9	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10737	wwPDB-VP
Average B, all atoms $(Å^2)$	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 32.07 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.9857e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, VH3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles	
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.35	0/799	0.59	0/1080	
1	D	0.35	0/808	0.61	0/1088	
1	G	0.36	0/760	0.63	0/1031	
1	J	0.34	0/793	0.57	0/1072	
2	В	0.33	0/696	0.56	0/940	
2	Е	0.32	0/711	0.51	0/960	
2	Н	0.31	0/684	0.49	0/925	
2	K	0.33	0/680	0.52	0/921	
3	С	0.34	0/1148	0.55	0/1570	
3	F	0.36	0/1154	0.57	0/1577	
3	I	0.35	0/1105	0.54	0/1518	
3	L	0.36	0/1124	0.57	0/1540	
All	All	0.34	0/10462	0.56	0/14222	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes	
1	A	803	0	775	12	0	

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
1	D	813	0	796	9	0
1	G	764	0	711	14	0
1	J	797	0	771	15	0
2	В	682	0	673	17	0
2	Е	697	0	687	11	0
2	Н	670	0	659	11	0
2	K	666	0	648	10	0
3	С	1128	0	1086	18	0
3	F	1134	0	1094	12	0
3	I	1086	0	1015	17	0
3	L	1105	0	1041	16	0
4	С	40	0	0	0	0
4	F	40	0	0	0	0
4	I	40	0	0	0	0
4	L	40	0	0	0	0
5	A	17	0	0	0	0
5	В	14	0	0	1	0
5	С	31	0	0	1	0
5	D	14	0	0	1	0
5	Ε	23	0	0	0	0
5	F	24	0	0	0	0
5	G	10	0	0	0	0
5	Н	11	0	0	3	0
5	I	31	0	0	1	0
5	J	25	0	0	0	0
5	K	12	0	0	1	0
5	L	20	0	0	0	0
All	All	10737	0	9956	131	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 131 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
2:H:97:PRO:HB3	3:I:169:LEU:HD21	1.60	0.82
3:F:73:GLN:H	3:F:141:ASN:HD21	1.37	0.71
3:L:73:GLN:OE1	3:L:108:ARG:NH2	2.22	0.71
1:G:94:SER:HG	2:H:68:HIS:HD1	1.35	0.71
2:E:63:ARG:HD2	3:L:104:GLY:HA3	1.72	0.70



There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	100/104~(96%)	95 (95%)	5 (5%)	0	100	100
1	D	$100/104\ (96\%)$	96 (96%)	4 (4%)	0	100	100
1	G	100/104~(96%)	95 (95%)	5 (5%)	0	100	100
1	J	$100/104\ (96\%)$	95 (95%)	4 (4%)	1 (1%)	15	21
2	В	84/97~(87%)	84 (100%)	0	0	100	100
2	E	85/97 (88%)	85 (100%)	0	0	100	100
2	Н	82/97~(84%)	82 (100%)	0	0	100	100
2	K	82/97~(84%)	82 (100%)	0	0	100	100
3	$\mathbf{C}$	140/162~(86%)	137 (98%)	3 (2%)	0	100	100
3	F	140/162~(86%)	138 (99%)	2 (1%)	0	100	100
3	I	$139/162\ (86\%)$	136 (98%)	3 (2%)	0	100	100
3	L	140/162 (86%)	138 (99%)	2 (1%)	0	100	100
All	All	1292/1452~(89%)	1263 (98%)	28 (2%)	1 (0%)	51	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	J	82	ASP

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	83/90 (92%)	83 (100%)	0	100	100
1	D	85/90 (94%)	85 (100%)	0	100	100
1	G	73/90 (81%)	73 (100%)	0	100	100
1	J	81/90 (90%)	80 (99%)	1 (1%)	71	84
2	В	74/86 (86%)	74 (100%)	0	100	100
2	$\mathbf{E}$	76/86 (88%)	76 (100%)	0	100	100
2	Н	73/86 (85%)	73 (100%)	0	100	100
2	K	72/86 (84%)	72 (100%)	0	100	100
3	С	120/147 (82%)	118 (98%)	2 (2%)	60	76
3	F	122/147 (83%)	122 (100%)	0	100	100
3	I	114/147 (78%)	114 (100%)	0	100	100
3	L	116/147 (79%)	116 (100%)	0	100	100
All	All	1089/1292 (84%)	1086 (100%)	3 (0%)	92	97

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	С	113	ARG
3	С	148	PHE
1	J	50	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
3	I	96	GLN
1	J	49	GLN
3	L	132	GLN
1	D	49	GLN
3	С	110	HIS

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trino	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	gles
Mol	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CAS	L	77	3	5,8,9	0.99	0	1,9,11	1.14	0
1	CAS	D	89	1	5,8,9	1.09	0	1,9,11	0.22	0
1	CAS	D	60	1	5,8,9	1.10	0	1,9,11	1.17	0
3	CAS	F	77	3	5,8,9	1.15	0	1,9,11	0.91	0
1	CAS	G	89	1	5,8,9	1.18	0	1,9,11	0.96	0
3	CAS	С	77	3	5,8,9	0.97	0	1,9,11	0.71	0
3	CAS	I	77	3	5,8,9	1.11	0	1,9,11	1.41	0
1	CAS	G	60	1	5,8,9	1.05	0	1,9,11	0.38	0
1	CAS	J	60	1	5,8,9	1.05	0	1,9,11	1.12	0
1	CAS	A	89	1	5,8,9	1.20	0	1,9,11	1.18	0
1	CAS	J	89	1	5,8,9	1.16	0	1,9,11	0.72	0
1	CAS	A	60	1	5,8,9	1.06	0	1,9,11	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CAS	L	77	3	-	0/0/7/9	-
1	CAS	D	89	1	-	0/0/7/9	-
1	CAS	D	60	1	-	0/0/7/9	-
3	CAS	F	77	3	-	0/0/7/9	-
1	CAS	G	89	1	-	0/0/7/9	-
3	CAS	С	77	3	-	0/0/7/9	-
3	CAS	I	77	3	-	0/0/7/9	-
1	CAS	G	60	1	-	0/0/7/9	-
1	CAS	J	60	1	-	0/0/7/9	-
1	CAS	A	89	1	-	0/0/7/9	-
1	CAS	J	89	1	-	0/0/7/9	-
1	CAS	A	60	1	-	0/0/7/9	-



There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	J	89	CAS	1	0

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain	n Res	Link	В	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
4	VH3	F	301	-	36,43,43	4.07	16 (44%)	48,66,66	1.49	8 (16%)	
4	VH3	С	301	-	36,43,43	4.07	16 (44%)	48,66,66	1.48	7 (14%)	
4	VH3	I	301	-	36,43,43	4.10	16 (44%)	48,66,66	1.49	7 (14%)	
4	VH3	L	301	-	36,43,43	4.07	16 (44%)	48,66,66	1.47	9 (18%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	VH3	F	301	-	-	3/39/58/58	0/4/4/4

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	VH3	С	301	-	-	3/39/58/58	0/4/4/4
4	VH3	I	301	-	-	4/39/58/58	0/4/4/4
4	VH3	L	301	-	-	5/39/58/58	0/4/4/4

The worst 5 of 64 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\operatorname{\mathring{A}})$	Ideal(A)
4	L	301	VH3	C03-C04	-10.68	1.33	1.53
4	I	301	VH3	C40-C02	-10.64	1.37	1.52
4	F	301	VH3	C03-C04	-10.64	1.33	1.53
4	С	301	VH3	C03-C04	-10.58	1.33	1.53
4	L	301	VH3	C40-C02	-10.57	1.37	1.52

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	I	301	VH3	O10-C09-C08	5.01	120.62	115.83
4	С	301	VH3	O10-C09-C08	4.67	120.30	115.83
4	F	301	VH3	O10-C09-C08	4.62	120.25	115.83
4	L	301	VH3	O10-C09-C08	4.55	120.18	115.83
4	L	301	VH3	C40-N25-C04	-4.35	105.42	111.70

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	С	301	VH3	C20-C13-C14-S19
4	F	301	VH3	C20-C13-C14-S19
4	I	301	VH3	C20-C13-C14-S19
4	L	301	VH3	C20-C13-C14-S19
4	С	301	VH3	C12-C09-O10-C11


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be




highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	A	102/104 (98%)	0.06	2 (1%) 65 66	34, 53, 71, 95	0
1	D	102/104 (98%)	0.02	2 (1%) 65 66	31, 43, 68, 87	0
1	G	102/104 (98%)	0.32	3 (2%) 51 53	38, 60, 87, 100	0
1	J	102/104 (98%)	0.10	1 (0%) 82 83	33, 46, 73, 87	0
2	В	88/97 (90%)	0.11	1 (1%) 80 81	34, 45, 68, 75	0
2	E	89/97 (91%)	0.08	0 100 100	31, 41, 64, 80	0
2	Н	86/97 (88%)	0.16	0 100 100	37, 51, 63, 70	0
2	K	86/97 (88%)	0.00	0 100 100	33, 43, 63, 70	0
3	С	142/162 (87%)	0.02	0 100 100	31, 42, 64, 89	0
3	F	142/162 (87%)	0.07	1 (0%) 87 88	26, 42, 60, 81	0
3	I	141/162 (87%)	0.16	1 (0%) 87 88	31, 44, 76, 90	0
3	L	142/162 (87%)	0.06	2 (1%) 75 77	26, 42, 69, 81	0
All	All	1324/1452 (91%)	0.10	13 (0%) 82 83	26, 45, 72, 100	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	102	VAL	4.8
3	I	182	ARG	4.0
3	L	180	ILE	2.8
1	A	81	ALA	2.7
2	В	16	MET	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
1	CAS	D	60	9/10	0.86	0.18	46,51,69,135	0
1	CAS	G	89	9/10	0.87	0.14	70,78,99,147	0
1	CAS	J	60	9/10	0.88	0.13	32,39,58,129	0
1	CAS	A	89	9/10	0.89	0.18	48,53,108,133	0
1	CAS	A	60	9/10	0.89	0.20	47,56,75,126	0
1	CAS	J	89	9/10	0.93	0.14	48,53,88,118	0
1	CAS	G	60	9/10	0.94	0.14	42,50,76,115	0
1	CAS	D	89	9/10	0.97	0.11	43,47,87,88	0
3	CAS	I	77	9/10	0.97	0.12	36,41,61,67	0
3	CAS	L	77	9/10	0.97	0.12	35,40,53,64	0
3	CAS	С	77	9/10	0.98	0.11	34,44,58,66	0
3	CAS	F	77	9/10	0.98	0.11	32,39,61,62	0

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	VH3	I	301	40/40	0.94	0.14	35,45,54,64	0
4	VH3	С	301	40/40	0.95	0.17	30,40,51,56	0
4	VH3	L	301	40/40	0.95	0.14	27,40,46,51	0
4	VH3	F	301	40/40	0.96	0.16	26,37,46,49	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around VH3 I 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around VH3 C 301:

 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












#### Electron density around VH3 L 301:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around VH3 F 301:

 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









## 6.5 Other polymers (i)

There are no such residues in this entry.

