



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 06:32 pm GMT

PDB ID : 5CSU
Title : Disproportionating enzyme 1 from Arabidopsis - acarviostatin soak
Authors : O'Neill, E.C.; Stevenson, C.E.M.; Tantanarat, K.; Latousakis, D.; Donaldson, M.I.; Rejzek, M.; Limpaseni, T.; Smith, A.M.; Field, R.A.; Lawson, D.M.
Deposited on : 2015-07-23
Resolution : 2.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	564	 3% 88% 8%
1	B	564	 % 88% 8%
2	C	2	 100%
2	E	2	 100%
3	D	3	 33% 67%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	3	 33% 67%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-alpha-glucanotransferase DPE1, chloroplastic/amyloplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	518	3973	2554	676	731	12	0	0	0
1	B	518	4019	2580	684	743	12	0	0	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	MET	-	initiating methionine	UNP Q9LV91
A	14	HIS	-	expression tag	UNP Q9LV91
A	15	HIS	-	expression tag	UNP Q9LV91
A	16	HIS	-	expression tag	UNP Q9LV91
A	17	HIS	-	expression tag	UNP Q9LV91
A	18	HIS	-	expression tag	UNP Q9LV91
A	19	HIS	-	expression tag	UNP Q9LV91
A	20	GLY	-	expression tag	UNP Q9LV91
A	21	LYS	-	expression tag	UNP Q9LV91
A	22	PRO	-	expression tag	UNP Q9LV91
A	23	ILE	-	expression tag	UNP Q9LV91
A	24	PRO	-	expression tag	UNP Q9LV91
A	25	ASN	-	expression tag	UNP Q9LV91
A	26	PRO	-	expression tag	UNP Q9LV91
A	27	LEU	-	expression tag	UNP Q9LV91
A	28	LEU	-	expression tag	UNP Q9LV91
A	29	GLY	-	expression tag	UNP Q9LV91
A	30	LEU	-	expression tag	UNP Q9LV91
A	31	ASP	-	expression tag	UNP Q9LV91
A	32	SER	-	expression tag	UNP Q9LV91
A	33	THR	-	expression tag	UNP Q9LV91
A	34	GLU	-	expression tag	UNP Q9LV91
A	35	ASN	-	expression tag	UNP Q9LV91
A	36	LEU	-	expression tag	UNP Q9LV91
A	37	TYR	-	expression tag	UNP Q9LV91

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	PHE	-	expression tag	UNP Q9LV91
A	39	GLN	-	expression tag	UNP Q9LV91
A	40	GLY	-	expression tag	UNP Q9LV91
A	41	ILE	-	expression tag	UNP Q9LV91
A	42	ASP	-	expression tag	UNP Q9LV91
A	43	PRO	-	expression tag	UNP Q9LV91
A	44	PHE	-	expression tag	UNP Q9LV91
A	45	THR	-	expression tag	UNP Q9LV91
B	13	MET	-	initiating methionine	UNP Q9LV91
B	14	HIS	-	expression tag	UNP Q9LV91
B	15	HIS	-	expression tag	UNP Q9LV91
B	16	HIS	-	expression tag	UNP Q9LV91
B	17	HIS	-	expression tag	UNP Q9LV91
B	18	HIS	-	expression tag	UNP Q9LV91
B	19	HIS	-	expression tag	UNP Q9LV91
B	20	GLY	-	expression tag	UNP Q9LV91
B	21	LYS	-	expression tag	UNP Q9LV91
B	22	PRO	-	expression tag	UNP Q9LV91
B	23	ILE	-	expression tag	UNP Q9LV91
B	24	PRO	-	expression tag	UNP Q9LV91
B	25	ASN	-	expression tag	UNP Q9LV91
B	26	PRO	-	expression tag	UNP Q9LV91
B	27	LEU	-	expression tag	UNP Q9LV91
B	28	LEU	-	expression tag	UNP Q9LV91
B	29	GLY	-	expression tag	UNP Q9LV91
B	30	LEU	-	expression tag	UNP Q9LV91
B	31	ASP	-	expression tag	UNP Q9LV91
B	32	SER	-	expression tag	UNP Q9LV91
B	33	THR	-	expression tag	UNP Q9LV91
B	34	GLU	-	expression tag	UNP Q9LV91
B	35	ASN	-	expression tag	UNP Q9LV91
B	36	LEU	-	expression tag	UNP Q9LV91
B	37	TYR	-	expression tag	UNP Q9LV91
B	38	PHE	-	expression tag	UNP Q9LV91
B	39	GLN	-	expression tag	UNP Q9LV91
B	40	GLY	-	expression tag	UNP Q9LV91
B	41	ILE	-	expression tag	UNP Q9LV91
B	42	ASP	-	expression tag	UNP Q9LV91
B	43	PRO	-	expression tag	UNP Q9LV91
B	44	PHE	-	expression tag	UNP Q9LV91
B	45	THR	-	expression tag	UNP Q9LV91


- Molecule 2 is an oligosaccharide called 4-amino-4,6-dideoxy-alpha-D-glucopyranose-(1-4)-al

pha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	Total	C	N	O	0	0	0
			21	12	1	8			
2	E	2	Total	C	N	O	0	0	0
			21	12	1	8			

- Molecule 3 is an oligosaccharide called 4-amino-4,6-dideoxy-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	3	Total	C	N	O	0	0	0
			33	18	1	14			
3	F	3	Total	C	N	O	0	0	0
			33	18	1	14			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is 5-HYDROXYMETHYL-CHONDURITOL (three-letter code: HMC) (formula: $C_7H_{12}O_5$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 11 7 4	0	0
5	A	1	Total C O 11 7 4	0	0
5	B	1	Total C O 11 7 4	0	0
5	B	1	Total C O 11 7 4	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	93	Total O 93 93	0	0
6	B	107	Total O 107 107	0	0

Chain D:  33% 67%

GLC1
GLC2
AGL3

- Molecule 3: 4-amino-4,6-dideoxy-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain F:  33% 67%

GLC1
GLC2
AGL3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	70.83Å 74.13Å 79.73Å 64.59° 69.76° 66.57°	Depositor
Resolution (Å)	40.95 – 2.53 40.95 – 2.53	Depositor EDS
% Data completeness (in resolution range)	95.7 (40.95-2.53) 95.7 (40.95-2.53)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.54Å)	Xtrriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.207 , 0.249 0.211 , 0.250	Depositor DCC
R_{free} test set	2181 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtrriage
Anisotropy	0.416	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8356	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AGL, EDO, BGC, GLC, HMC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/4086	0.72	2/5576 (0.0%)
1	B	0.55	0/4133	0.73	2/5631 (0.0%)
All	All	0.54	0/8219	0.73	4/11207 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	478	ARG	NE-CZ-NH1	9.06	124.83	120.30
1	B	478	ARG	NE-CZ-NH2	-7.22	116.69	120.30
1	A	376	ARG	NE-CZ-NH1	6.48	123.54	120.30
1	A	376	ARG	NE-CZ-NH2	-5.66	117.47	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3973	0	3733	6	0
1	B	4019	0	3810	6	0
2	C	21	0	18	0	0
2	E	21	0	18	0	0
3	D	33	0	29	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	33	0	29	1	0
4	A	8	0	12	0	0
4	B	4	0	6	0	0
5	A	22	0	19	0	0
5	B	22	0	19	0	0
6	A	93	0	0	0	0
6	B	107	0	0	0	0
All	All	8356	0	7693	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:TRP:O	1:A:349:GLU:HG3	2.09	0.52
1:A:59:ILE:HG22	1:A:63:GLU:HB2	1.92	0.51
1:B:163:ASP:OD1	1:B:163:ASP:N	2.45	0.48
1:B:246:ARG:HG2	1:B:306:TRP:CZ2	2.52	0.45
1:B:478:ARG:HH11	1:B:478:ARG:CG	2.29	0.45
1:B:422:LEU:HD11	1:B:441:MET:HE2	1.99	0.45
1:A:246:ARG:HG2	1:A:306:TRP:CZ2	2.52	0.44
1:A:422:LEU:HD11	1:A:441:MET:HE2	2.00	0.43
1:A:296:ILE:HG23	1:A:297:TYR:CD2	2.54	0.43
1:B:296:ILE:HG23	1:B:297:TYR:CD2	2.55	0.42
1:A:124:LEU:HB3	1:A:125:PRO:HD2	2.02	0.41
1:B:423:GLY:HA2	3:F:2:GLC:H3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	516/564 (92%)	505 (98%)	11 (2%)	0	100	100
1	B	516/564 (92%)	504 (98%)	11 (2%)	1 (0%)	47	67
All	All	1032/1128 (92%)	1009 (98%)	22 (2%)	1 (0%)	51	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	499	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	394/480 (82%)	385 (98%)	9 (2%)	50	74
1	B	406/480 (85%)	392 (97%)	14 (3%)	37	61
All	All	800/960 (83%)	777 (97%)	23 (3%)	42	67

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	GLU
1	A	216	CYS
1	A	248	LEU
1	A	309	LYS
1	A	376	ARG
1	A	395	ARG
1	A	462	GLU
1	A	533	SER
1	A	536	MET
1	B	60	SER
1	B	130	ASP
1	B	163	ASP
1	B	248	LEU
1	B	309	LYS
1	B	311	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	349	GLU
1	B	376	ARG
1	B	392	MET
1	B	462	GLU
1	B	478	ARG
1	B	482	ASP
1	B	533	SER
1	B	536	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	C	1	2,5	11,11,12	0.36	0	15,15,17	1.11	1 (6%)
2	AGL	C	2	2,5	9,10,11	0.40	0	14,14,16	1.03	1 (7%)
3	BGC	D	1	3	12,12,12	0.29	0	17,17,17	0.47	0
3	GLC	D	2	3	11,11,12	0.43	0	15,15,17	0.79	1 (6%)
3	AGL	D	3	3,5	9,10,11	0.41	0	14,14,16	1.94	3 (21%)
2	GLC	E	1	2,5	11,11,12	0.38	0	15,15,17	1.04	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AGL	E	2	2,5	9,10,11	0.45	0	14,14,16	1.03	1 (7%)
3	BGC	F	1	3	12,12,12	0.31	0	17,17,17	0.59	0
3	GLC	F	2	3	11,11,12	0.45	0	15,15,17	0.74	0
3	AGL	F	3	3,5	9,10,11	0.42	0	14,14,16	1.90	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2,5	-	2/2/19/22	0/1/1/1
2	AGL	C	2	2,5	-	-	0/1/1/1
3	BGC	D	1	3	-	0/2/22/22	0/1/1/1
3	GLC	D	2	3	-	0/2/19/22	0/1/1/1
3	AGL	D	3	3,5	-	-	0/1/1/1
2	GLC	E	1	2,5	-	0/2/19/22	0/1/1/1
2	AGL	E	2	2,5	-	-	0/1/1/1
3	BGC	F	1	3	-	0/2/22/22	0/1/1/1
3	GLC	F	2	3	-	0/2/19/22	0/1/1/1
3	AGL	F	3	3,5	-	-	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	3	AGL	C3-C4-N4	4.62	120.51	111.05
3	F	3	AGL	C3-C4-N4	4.42	120.10	111.05
3	F	3	AGL	C3-C4-C5	-3.93	102.50	110.67
3	D	3	AGL	C3-C4-C5	-3.62	103.13	110.67
2	C	1	GLC	C1-O5-C5	3.32	116.69	112.19
3	D	3	AGL	C5-C4-N4	-2.99	104.73	110.92
3	F	3	AGL	C5-C4-N4	-2.80	105.11	110.92
2	E	2	AGL	C3-C4-C5	-2.43	105.61	110.67
3	D	2	GLC	C1-O5-C5	2.37	115.41	112.19
2	E	1	GLC	C1-O5-C5	2.37	115.40	112.19
2	E	1	GLC	C1-C2-C3	2.06	112.20	109.67
2	C	2	AGL	C3-C4-C5	-2.01	106.49	110.67

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	GLC	C4-C5-C6-O6
2	C	1	GLC	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	B	601	-	3,3,3	0.47	0	2,2,2	0.27	0
5	HMC	A	606	3,2	11,11,12	1.71	2 (18%)	9,15,17	0.69	0
5	HMC	B	602	2	11,11,12	0.65	0	9,15,17	0.78	0
4	EDO	A	601	-	3,3,3	0.54	0	2,2,2	0.05	0
5	HMC	B	605	3,2	11,11,12	1.31	2 (18%)	9,15,17	0.92	0
4	EDO	A	602	-	3,3,3	0.42	0	2,2,2	0.09	0
5	HMC	A	603	2	11,11,12	0.62	0	9,15,17	0.84	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	601	-	-	1/1/1/1	-
5	HMC	A	606	3,2	-	2/2/19/22	0/1/1/1
5	HMC	B	602	2	-	2/2/19/22	0/1/1/1
4	EDO	A	601	-	-	0/1/1/1	-
5	HMC	B	605	3,2	-	2/2/19/22	0/1/1/1
4	EDO	A	602	-	-	0/1/1/1	-
5	HMC	A	603	2	-	2/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	606	HMC	C4-C5	4.54	1.55	1.51
5	B	605	HMC	C4-C5	3.04	1.53	1.51
5	B	605	HMC	C1-C6	2.41	1.55	1.50
5	A	606	HMC	C1-C6	2.41	1.55	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	603	HMC	C1-C6-C5	-2.33	117.83	123.20

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	603	HMC	C4-C5-C7-O7
5	A	603	HMC	C6-C5-C7-O7
5	A	606	HMC	C6-C5-C7-O7
5	B	602	HMC	C4-C5-C7-O7
5	B	605	HMC	C4-C5-C7-O7
5	B	605	HMC	C6-C5-C7-O7
5	B	602	HMC	C6-C5-C7-O7
4	B	601	EDO	O1-C1-C2-O2
5	A	606	HMC	C4-C5-C7-O7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	518/564 (91%)	0.05	15 (2%) 51 55	26, 43, 65, 81	0
1	B	518/564 (91%)	-0.04	7 (1%) 75 78	25, 43, 63, 75	0
All	All	1036/1128 (91%)	0.01	22 (2%) 63 67	25, 43, 63, 81	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	544	GLY	4.0
1	A	67	SER	3.4
1	A	75	VAL	3.2
1	A	393	VAL	3.1
1	B	409	SER	3.0
1	A	552	SER	3.0
1	A	65	PHE	2.8
1	A	68	GLU	2.8
1	A	334	THR	2.5
1	B	385	PRO	2.4
1	A	576	LEU	2.4
1	A	197	GLY	2.4
1	B	576	LEU	2.4
1	A	409	SER	2.3
1	B	552	SER	2.3
1	B	59	ILE	2.2
1	A	212	PRO	2.2
1	A	64	ASP	2.2
1	B	64	ASP	2.1
1	A	385	PRO	2.1
1	A	207	ASP	2.1
1	A	123	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	AGL	D	3	10/11	0.89	0.20	31,33,36,41	0
3	AGL	F	3	10/11	0.89	0.22	37,38,40,45	0
3	GLC	D	2	11/12	0.92	0.13	34,37,39,40	0
3	BGC	D	1	12/12	0.93	0.14	42,45,52,53	0
2	GLC	C	1	11/12	0.93	0.12	42,46,48,49	0
2	GLC	E	1	11/12	0.94	0.12	40,42,43,43	0
2	AGL	E	2	10/11	0.94	0.13	47,53,57,65	0
3	BGC	F	1	12/12	0.94	0.11	40,44,47,49	0
2	AGL	C	2	10/11	0.94	0.17	48,50,56,61	0
3	GLC	F	2	11/12	0.95	0.11	38,39,40,43	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	HMC	A	603	11/12	0.75	0.33	69,72,74,77	0
5	HMC	B	602	11/12	0.81	0.33	74,77,80,82	0
5	HMC	A	606	11/12	0.83	0.21	43,45,47,48	0
4	EDO	B	601	4/4	0.87	0.19	56,57,57,57	0
4	EDO	A	602	4/4	0.88	0.15	63,64,64,65	0
5	HMC	B	605	11/12	0.89	0.19	45,47,50,51	0
4	EDO	A	601	4/4	0.93	0.13	60,62,63,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.