

wwPDB X-ray Structure Validation Summary Report (i)

Sep 26, 2023 – 10:51 PM EDT

PDB ID : 6CSB

Title: V308E mutant of cytochrome P450 2D6 complexed with thioridazine

Authors: Yang, Y.T.; Fujita, K.; Wang, P.F.; Im, S.C.; Pearl, N.M.; Meagher, J.;

Stuckey, J.; Waskell, L.

Deposited on : 2018-03-20

Resolution : 2.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	٨	470	4%		
1	A	479	85%	9%	• 6%
1	В	479	87%	8%	5%
1	С	479	8%	15%	5%
1	D	479	83%	11%	• 5%



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 15058 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cytochrome P450 2D6.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	Λ	452	Total	С	N	О	S	0	9	0
1	A	452	3527	2260	619	633	15	U	2	
1	В	455	Total	С	N	О	S	0	5	0
1	Ъ	400	3550	2273	631	631	15	U		
1	С	454	Total	С	N	О	S	0	3	0
1		404	3519	2255	617	632	15	U		
1	D	453	Total	С	N	О	S	0	3	0
1	ש	400	3501	2243	610	633	15	0	3	

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	initiating methionine	UNP P10635
A	24	ALA	-	expression tag	UNP P10635
A	25	LYS	-	expression tag	UNP P10635
A	26	LYS	-	expression tag	UNP P10635
A	27	THR	-	expression tag	UNP P10635
A	28	SER	-	expression tag	UNP P10635
A	29	SER	-	expression tag	UNP P10635
A	30	LYS	-	expression tag	UNP P10635
A	31	GLY	-	expression tag	UNP P10635
A	32	LYS	-	expression tag	UNP P10635
A	33	LEU	-	expression tag	UNP P10635
A	308	GLU	VAL	engineered mutation	UNP P10635
A	498	HIS	-	expression tag	UNP P10635
A	499	HIS	-	expression tag	UNP P10635
A	500	HIS	-	expression tag	UNP P10635
A	501	HIS	-	expression tag	UNP P10635
В	23	MET	-	initiating methionine	UNP P10635
В	24	ALA	-	expression tag	UNP P10635
В	25	LYS	-	expression tag	UNP P10635
В	26	LYS		expression tag	UNP P10635
В	27	THR	-	expression tag	UNP P10635



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	28	SER	-	expression tag	UNP P10635
В	29	SER	-	expression tag	UNP P10635
В	30	LYS	-	expression tag	UNP P10635
В	31	GLY	-	expression tag	UNP P10635
В	32	LYS	-	expression tag	UNP P10635
В	33	LEU	-	expression tag	UNP P10635
В	308	GLU	VAL	engineered mutation	UNP P10635
В	498	HIS	-	expression tag	UNP P10635
В	499	HIS	-	expression tag	UNP P10635
В	500	HIS	-	expression tag	UNP P10635
В	501	HIS	-	expression tag	UNP P10635
С	23	MET	-	initiating methionine	UNP P10635
С	24	ALA	-	expression tag	UNP P10635
С	25	LYS	-	expression tag	UNP P10635
С	26	LYS	-	expression tag	UNP P10635
С	27	THR	-	expression tag	UNP P10635
С	28	SER	-	expression tag	UNP P10635
С	29	SER	-	expression tag	UNP P10635
С	30	LYS	-	expression tag	UNP P10635
С	31	GLY	-	expression tag	UNP P10635
С	32	LYS	-	expression tag	UNP P10635
С	33	LEU	-	expression tag	UNP P10635
С	308	GLU	VAL	engineered mutation	UNP P10635
С	498	HIS	-	expression tag	UNP P10635
С	499	HIS	-	expression tag	UNP P10635
С	500	HIS	-	expression tag	UNP P10635
С	501	HIS	-	expression tag	UNP P10635
D	23	MET	-	initiating methionine	UNP P10635
D	24	ALA	-	expression tag	UNP P10635
D	25	LYS	-	expression tag	UNP P10635
D	26	LYS	-	expression tag	UNP P10635
D	27	THR	-	expression tag	UNP P10635
D	28	SER	-	expression tag	UNP P10635
D	29	SER	-	expression tag	UNP P10635
D	30	LYS	-	expression tag	UNP P10635
D	31	GLY	-	expression tag	UNP P10635
D	32	LYS	-	expression tag	UNP P10635
D	33	LEU	-	expression tag	UNP P10635
D	308	GLU	VAL	engineered mutation	UNP P10635
D	498	HIS	-	expression tag	UNP P10635
D	499	HIS	-	expression tag	UNP P10635
D	500	HIS	-	expression tag	UNP P10635



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	501	HIS	-	expression tag	UNP P10635


• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	Λ	1	Total	С	Fe	N	О	0	0	
	А	1	43	34	1	4	4	0		
2	В	1	Total	С	Fe	N	О	0	0	
	Б	1	43	34	1	4	4			
2	С	1	Total	С	Fe	N	О	0	0	
2	C	1	43	34	1	4	4	0	U	
2	D	1	Total	С	Fe	N	О	0	0	
	2 D	1	43	34	1	4	4			

 \bullet Molecule 3 is 10-{2-[(2R)-1-methylpiperidin-2-yl]ethyl}-2-(methylsulfanyl)-10H-phenothiazi ne (three-letter code: RTZ) (formula: $C_{21}H_{26}N_2S_2).$






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	С	N	S	0	0
3	А	1	25	21	2	2	U	0
3	A	1	Total	С	N	S	0	0
3	А	1	25	21	2	2	U	0
3	В	1	Total	С	N	S	0	0
3	Ъ	1	25	21	2	2	U	0
3	В	1	Total	С	N	S	0	0
3	D	1	25	21	2	2		
3	С	1	Total	С	N	S	0	0
3	C	1	25	21	2	2	0	0
3	С	1	Total	С	N	S	0	0
J	O	1	25	21	2	2	U	U
3	D	1	Total	С	N	S	0	0
		1	25	21	2	2		0
3	D	1	Total	С	N	S	0	0
3	ט	1	25	21	2	2		

 \bullet Molecule 4 is HEGA-10 (three-letter code: 2CV) (formula: C18H37NO7).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 10 6 1 3	0	0
4	В	1	Total C O 7 4 3	0	0
4	С	1	Total C O 7 4 3	0	0

 \bullet Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Zn 3 3	0	0
5	В	3	Total Zn 3 3	0	0
5	С	2	Total Zn 2 2	0	0
5	D	1	Total Zn 1 1	0	0


 \bullet Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	В	1	Total 4	C 2	O 2	0	0

 \bullet Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $\mathrm{C_3H_8O_3}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	С	1	Total C O 6 3 3	0	0

• Molecule 8 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	149	Total O 149 149	0	0
8	В	178	Total O 178 178	0	0
8	С	99	Total O 99 99	0	0
8	D	120	Total O 120 120	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cytochrome P450 2D6







• Molecule 1: Cytochrome P450 2D6





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	57.74Å 192.00Å 250.46Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.47 - 2.39	Depositor
resolution (A)	48.47 - 2.39	EDS
% Data completeness	95.2 (48.47-2.39)	Depositor
(in resolution range)	95.4 (48.47-2.39)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.25 (at 2.39Å)	Xtriage
Refinement program	PHENIX (1.12_2829)	Depositor
P.P.	0.193 , 0.229	Depositor
R, R_{free}	0.192 , 0.229	DCC
R_{free} test set	5234 reflections $(4.95%)$	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.460	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 54.0	EDS
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15058	wwPDB-VP
Average B, all atoms $(Å^2)$	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 20.65 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.2963e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, RTZ, ZN, 2CV, ACT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Во	ond angles
IVIOI	Cham	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	0.45	0/3626	0.59	3/4936 (0.1%)
1	В	0.44	0/3658	0.58	0/4977
1	С	0.41	0/3620	0.59	0/4927
1	D	0.43	1/3601 (0.0%)	0.60	2/4902 (0.0%)
All	All	0.43	1/14505~(0.0%)	0.59	5/19742 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
1	D	32	LYS	C-N	5.40	1.46	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	441	ARG	NE-CZ-NH2	-6.73	116.94	120.30
1	D	441	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	A	224	LEU	CB-CG-CD1	-5.84	101.07	111.00
1	D	115	ARG	NE-CZ-NH1	5.60	123.10	120.30
1	A	422	ASP	CB-CG-OD1	5.03	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3527	0	3469	33	0
1	В	3550	0	3488	28	0
1	С	3519	0	3456	53	0
1	D	3501	0	3406	49	0
2	A	43	0	30	0	0
2	В	43	0	30	3	0
2	С	43	0	30	4	0
2	D	43	0	30	4	0
3	A	50	0	52	1	0
3	В	50	0	52	4	0
3	С	50	0	52	5	0
3	D	50	0	52	8	0
4	A	10	0	8	3	0
4	В	7	0	6	1	0
4	С	7	0	6	4	0
5	A	3	0	0	0	0
5	В	3	0	0	0	0
5	С	2	0	0	0	0
5	D	1	0	0	0	0
6	В	4	0	3	0	0
7	С	6	0	8	0	0
8	A	149	0	0	1	0
8	В	178	0	0	4	1
8	С	99	0	0	1	0
8	D	120	0	0	3	0
All	All	15058	0	14178	173	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 173 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:D:220:LEU:HD22	1:D:240:VAL:CG1	1.71	1.18
1:D:217:SER:HB3	1:D:483:PHE:HE1	1.34	0.92
1:D:220:LEU:HD22	1:D:240:VAL:HG12	1.58	0.83
1:D:220:LEU:HD22	1:D:240:VAL:HG13	1.60	0.83
3:D:603:RTZ:HAH	3:D:603:RTZ:HAA	1.62	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ (\rm \AA) \end{array}$	Clash overlap (Å)
8:B:702:HOH:O	8:B:769:HOH:O[4_545]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	450/479~(94%)	434 (96%)	16 (4%)	0	100	100
1	В	456/479~(95%)	441 (97%)	15 (3%)	0	100	100
1	С	453/479~(95%)	434 (96%)	18 (4%)	1 (0%)	47	62
1	D	452/479~(94%)	434 (96%)	17 (4%)	1 (0%)	47	62
All	All	1811/1916 (94%)	1743 (96%)	66 (4%)	2 (0%)	51	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	483	PHE
1	С	240	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	374/409~(91%)	369 (99%)	5 (1%)	69	84
1	В	372/409 (91%)	366 (98%)	6 (2%)	62	79
1	С	371/409 (91%)	366 (99%)	5 (1%)	69	84



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	364/409 (89%)	358 (98%)	6 (2%)	62 79
All	All	1481/1636 (90%)	1459 (98%)	22 (2%)	71 80

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	450	ARG
1	D	146[B]	LYS
1	D	146[A]	LYS
1	D	213	LEU
1	В	146[A]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	210	GLN
1	С	350	GLN
1	D	424	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 9 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	gles
WIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	В	601	1,4	41,50,50	1.67	7 (17%)	45,82,82	1.47	6 (13%)
3	RTZ	С	603	-	27,28,28	0.20	0	35,39,39	0.76	0
2	HEM	A	601	1,4	41,50,50	1.61	8 (19%)	45,82,82	1.35	5 (11%)
2	HEM	D	601	1,8	41,50,50	1.55	7 (17%)	45,82,82	1.65	12 (26%)
3	RTZ	В	603	-	27,28,28	0.19	0	35,39,39	0.79	2 (5%)
3	RTZ	С	602	-	27,28,28	0.23	0	35,39,39	0.52	0
2	HEM	С	601	1,4	41,50,50	1.59	6 (14%)	45,82,82	1.39	5 (11%)
3	RTZ	В	602	-	27,28,28	0.24	0	35,39,39	0.49	0
3	RTZ	D	602	-	27,28,28	0.25	0	35,39,39	0.45	0
4	2CV	A	604	2	9,9,25	0.31	0	10,10,30	0.83	0
3	RTZ	D	603	-	27,28,28	0.18	0	35,39,39	1.47	1 (2%)
6	ACT	В	604	-	3,3,3	1.28	0	3,3,3	1.49	0
7	GOL	С	604	-	5,5,5	0.94	0	5,5,5	0.88	0
3	RTZ	A	602	-	27,28,28	0.22	0	35,39,39	0.48	0
4	2CV	С	605	2	6,6,25	0.42	0	7,7,30	0.61	0
4	2CV	В	605	2	6,6,25	0.41	0	7,7,30	1.05	1 (14%)
3	RTZ	A	603	-	27,28,28	0.23	0	35,39,39	0.97	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	В	601	1,4	-	4/12/54/54	-
3	RTZ	С	603	-	-	1/7/30/30	0/4/4/4
2	HEM	A	601	1,4	-	4/12/54/54	-
2	HEM	D	601	1,8	-	3/12/54/54	-
3	RTZ	В	603	-	-	3/7/30/30	0/4/4/4
3	RTZ	С	602	1	-	4/7/30/30	0/4/4/4
2	HEM	С	601	1,4	-	3/12/54/54	-
3	RTZ	В	602	-	-	4/7/30/30	0/4/4/4
3	RTZ	D	602	-	-	4/7/30/30	0/4/4/4
4	2CV	A	604	2	-	6/10/10/34	-
3	RTZ	D	603	-	-	5/7/30/30	0/4/4/4



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	С	604	-	-	2/4/4/4	-
3	RTZ	A	602	-	-	4/7/30/30	0/4/4/4
4	2CV	С	605	2	-	5/6/6/34	-
4	2CV	В	605	2	-	0/6/6/34	-
3	RTZ	A	603	-	-	2/7/30/30	0/4/4/4

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\operatorname{Observed}(\mathring{A})$	Ideal(Å)
2	С	601	HEM	C3C-CAC	4.44	1.56	1.47
2	A	601	HEM	C3C-CAC	4.40	1.56	1.47
2	D	601	HEM	C3C-CAC	4.28	1.56	1.47
2	В	601	HEM	C3C-CAC	4.18	1.56	1.47
2	В	601	HEM	C3C-C2C	-3.91	1.34	1.40

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	D	603	RTZ	CAL-CAP-CAG	-7.23	101.57	112.58
2	D	601	HEM	C4D-ND-C1D	3.96	109.17	105.07
2	В	601	HEM	C4C-CHD-C1D	3.77	127.53	122.56
2	В	601	HEM	C1B-NB-C4B	3.54	108.73	105.07
2	С	601	HEM	C1B-NB-C4B	3.34	108.52	105.07

There are no chirality outliers.

5 of 54 torsion outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$
3	A	602	RTZ	CAP-CAL-CAN-NAW
3	A	602	RTZ	CAN-CAL-CAP-CAG
3	A	602	RTZ	CAN-CAL-CAP-NAV
3	A	603	RTZ	CAN-CAL-CAP-CAG
3	A	603	RTZ	CAN-CAL-CAP-NAV

There are no ring outliers.

13 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	601	HEM	3	0
3	С	603	RTZ	3	0




Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	601	HEM	4	0
3	В	603	RTZ	3	0
3	С	602	RTZ	2	0
2	С	601	HEM	4	0
3	В	602	RTZ	1	0
3	D	602	RTZ	1	0
4	A	604	2CV	3	0
3	D	603	RTZ	7	0
4	С	605	2CV	4	0
4	В	605	2CV	1	0
3	A	603	RTZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.










































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	452/479 (94%)	0.23	18 (3%) 38 37	23, 43, 90, 111	0
1	В	455/479 (94%)	0.23	30 (6%) 18 17	23, 44, 85, 113	0
1	С	454/479 (94%)	0.53	36 (7%) 12 11	28, 51, 97, 135	0
1	D	453/479 (94%)	0.54	53 (11%) 4 4	28, 49, 98, 124	0
All	All	1814/1916 (94%)	0.38	137 (7%) 13 12	23, 46, 93, 135	0

The worst 5 of 137 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	240	VAL	8.6
1	С	33	LEU	6.6
1	С	38	LEU	5.9
1	С	342	VAL	5.7
1	В	240	VAL	5.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
5	ZN	С	607	1/1	0.43	0.10	139,139,139,139	0
3	RTZ	D	603	25/25	0.67	0.35	87,124,166,168	0
3	RTZ	С	603	25/25	0.73	0.28	73,118,134,151	0
3	RTZ	A	603	25/25	0.75	0.27	72,98,136,163	0
5	ZN	В	608	1/1	0.77	0.08	121,121,121,121	0
5	ZN	A	607	1/1	0.84	0.12	104,104,104,104	0
3	RTZ	В	603	25/25	0.84	0.19	67,89,117,127	0
3	RTZ	С	602	25/25	0.84	0.21	58,87,113,122	0
3	RTZ	D	602	25/25	0.85	0.17	54,74,92,95	0
4	2CV	A	604	10/26	0.88	0.30	39,75,98,105	0
4	2CV	В	605	7/26	0.89	0.31	37,93,115,125	0
7	GOL	С	604	6/6	0.89	0.18	55,75,78,82	0
3	RTZ	A	602	25/25	0.90	0.16	55,72,85,102	0
4	2CV	С	605	7/26	0.90	0.35	41,68,80,81	0
3	RTZ	В	602	25/25	0.91	0.14	44,59,70,75	0
5	ZN	В	607	1/1	0.92	0.17	110,110,110,110	0
6	ACT	В	604	4/4	0.93	0.17	60,68,74,98	0
2	HEM	В	601	43/43	0.98	0.16	18,25,30,34	0
2	HEM	С	601	43/43	0.98	0.18	23,31,36,44	0
2	HEM	D	601	43/43	0.98	0.17	18,28,32,34	0
2	HEM	A	601	43/43	0.98	0.16	18,26,29,32	0
5	ZN	A	606	1/1	0.99	0.11	46,46,46,46	0
5	ZN	A	605	1/1	0.99	0.14	38,38,38,38	0
5	ZN	С	606	1/1	0.99	0.10	36,36,36,36	0
5	ZN	В	606	1/1	1.00	0.12	38,38,38,38	0
5	ZN	D	604	1/1	1.00	0.12	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


































Electron density around 2CV A 604:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 2CV B 605:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









































6.5 Other polymers (i)

There are no such residues in this entry.

