



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 18, 2023 – 10:23 PM EDT

PDB ID : 5CVN
Title : WDR48 (2-580):USP46 ubiquitin ternary complex
Authors : Harris, S.F.; Yin, J.
Deposited on : 2015-07-27
Resolution : 3.36 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1558 (3.42-3.30)
Clashscore	141614	1627 (3.42-3.30)
Ramachandran outliers	138981	1599 (3.42-3.30)
Sidechain outliers	138945	1598 (3.42-3.30)
RSRZ outliers	127900	1507 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	598	 2% 66% 18% 14%
2	B	358	 % 70% 16% 10%
3	D	96	 % 59% 16% 21%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7323 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WD repeat-containing protein 48.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	514	4048	2559	710	757	22	0	0	0

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	MET	-	expression tag	UNP Q8TAF3
A	-13	HIS	-	expression tag	UNP Q8TAF3
A	-12	HIS	-	expression tag	UNP Q8TAF3
A	-11	HIS	-	expression tag	UNP Q8TAF3
A	-10	HIS	-	expression tag	UNP Q8TAF3
A	-9	HIS	-	expression tag	UNP Q8TAF3
A	-8	HIS	-	expression tag	UNP Q8TAF3
A	-7	GLY	-	expression tag	UNP Q8TAF3
A	-6	GLU	-	expression tag	UNP Q8TAF3
A	-5	ASN	-	expression tag	UNP Q8TAF3
A	-4	LEU	-	expression tag	UNP Q8TAF3
A	-3	TYR	-	expression tag	UNP Q8TAF3
A	-2	PHE	-	expression tag	UNP Q8TAF3
A	-1	GLN	-	expression tag	UNP Q8TAF3
A	0	GLY	-	expression tag	UNP Q8TAF3
A	1	SER	-	expression tag	UNP Q8TAF3
A	581	GLY	-	expression tag	UNP Q8TAF3
A	582	ASN	-	expression tag	UNP Q8TAF3
A	583	SER	-	expression tag	UNP Q8TAF3

- Molecule 2 is a protein called Ubiquitin carboxyl-terminal hydrolase 46.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	321	2640	1677	449	497	17	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	9	MET	-	expression tag	UNP P62068
B	10	HIS	-	expression tag	UNP P62068
B	11	HIS	-	expression tag	UNP P62068
B	12	HIS	-	expression tag	UNP P62068
B	13	HIS	-	expression tag	UNP P62068
B	14	HIS	-	expression tag	UNP P62068
B	15	HIS	-	expression tag	UNP P62068
B	16	GLY	-	expression tag	UNP P62068
B	17	GLU	-	expression tag	UNP P62068
B	18	ASN	-	expression tag	UNP P62068
B	19	LEU	-	expression tag	UNP P62068
B	20	TYR	-	expression tag	UNP P62068
B	21	PHE	-	expression tag	UNP P62068
B	22	GLN	-	expression tag	UNP P62068
B	23	GLY	-	expression tag	UNP P62068
B	24	SER	-	expression tag	UNP P62068

- Molecule 3 is a protein called Polyubiquitin-B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	76	600	378	105	116	1	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	MET	-	expression tag	UNP P0CG47
D	-18	GLY	-	expression tag	UNP P0CG47
D	-17	SER	-	expression tag	UNP P0CG47
D	-16	SER	-	expression tag	UNP P0CG47
D	-15	HIS	-	expression tag	UNP P0CG47
D	-14	HIS	-	expression tag	UNP P0CG47
D	-13	HIS	-	expression tag	UNP P0CG47
D	-12	HIS	-	expression tag	UNP P0CG47
D	-11	HIS	-	expression tag	UNP P0CG47
D	-10	HIS	-	expression tag	UNP P0CG47
D	-9	SER	-	expression tag	UNP P0CG47
D	-8	SER	-	expression tag	UNP P0CG47
D	-7	GLY	-	expression tag	UNP P0CG47
D	-6	LEU	-	expression tag	UNP P0CG47
D	-5	VAL	-	expression tag	UNP P0CG47

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-4	PRO	-	expression tag	UNP P0CG47
D	-3	ARG	-	expression tag	UNP P0CG47
D	-2	GLY	-	expression tag	UNP P0CG47
D	-1	SER	-	expression tag	UNP P0CG47
D	0	HIS	-	expression tag	UNP P0CG47

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Zn 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	15	Total O 15 15	0	0
5	B	13	Total O 13 13	0	0
5	D	6	Total O 6 6	0	0

Chain D:  59% 16% • 21%



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	103.44Å 154.19Å 182.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.36 – 3.36 47.36 – 3.36	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.36-3.36) 99.6 (47.36-3.36)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 3.40Å)	Xtrriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.175 , 0.222 0.182 , 0.233	Depositor DCC
R_{free} test set	1060 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	88.8	Xtrriage
Anisotropy	0.471	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 65.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7323	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/4132	0.71	1/5608 (0.0%)
2	B	0.47	0/2693	0.72	0/3631
3	D	0.42	0/606	0.72	0/815
All	All	0.45	0/7431	0.71	1/10054 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	LYS	C-N-CA	6.43	137.77	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4048	0	4050	48	0
2	B	2640	0	2590	25	0
3	D	600	0	629	10	0
4	B	1	0	0	0	0
5	A	15	0	0	1	0
5	B	13	0	0	1	0
5	D	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7323	0	7269	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:ASN:HA	1:A:55:ARG:HH11	1.44	0.82
3:D:8:LEU:HD11	3:D:71:LEU:HD22	1.65	0.79
2:B:340:ILE:HG13	2:B:341:GLU:H	1.47	0.78
1:A:370:ASN:HB2	1:A:412:LYS:HB2	1.70	0.72
1:A:192:LEU:HG	1:A:206:LEU:HB2	1.74	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/598 (85%)	472 (93%)	29 (6%)	7 (1%)	11	40
2	B	317/358 (88%)	288 (91%)	24 (8%)	5 (2%)	9	38
3	D	74/96 (77%)	71 (96%)	3 (4%)	0	100	100
All	All	899/1052 (86%)	831 (92%)	56 (6%)	12 (1%)	12	42

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	159	VAL
2	B	346	LEU
1	A	27	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	89	THR
2	B	354	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	450/521 (86%)	407 (90%)	43 (10%)	8	30
2	B	297/330 (90%)	256 (86%)	41 (14%)	3	15
3	D	68/85 (80%)	59 (87%)	9 (13%)	4	17
All	All	815/936 (87%)	722 (89%)	93 (11%)	5	22

5 of 93 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	202	VAL
2	B	273	ARG
2	B	208	ILE
2	B	236	GLN
2	B	317	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	68	HIS
2	B	239	GLN
2	B	119	GLN
1	A	381	ASN
2	B	146	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	514/598 (85%)	-0.05	14 (2%) 54 57	65, 107, 162, 208	0
2	B	321/358 (89%)	-0.16	4 (1%) 79 82	56, 85, 160, 205	0
3	D	76/96 (79%)	-0.33	0 100 100	58, 71, 89, 96	0
All	All	911/1052 (86%)	-0.11	18 (1%) 65 68	56, 97, 161, 208	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	SER	3.6
1	A	332	PHE	3.5
2	B	348	SER	3.4
1	A	157	ASN	3.2
1	A	560	PRO	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ZN	B	900	1/1	1.00	0.15	88,88,88,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.