

wwPDB X-ray Structure Validation Summary Report (i)

Aug 22, 2020 – 05:51 PM BST

PDB ID : 6CZ6

Title: Mycobacterium tuberculosis transcriptional regulator

Authors : Tang, S.; Sacchettini, J.

Deposited on : 2018-04-08

Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as 541 be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.13.1

 $buster\text{-report} \quad : \quad 1.1.7 \ (2018)$

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4: 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	429	2% 6 7%	10%	24%			
1	В	429	67%	9%	24%			
1	С	429	²⁹ 68%	8%	24%			
1	D	429	66%	10%	24%			



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 10643 atoms, of which 128 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called HTH-type transcriptional regulator PrpR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	Λ	328	Total	С	N	О	S	Se	0	0	0
1	A	320	2574	1625	471	465	6	7	0	U	0
1	В	328	Total	С	N	О	S	Se	0	0	0
1	Ъ	320	2574	1625	471	465	6	7	U	U	U
1	C	327	Total	С	N	О	S	Se	0	0	0
1		321	2563	1619	467	464	6	7	0	U	
1	D	326	Total	С	N	О	S	Se	0	0	0
1	D	320	2556	1614	466	463	6	7	U	U	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	58	HIS	-	expression tag	UNP O06581
A	59	HIS	-	expression tag	UNP O06581
A	60	HIS	-	expression tag	UNP O06581
A	61	HIS	_	expression tag	UNP O06581
A	62	HIS	_	expression tag	UNP O06581
A	63	HIS	_	expression tag	UNP O06581
A	64	SER	_	expression tag	UNP O06581
A	65	SER	-	expression tag	UNP O06581
A	66	GLY	_	expression tag	UNP O06581
A	67	VAL	_	expression tag	UNP O06581
A	68	ASP	_	expression tag	UNP O06581
A	69	LEU	_	expression tag	UNP O06581
A	70	GLY	_	expression tag	UNP O06581
A	71	THR	_	expression tag	UNP O06581
A	72	GLU	-	expression tag	UNP O06581
A	73	ASN	_	expression tag	UNP O06581
A	74	LEU	=	expression tag	UNP O06581
A	75	TYR	-	expression tag	UNP O06581
A	76	PHE	-	expression tag	UNP O06581
A	77	GLN	-	expression tag	UNP O06581
A	78	SER	-	expression tag	UNP O06581



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	ASN	-	expression tag	UNP O06581
A	80	ALA	-	expression tag	UNP O06581
A	337	ILE	VAL	conflict	UNP O06581
В	58	HIS	-	expression tag	UNP O06581
В	59	HIS	-	expression tag	UNP O06581
В	60	HIS	-	expression tag	UNP O06581
В	61	HIS	-	expression tag	UNP 006581
В	62	HIS	-	expression tag	UNP O06581
В	63	HIS	-	expression tag	UNP O06581
В	64	SER	-	expression tag	UNP O06581
В	65	SER	-	expression tag	UNP O06581
В	66	GLY	-	expression tag	UNP O06581
В	67	VAL	-	expression tag	UNP O06581
В	68	ASP	-	expression tag	UNP O06581
В	69	LEU	-	expression tag	UNP O06581
В	70	GLY	-	expression tag	UNP O06581
В	71	THR	_	expression tag	UNP O06581
В	72	GLU	-	expression tag	UNP O06581
В	73	ASN	-	expression tag	UNP O06581
В	74	LEU	_	expression tag	UNP O06581
В	75	TYR	-	expression tag	UNP O06581
В	76	PHE	-	expression tag	UNP O06581
В	77	GLN	-	expression tag	UNP O06581
В	78	SER	-	expression tag	UNP O06581
В	79	ASN	-	expression tag	UNP O06581
В	80	ALA	-	expression tag	UNP O06581
В	337	ILE	VAL	conflict	UNP O06581
С	58	HIS	-	expression tag	UNP O06581
С	59	HIS	-	expression tag	UNP O06581
С	60	HIS	-	expression tag	UNP O06581
С	61	HIS	-	expression tag	UNP O06581
С	62	HIS	-	expression tag	UNP O06581
С	63	HIS	-	expression tag	UNP O06581
С	64	SER	-	expression tag	UNP 006581
С	65	SER	-	expression tag	UNP O06581
С	66	GLY	-	expression tag	UNP O06581
С	67	VAL	-	expression tag	UNP O06581
C	68	ASP	-	expression tag	UNP 006581
С	69	LEU	-	expression tag	UNP 006581
C	70	GLY	-	expression tag	UNP 006581
С	71	THR	-	expression tag	UNP O06581
С	72	GLU	-	expression tag	UNP O06581




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	73	ASN	-	expression tag	UNP O06581
С	74	LEU	_	expression tag	UNP 006581
С	75	TYR	-	expression tag	UNP 006581
С	76	PHE	_	expression tag	UNP 006581
С	77	GLN	-	expression tag	UNP O06581
С	78	SER	-	expression tag	UNP O06581
С	79	ASN	-	expression tag	UNP O06581
С	80	ALA	-	expression tag	UNP O06581
С	337	ILE	VAL	conflict	UNP 006581
D	58	HIS	-	expression tag	UNP O06581
D	59	HIS	-	expression tag	UNP O06581
D	60	HIS	-	expression tag	UNP O06581
D	61	HIS	-	expression tag	UNP O06581
D	62	HIS	-	expression tag	UNP O06581
D	63	HIS	-	expression tag	UNP O06581
D	64	SER	-	expression tag	UNP O06581
D	65	SER	-	expression tag	UNP 006581
D	66	GLY	_	expression tag	UNP O06581
D	67	VAL	-	expression tag	UNP O06581
D	68	ASP	-	expression tag	UNP O06581
D	69	LEU	-	expression tag	UNP O06581
D	70	GLY	-	expression tag	UNP O06581
D	71	THR	-	expression tag	UNP O06581
D	72	GLU	-	expression tag	UNP O06581
D	73	ASN	-	expression tag	UNP O06581
D	74	LEU	-	expression tag	UNP O06581
D	75	TYR	-	expression tag	UNP O06581
D	76	PHE	-	expression tag	UNP O06581
D	77	GLN	-	expression tag	UNP O06581
D	78	SER	-	expression tag	UNP O06581
D	79	ASN	-	expression tag	UNP O06581
D	80	ALA	-	expression tag	UNP O06581
D	337	ILE	VAL	conflict	UNP O06581


 \bullet Molecule 2 is COENZYME A (three-letter code: COA) (formula: $\mathrm{C_{21}H_{36}N_7O_{16}P_3S}).$





Mol	Chain	Residues			Ato	ms				ZeroOcc	AltConf
2	Α	1	Total	С	Н	N	О	Р	S	0	0
	A	1	80	21	32	7	16	3	1	0	U
2	В	1	Total	С	Н	N	О	Р	S	0	0
	Б	1	80	21	32	7	16	3	1	0	U
2	С	1	Total	С	Н	N	О	Р	S	0	0
		1	80	21	32	7	16	3	1	0	U
9	D	1	Total	С	H	N	О	Р	S	0	0
	ש	1	80	21	32	7	16	3	1	U	0

 \bullet Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe $_4$ S4).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe S 8 4 4	0	0
3	В	1	Total Fe S	0	0
2		-	8 4 4 Total Fe S	0	0
3		1	8 4 4 Total Fe S	U	U
3	D	1	Total Fe S 8 4 4	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	7	Total O 7 7	0	0
4	В	6	Total O 6 6	0	0
4	С	4	Total O 4 4	0	0
4	D	7	Total O 7 7	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HTH-type transcriptional regulator PrpR









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	$96.19 { $	Denogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.68 - 2.70	Depositor
Resolution (A)	45.68 - 2.70	EDS
% Data completeness	99.4 (45.68-2.70)	Depositor
(in resolution range)	$99.4 \ (45.68 - 2.70)$	EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.95 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.13rc1_2954	Depositor
D D.	0.182 , 0.212	Depositor
R, R_{free}	0.182 , 0.212	DCC
R_{free} test set	2894 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	65.6	Xtriage
Anisotropy	0.417	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 36.9	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.176 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10643	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	$\mid \text{RMSZ} \mid \# Z > 5 \mid$		RMSZ	# Z > 5	
1	A	0.29	0/2628	0.50	0/3548	
1	В	0.28	0/2628	0.50	0/3548	
1	С	0.29	0/2616	0.50	0/3531	
1	D	0.29	0/2609	0.50	0/3521	
All	All	0.29	0/10481	0.50	0/14148	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2574	0	2523	30	0
1	В	2574	0	2523	33	1
1	С	2563	0	2509	37	1
1	D	2556	0	2500	38	0
2	A	48	32	32	0	0
2	В	48	32	32	0	0
2	С	48	32	32	0	0
2	D	48	32	32	0	0
3	A	8	0	0	0	0



$\alpha \cdots$	· ·	•	
Continued	trom	nromanne	naae
-	110111	picolous	payc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	8	0	0	0	0
3	С	8	0	0	0	0
3	D	8	0	0	0	0
4	A	7	0	0	0	0
4	В	6	0	0	1	0
4	С	4	0	0	0	0
4	D	7	0	0	0	0
All	All	10515	128	10183	122	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance } (ext{Å}) \end{array}$	$egin{array}{c} { m Clash} \ { m overlap} \ ({ m \AA}) \end{array}$
1:C:186:ARG:NH1	1:C:187:THR:O	1.92	1.02
1:C:186:ARG:NH1	1:C:188:GLY:HA3	1.77	0.99
1:D:445:ALA:HB3	1:D:449:ILE:HD11	1.55	0.88
1:C:186:ARG:HH11	1:C:188:GLY:HA3	1.39	0.86
1:B:376:GLU:HG3	1:D:376:GLU:HG3	1.57	0.84

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	Clash overlap (Å)
1:B:186:ARG:NH1	1:C:223:ASP:OD2[2_555]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$326/429 \ (76\%)$	320 (98%)	6 (2%)	0	100	100
1	В	$326/429 \ (76\%)$	317 (97%)	9 (3%)	0	100	100
1	С	$323/429 \ (75\%)$	315 (98%)	8 (2%)	0	100	100
1	D	$322/429 \ (75\%)$	315 (98%)	7 (2%)	0	100	100
All	All	1297/1716 (76%)	1267 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	$267/342 \ (78\%)$	263 (98%)	4 (2%)	65 86		
1	В	267/342~(78%)	265 (99%)	2 (1%)	84 94		
1	С	$266/342 \ (78\%)$	264 (99%)	2 (1%)	81 93		
1	D	$265/342 \ (78\%)$	265 (100%)	0	100 100		
All	All	1065/1368~(78%)	1057 (99%)	8 (1%)	81 93		

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	359	VAL
1	С	190	LEU
1	В	433	VAL
1	A	264	GLN
1	В	394	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Chain	Dog	Link	Bo	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	m Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	COA	С	501	_	41,50,50	0.96	1 (2%)	52,75,75	0.84	0
3	SF4	A	502	1	0,12,12	0.00	-	-		
2	COA	A	501	_	41,50,50	1.16	4 (9%)	52,75,75	0.83	0
2	COA	В	501	-	41,50,50	0.94	1 (2%)	52,75,75	0.89	2 (3%)
3	SF4	D	502	1	0,12,12	0.00	-	-		<u> </u>
3	SF4	В	502	1	0,12,12	0.00	-	-		
2	COA	D	501	_	41,50,50	1.14	4 (9%)	52,75,75	0.94	2 (3%)
3	SF4	С	502	1	0,12,12	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	С	501	-	-	5/44/64/64	0/3/3/3
3	SF4	A	502	1	-	-	0/6/5/5
2	COA	A	501	-	-	1/44/64/64	0/3/3/3
2	COA	В	501	-	-	5/44/64/64	0/3/3/3
3	SF4	D	502	1	=	-	0/6/5/5
3	SF4	В	502	1	-	-	0/6/5/5



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	D	501	-	_	16/44/64/64	0/3/3/3
3	SF4	С	502	1	-	-	0/6/5/5

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\mathring{\mathrm{A}})$	$Ideal(\AA)$
2	A	501	COA	P3B-O3B	3.08	1.65	1.59
2	A	501	COA	O4B-C1B	2.75	1.44	1.41
2	D	501	COA	P3B-O3B	2.50	1.64	1.59
2	A	501	COA	C3B-C4B	2.49	1.59	1.52
2	D	501	COA	OAP-CAP	2.47	1.46	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	В	501	COA	CDP-CBP-CAP	2.42	113.01	108.82
2	D	501	COA	CAP-C9P-N8P	2.08	120.72	116.58
2	В	501	COA	O5P-C5P-N4P	-2.05	119.14	123.01
2	D	501	COA	C7P-N8P-C9P	-2.03	118.97	122.59

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	501	COA	C5B-O5B-P1A-O1A
2	С	501	COA	C5B-O5B-P1A-O3A
2	С	501	COA	P1A-O3A-P2A-O6A
2	В	501	COA	C5B-O5B-P1A-O3A
2	В	501	COA	P1A-O3A-P2A-O6A


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	321/429 (74%)	-0.04	10 (3%) 49 49	41, 63, 93, 128	0
1	В	321/429 (74%)	0.01	10 (3%) 49 49	42, 64, 95, 139	0
1	С	320/429 (74%)	0.01	7 (2%) 62 63	43, 66, 97, 116	0
1	D	319/429 (74%)	-0.03	11 (3%) 45 45	43, 64, 94, 116	0
All	All	1281/1716 (74%)	-0.02	38 (2%) 50 51	41, 64, 95, 139	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	375	PRO	3.6
1	С	375	PRO	3.6
1	D	375	PRO	3.4
1	D	401	ASP	3.3
1	A	399	ALA	3.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	COA	A	501	48/48	0.94	0.18	59,75,98,100	0
2	COA	D	501	48/48	0.94	0.18	59,79,126,130	0
2	COA	В	501	48/48	0.96	0.17	58,73,110,128	0
2	COA	С	501	48/48	0.96	0.16	57,72,110,122	0
3	SF4	D	502	8/8	0.99	0.16	55,62,68,68	0
3	SF4	С	502	8/8	0.99	0.16	52,60,67,68	0
3	SF4	A	502	8/8	1.00	0.16	55,59,61,66	0
3	SF4	В	502	8/8	1.00	0.16	55,59,60,61	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

