



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 27, 2024 – 06:29 PM EST

PDB ID : 1D0O  
Title : BOVINE ENDOTHELIAL NITRIC OXIDE SYNTHASE HEME DOMAIN COMPLEXED WITH 3-BROMO-7-NITROINDAZOLE (H4B PRESENT)  
Authors : Raman, C.S.; Li, H.; Martasek, P.; Southan, G.J.; Masters, B.S.S.; Poulos, T.L.  
Deposited on : 1999-09-13  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	 7% 80% 13% • 6%
1	B	444	 8% 79% 13% • 7%

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7192 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOVINE ENDOTHELIAL NITRIC OXIDE SYNTHASE HEME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	416	3302	2099	584	603	16	0	0	0
1	B	414	3291	2092	582	601	16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	conflict	UNP P29473
B	100	ARG	CYS	conflict	UNP P29473

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C Fe N O 43 34 1 4 4	0	0
4	B	1	Total C Fe N O 43 34 1 4 4	0	0

- Molecule 5 is 3-BROMO-7-NITROINDAZOLE (three-letter code: INE) (formula:  $C_7H_4BrN_3O_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	Br	C	N	O		
5	A	1	Total	Br	C	N	O	0	0
			13	1	7	3	2		
5	A	1	Total	Br	C	N	O	0	0
			13	1	7	3	2		
5	B	1	Total	Br	C	N	O	0	0
			13	1	7	3	2		
5	B	1	Total	Br	C	N	O	0	0
			13	1	7	3	2		

- Molecule 6 is CACODYLIC ACID (three-letter code: CAD) (formula: C<sub>2</sub>H<sub>7</sub>AsO<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	As	C	0	0
			3	1	2		
6	B	1	Total	As	C	0	0
			3	1	2		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	234	Total	O	0	0
			234	234		
8	B	192	Total	O	0	0
			192	192		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BOVINE ENDOTHELIAL NITRIC OXIDE SYNTHASE HEME



- Molecule 1: BOVINE ENDOTHELIAL NITRIC OXIDE SYNTHASE HEME



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.59Å 106.06Å 155.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.95 31.19 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.95) 89.8 (31.19-1.95)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 1.95Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.200 , 0.247 0.200 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.3	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 54.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7192	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAD, INE, ACT, HEM, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3397	0.63	0/4631
1	B	0.37	0/3385	0.65	3/4614 (0.1%)
All	All	0.37	0/6782	0.64	3/9245 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	360	MET	N-CA-C	-5.60	95.87	111.00
1	B	99	ARG	NE-CZ-NH1	5.57	123.08	120.30
1	B	243	PHE	N-CA-C	-5.02	97.44	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3302	0	3215	36	0
1	B	3291	0	3205	38	0
2	A	8	0	6	0	0
2	B	8	0	6	0	0
3	A	1	0	0	0	0
4	A	43	0	30	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	43	0	30	0	0
5	A	26	0	8	0	0
5	B	26	0	8	0	0
6	A	3	0	0	0	0
6	B	3	0	0	0	0
7	A	6	0	8	0	0
7	B	6	0	8	1	0
8	A	234	0	0	3	0
8	B	192	0	0	2	0
All	All	7192	0	6524	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:LEU:HD11	1:A:156:GLU:HG2	1.69	0.75
1:B:126:LEU:HD11	1:B:156:GLU:HG2	1.69	0.74
1:B:114:ARG:HD3	1:B:115:PRO:HD2	1.68	0.74
1:B:114:ARG:HA	1:B:114:ARG:HH11	1.54	0.73
1:B:115:PRO:O	1:B:117:PRO:HD3	1.93	0.69
1:B:99:ARG:HB2	1:B:99:ARG:HH11	1.61	0.65
1:B:126:LEU:O	1:B:130:ARG:HG3	1.98	0.63
1:A:120:PRO:HB2	1:A:125:LEU:HB2	1.79	0.62
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.83	0.60
1:A:122:ALA:O	1:A:126:LEU:HB2	2.02	0.60
1:B:152:LEU:O	1:B:156:GLU:HG3	2.02	0.59
1:A:119:PRO:HG3	1:A:238:PRO:CG	2.33	0.58
1:A:378:LEU:HB2	8:A:1181:HOH:O	2.01	0.58
1:A:119:PRO:HG3	1:A:238:PRO:HG3	1.86	0.58
1:B:378:LEU:HB2	8:B:1052:HOH:O	2.03	0.58
1:B:322:LEU:HD13	1:B:324:TRP:CZ2	2.38	0.57
1:A:119:PRO:HG3	1:A:238:PRO:HB3	1.87	0.56
1:A:117:PRO:HG2	8:A:1087:HOH:O	2.05	0.56
1:A:340:ASN:H	1:A:340:ASN:HD22	1.53	0.55
1:A:455:SER:HB3	1:A:458:LEU:HD12	1.88	0.54
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.47	0.54
1:B:270:VAL:O	1:B:274:GLU:HG2	2.09	0.53
1:B:249:GLN:HB2	1:B:252:ARG:CG	2.39	0.53
1:B:236:ARG:HD3	1:B:351:SER:HB3	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:LEU:HD11	1:A:349:GLU:HB3	1.90	0.52
1:A:231:THR:O	1:A:353:ALA:HA	2.12	0.49
1:B:257:ARG:NH1	1:B:270:VAL:HG11	2.27	0.49
1:A:126:LEU:O	1:A:130:ARG:HG3	2.12	0.49
1:A:452:PRO:HG2	1:A:459:THR:HG21	1.94	0.49
1:A:119:PRO:HG3	1:A:238:PRO:CB	2.43	0.48
1:A:281:TRP:HB2	1:A:304:LEU:HD21	1.95	0.48
1:A:99:ARG:HD3	8:A:1136:HOH:O	2.13	0.48
1:A:249:GLN:HB2	1:A:252:ARG:HG2	1.96	0.48
1:A:180:TRP:CE3	1:A:192:TRP:HA	2.49	0.47
1:B:112:GLN:HG3	1:B:113:THR:H	1.77	0.47
1:B:249:GLN:HA	1:B:337:ALA:O	2.13	0.47
1:B:70:PHE:HA	1:B:71:PRO:HD3	1.74	0.47
1:A:263:VAL:HG11	1:A:267:PRO:HA	1.97	0.47
1:A:319:HIS:HB2	1:A:325:PHE:CB	2.45	0.47
1:A:109:ARG:O	1:A:110:LYS:HD2	2.15	0.47
1:A:213:ILE:O	1:A:217:ILE:HG13	2.15	0.46
1:B:169:GLU:O	1:B:173:VAL:HG23	2.14	0.46
1:A:322:LEU:HD13	1:A:324:TRP:CZ2	2.51	0.46
1:B:231:THR:O	1:B:353:ALA:HA	2.15	0.46
1:B:114:ARG:HD3	1:B:115:PRO:CD	2.41	0.46
1:B:409:LEU:O	1:B:413:GLN:HG3	2.16	0.45
1:B:455:SER:HB3	1:B:458:LEU:HD22	1.96	0.45
1:B:99:ARG:HH11	1:B:99:ARG:CB	2.27	0.45
7:B:881:GOL:H32	8:B:1007:HOH:O	2.15	0.45
1:B:381:VAL:O	1:B:385:MET:HG3	2.16	0.45
1:B:150:GLU:O	1:B:154:GLU:HG3	2.18	0.44
1:B:99:ARG:HH11	1:B:99:ARG:CG	2.30	0.44
1:B:472:SER:HA	1:B:473:PRO:C	2.38	0.44
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.48	0.43
1:A:390:ARG:HA	1:A:390:ARG:HD3	1.90	0.43
1:A:249:GLN:HA	1:A:337:ALA:O	2.18	0.42
1:A:381:VAL:O	1:A:385:MET:HG3	2.19	0.42
1:B:257:ARG:HH11	1:B:270:VAL:HG11	1.83	0.42
1:A:325:PHE:O	1:A:328:LEU:HB2	2.20	0.42
1:B:218:LYS:HE2	1:B:218:LYS:HB3	1.89	0.42
1:B:233:PHE:HB2	1:B:352:ALA:HB3	2.01	0.42
1:B:281:TRP:HB2	1:B:304:LEU:HD21	2.02	0.42
1:A:126:LEU:HD22	1:A:126:LEU:HA	1.84	0.41
1:B:236:ARG:HD2	1:B:242:ASP:OD2	2.21	0.41
1:A:379:GLU:O	1:A:383:VAL:HG23	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:398:ASP:O	1:A:402:VAL:HG23	2.21	0.41
1:B:140:ILE:HG13	1:B:142:ARG:HG2	2.03	0.41
1:A:472:SER:HA	1:A:473:PRO:C	2.41	0.41
1:B:366:THR:O	1:B:370:CYS:HB2	2.20	0.41
1:A:458:LEU:HD22	1:B:378:LEU:HD21	2.02	0.41
1:B:471:LEU:O	1:B:474:ALA:HB2	2.21	0.40
1:A:151:ARG:HD3	1:A:168:ARG:HE	1.86	0.40
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.62	0.40
1:B:294:LEU:HD23	1:B:294:LEU:HA	1.97	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	414/444 (93%)	394 (95%)	19 (5%)	1 (0%)	47	38
1	B	412/444 (93%)	391 (95%)	18 (4%)	3 (1%)	22	11
All	All	826/888 (93%)	785 (95%)	37 (4%)	4 (0%)	29	17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	259	GLN
1	B	108	PRO
1	A	115	PRO
1	B	115	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	354/377 (94%)	341 (96%)	13 (4%)	34	22
1	B	353/377 (94%)	342 (97%)	11 (3%)	40	28
All	All	707/754 (94%)	683 (97%)	24 (3%)	37	25

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	ARG
1	A	100	ARG
1	A	114	ARG
1	A	115	PRO
1	A	126	LEU
1	A	257	ARG
1	A	259	GLN
1	A	285	ASN
1	A	328	LEU
1	A	340	ASN
1	A	376	ASN
1	A	416	LYS
1	A	468	ASN
1	B	69	LYS
1	B	72	ARG
1	B	99	ARG
1	B	100	ARG
1	B	109	ARG
1	B	112	GLN
1	B	114	ARG
1	B	139	SER
1	B	259	GLN
1	B	260	ASP
1	B	340	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	GLN
1	A	153	GLN
1	A	191	GLN
1	A	278	GLN
1	A	285	ASN
1	A	340	ASN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	405	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	INE	B	765	-	12,14,14	3.29	3 (25%)	7,20,20	1.62	1 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	HEM	A	500	1	41,50,50	1.16	4 (9%)	45,82,82	1.72	9 (20%)
2	ACT	B	861	-	3,3,3	0.79	0	3,3,3	0.90	0
2	ACT	A	850	-	3,3,3	0.98	0	3,3,3	0.79	0
6	CAD	B	951	1	0,2,4	-	-	0,1,6	-	-
2	ACT	B	851	-	3,3,3	0.91	0	3,3,3	0.74	0
7	GOL	B	881	-	5,5,5	0.32	0	5,5,5	0.25	0
5	INE	A	766	-	12,14,14	3.30	3 (25%)	7,20,20	1.60	1 (14%)
6	CAD	A	950	1	0,2,4	-	-	0,1,6	-	-
2	ACT	A	860	-	3,3,3	0.90	0	3,3,3	0.77	0
7	GOL	A	880	-	5,5,5	0.26	0	5,5,5	0.29	0
5	INE	B	761	-	12,14,14	3.27	4 (33%)	7,20,20	1.57	1 (14%)
4	HEM	B	500	1	41,50,50	1.34	7 (17%)	45,82,82	1.41	6 (13%)
5	INE	A	760	-	12,14,14	3.72	4 (33%)	7,20,20	1.52	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	INE	B	765	-	-	0/2/4/4	0/2/2/2
4	HEM	A	500	1	-	4/12/54/54	-
7	GOL	B	881	-	-	1/4/4/4	-
5	INE	A	766	-	-	0/2/4/4	0/2/2/2
7	GOL	A	880	-	-	1/4/4/4	-
5	INE	B	761	-	-	0/2/4/4	0/2/2/2
4	HEM	B	500	1	-	3/12/54/54	-
5	INE	A	760	-	-	0/2/4/4	0/2/2/2

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	766	INE	C3-N2	8.81	1.46	1.34
5	B	765	INE	C3-N2	8.51	1.46	1.34
5	B	761	INE	C3-N2	8.17	1.45	1.34
5	A	760	INE	C3-N2	7.52	1.44	1.34
5	A	760	INE	BR-C3	-7.26	1.79	1.90
5	A	760	INE	C7-N10	-6.69	1.33	1.45
5	B	765	INE	C7-N10	-6.54	1.33	1.45
5	B	761	INE	C7-N10	-6.20	1.34	1.45
5	A	766	INE	C7-N10	-6.12	1.34	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	500	HEM	C3C-CAC	-3.32	1.41	1.47
5	B	761	INE	BR-C3	-3.23	1.85	1.90
4	A	500	HEM	C3C-CAC	-2.96	1.41	1.47
5	B	761	INE	C8-N1	-2.68	1.31	1.37
5	B	765	INE	C8-N1	-2.57	1.31	1.37
4	B	500	HEM	CAA-C2A	2.55	1.55	1.52
5	A	766	INE	C8-N1	-2.54	1.31	1.37
4	B	500	HEM	CAB-C3B	-2.47	1.40	1.47
5	A	760	INE	C8-N1	-2.46	1.31	1.37
4	B	500	HEM	CMD-C2D	2.36	1.55	1.50
4	B	500	HEM	CHA-C4D	2.35	1.41	1.35
4	B	500	HEM	C2C-C1C	2.25	1.47	1.42
4	B	500	HEM	C3C-C2C	-2.22	1.37	1.40
4	A	500	HEM	CAB-C3B	-2.22	1.41	1.47
4	A	500	HEM	CHA-C4D	2.07	1.40	1.35
4	A	500	HEM	C3C-C2C	-2.05	1.37	1.40

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	500	HEM	C2C-C3C-C4C	-5.38	103.14	106.90
4	B	500	HEM	C4B-CHC-C1C	4.27	128.19	122.56
4	A	500	HEM	CBA-CAA-C2A	4.04	119.52	112.62
5	B	765	INE	C6-C7-N10	4.01	120.76	116.47
5	A	766	INE	C6-C7-N10	3.87	120.61	116.47
5	B	761	INE	C6-C7-N10	3.73	120.46	116.47
4	A	500	HEM	C4B-CHC-C1C	3.52	127.21	122.56
5	A	760	INE	C6-C7-N10	3.49	120.20	116.47
4	A	500	HEM	CMD-C2D-C1D	3.47	130.32	125.04
4	B	500	HEM	C2C-C3C-C4C	-3.17	104.68	106.90
4	B	500	HEM	CBA-CAA-C2A	3.15	117.99	112.62
4	A	500	HEM	CAA-C2A-C3A	-2.72	119.43	127.25
4	A	500	HEM	CMC-C2C-C3C	2.44	129.25	124.68
4	B	500	HEM	CMD-C2D-C1D	2.35	128.62	125.04
4	B	500	HEM	C4C-CHD-C1D	2.33	125.64	122.56
4	A	500	HEM	C4D-ND-C1D	-2.26	102.74	105.07
4	B	500	HEM	CMA-C3A-C4A	-2.24	125.02	128.46
4	A	500	HEM	C4B-C3B-C2B	-2.13	105.43	107.11
4	A	500	HEM	C2D-C1D-ND	2.05	112.34	109.88

There are no chirality outliers.

All (9) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
4	B	500	HEM	C2B-C3B-CAB-CBB
7	A	880	GOL	O1-C1-C2-C3
4	A	500	HEM	CAD-CBD-CGD-O2D
4	B	500	HEM	CAA-CBA-CGA-O2A
4	A	500	HEM	CAD-CBD-CGD-O1D
4	B	500	HEM	CAA-CBA-CGA-O1A
4	A	500	HEM	CAA-CBA-CGA-O2A
7	B	881	GOL	O1-C1-C2-O2
4	A	500	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	881	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	416/444 (93%)	0.56	33 (7%) <b>12</b> <b>19</b>	19, 32, 68, 100	0
1	B	414/444 (93%)	0.61	37 (8%) <b>9</b> <b>15</b>	19, 34, 71, 100	0
All	All	830/888 (93%)	0.58	70 (8%) <b>11</b> <b>17</b>	19, 33, 69, 100	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	113	THR	13.4
1	B	115	PRO	11.5
1	A	113	THR	11.2
1	B	111	LEU	10.5
1	A	117	PRO	10.0
1	A	119	PRO	9.2
1	A	111	LEU	9.1
1	B	117	PRO	8.6
1	A	118	GLY	8.5
1	A	239	GLY	7.9
1	B	114	ARG	7.7
1	B	110	LYS	7.0
1	B	118	GLY	6.8
1	A	120	PRO	6.7
1	A	110	LYS	6.7
1	A	116	SER	6.4
1	A	115	PRO	6.4
1	A	114	ARG	6.1
1	B	112	GLN	5.9
1	B	119	PRO	5.6
1	B	259	GLN	5.2
1	A	259	GLN	5.0
1	B	142	ARG	4.9
1	B	120	PRO	4.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	123	GLU	4.2
1	B	147	ALA	4.1
1	A	121	PRO	4.0
1	B	121	PRO	3.9
1	B	116	SER	3.9
1	A	122	ALA	3.7
1	B	140	ILE	3.6
1	A	146	GLN	3.5
1	B	261	GLY	3.3
1	A	160	ALA	3.2
1	B	122	ALA	3.1
1	B	146	GLN	3.1
1	A	112	GLN	3.1
1	A	261	GLY	3.1
1	B	260	ASP	2.8
1	B	223	ARG	2.8
1	A	67	GLY	2.8
1	B	109	ARG	2.8
1	A	270	VAL	2.8
1	A	257	ARG	2.7
1	A	258	GLN	2.7
1	B	414	LEU	2.6
1	B	262	SER	2.6
1	B	141	LYS	2.6
1	A	260	ASP	2.6
1	B	390	ARG	2.6
1	B	124	GLN	2.5
1	A	414	LEU	2.5
1	B	123	GLU	2.4
1	A	69	LYS	2.4
1	B	389	THR	2.4
1	B	69	LYS	2.3
1	B	149	GLU	2.3
1	B	232	VAL	2.3
1	B	145	SER	2.3
1	A	124	GLN	2.2
1	A	194	LYS	2.2
1	B	453	PRO	2.1
1	A	99	ARG	2.1
1	B	258	GLN	2.1
1	A	449	TRP	2.1
1	A	250	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	353	ALA	2.1
1	A	126	LEU	2.1
1	A	150	GLU	2.0
1	B	150	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	GOL	A	880	6/6	0.85	0.22	51,53,55,57	0
7	GOL	B	881	6/6	0.91	0.20	50,52,53,56	0
2	ACT	A	850	4/4	0.92	0.25	41,43,45,46	0
2	ACT	B	861	4/4	0.92	0.11	34,37,38,39	0
5	INE	B	765	13/13	0.96	0.14	38,43,47,58	0
2	ACT	B	851	4/4	0.96	0.17	30,36,39,40	0
5	INE	A	766	13/13	0.96	0.11	33,36,46,56	0
4	HEM	B	500	43/43	0.97	0.17	21,29,41,47	0
6	CAD	A	950	3/5	0.98	0.08	52,52,53,64	0
6	CAD	B	951	3/5	0.98	0.10	68,68,69,72	0
4	HEM	A	500	43/43	0.98	0.13	17,22,41,42	0
2	ACT	A	860	4/4	0.98	0.11	35,37,37,40	0
5	INE	A	760	13/13	0.99	0.11	18,22,27,31	0
5	INE	B	761	13/13	0.99	0.16	31,34,38,40	0
3	ZN	A	900	1/1	0.99	0.07	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around HEM B 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.