



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 07:48 PM EDT

PDB ID : 2D3M
Title : Pentaketide chromone synthase complexed with coenzyme A
Authors : Morita, H.; Kondo, S.; Oguro, S.; Noguchi, H.; Sugio, S.; Abe, I.; Kohno, T.
Deposited on : 2005-09-29
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	
1	B	406	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called pentaketide chromone synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	405	3192	2031	540	594	27	0	8	0
1	B	395	3105	1981	526	572	26	0	4	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	GB 60686902
A	2	PRO	-	cloning artifact	GB 60686902
A	3	GLY	-	cloning artifact	GB 60686902
A	177	CSD	CYS	modified residue	GB 60686902
B	1	GLY	-	cloning artifact	GB 60686902
B	2	PRO	-	cloning artifact	GB 60686902
B	3	GLY	-	cloning artifact	GB 60686902
B	177	CSD	CYS	modified residue	GB 60686902

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	48	21	7	16	3	1	0	0
2	B	1	48	21	7	16	3	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	307	307	307	0	0
3	B	259	259	259	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: pentaketide chromone synthase



- Molecule 1: pentaketide chromone synthase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.15Å 88.40Å 70.04Å 90.00° 95.58° 90.00°	Depositor
Resolution (Å)	30.00 – 1.60 30.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.60) 99.8 (30.00-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.48 (at 1.60Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.198 , 0.207 0.195 , 0.204	Depositor DCC
R_{free} test set	5886 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	10.5	Xtrriage
Anisotropy	0.521	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 26.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.014 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6959	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/3255	0.71	0/4411
1	B	0.57	0/3167	0.70	0/4291
All	All	0.57	0/6422	0.71	0/8702

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	170	TYR	Sidechain
1	B	170	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3192	0	3194	12	0
1	B	3105	0	3114	12	0
2	A	48	0	31	0	0
2	B	48	0	31	0	0
3	A	307	0	0	0	0
3	B	259	0	0	0	0
All	All	6959	0	6370	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:143:CYS:SG	1:B:179:ALA:HB1	2.39	0.63
1:A:143:CYS:SG	1:A:179:ALA:HB1	2.44	0.57
1:B:18:GLN:HE21	1:B:22:LYS:HE2	1.74	0.52
1:A:205:GLU:HG3	1:A:354:VAL:HB	1.93	0.50
1:B:205:GLU:HG3	1:B:354:VAL:HB	1.93	0.50
1:A:263:THR:HG21	1:A:287:THR:HG21	1.93	0.49
1:A:386:LEU:HB3	1:A:398:ILE:HB	1.96	0.48
1:B:386:LEU:HB3	1:B:398:ILE:HB	1.95	0.48
1:A:150[B]:MET:HB2	1:B:269:LEU:HB2	1.96	0.46
1:A:72:ILE:O	1:A:76:THR:HG23	2.15	0.45
1:A:269:LEU:HB2	1:B:150[B]:MET:HB2	1.99	0.44
1:A:150[A]:MET:HB2	1:B:269:LEU:HB2	1.99	0.43
1:A:255:THR:HG22	1:A:398:ILE:HG23	2.01	0.42
1:B:320:PRO:HD3	1:B:361:ILE:HD11	2.02	0.42
1:A:269:LEU:HB2	1:B:150[A]:MET:HB2	2.00	0.42
1:B:186:TYR:O	1:B:190:LEU:HG	2.19	0.42
1:A:320:PRO:HD3	1:A:361:ILE:HD11	2.03	0.41
1:B:72:ILE:O	1:B:76:THR:HG23	2.20	0.41
1:A:172:ILE:HD12	1:A:183:VAL:HG12	2.01	0.41
1:B:172:ILE:HD12	1:B:183:VAL:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/406 (101%)	399 (97%)	9 (2%)	2 (0%)	25	10
1	B	396/406 (98%)	387 (98%)	8 (2%)	1 (0%)	37	20
All	All	806/812 (99%)	786 (98%)	17 (2%)	3 (0%)	30	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	354	VAL
1	B	354	VAL
1	A	15	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/339 (102%)	342 (99%)	4 (1%)	67	50
1	B	334/339 (98%)	331 (99%)	3 (1%)	75	62
All	All	680/678 (100%)	673 (99%)	7 (1%)	73	57

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	286	MET
1	A	299	VAL
1	A	366	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	405	LEU
1	B	64	GLU
1	B	286	MET
1	B	366	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	ASN
1	B	164	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	177	1	4,7,8	12.00	1 (25%)	1,8,10	0.81	0
1	CSD	B	177	1	4,7,8	12.13	1 (25%)	1,8,10	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	177	1	-	1/2/6/8	-
1	CSD	B	177	1	-	1/2/6/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	177	CSD	OD1-SG	24.22	1.69	1.47
1	A	177	CSD	OD1-SG	23.96	1.69	1.47

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	177	CSD	CA-CB-SG-OD1
1	B	177	CSD	CA-CB-SG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	COA	A	500	-	43,50,50	3.14	19 (44%)	56,75,75	2.86	22 (39%)
2	COA	B	501	-	43,50,50	3.15	20 (46%)	56,75,75	2.88	22 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	500	-	-	5/44/64/64	0/3/3/3
2	COA	B	501	-	-	6/44/64/64	0/3/3/3

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	COA	O5P-C5P	-10.26	1.02	1.23
2	A	500	COA	O5P-C5P	-10.19	1.03	1.23
2	A	500	COA	C4A-N3A	6.27	1.44	1.35
2	B	501	COA	C4A-N3A	6.09	1.43	1.35
2	B	501	COA	P1A-O5B	-6.04	1.35	1.59
2	A	500	COA	P1A-O5B	-5.96	1.36	1.59
2	A	500	COA	CCP-CBP	5.76	1.62	1.52
2	B	501	COA	CCP-CBP	5.72	1.61	1.52
2	B	501	COA	O4B-C1B	5.46	1.48	1.40
2	A	500	COA	C1B-N9A	5.23	1.62	1.49
2	A	500	COA	O4B-C1B	5.20	1.47	1.40
2	B	501	COA	C1B-N9A	5.19	1.62	1.49
2	A	500	COA	C2A-N1A	4.91	1.42	1.33
2	B	501	COA	C2A-N1A	4.77	1.42	1.33
2	A	500	COA	C7P-N8P	4.30	1.55	1.46
2	B	501	COA	C7P-N8P	4.28	1.55	1.46
2	B	501	COA	C3P-N4P	3.84	1.54	1.46
2	B	501	COA	C2A-N3A	3.81	1.38	1.32
2	A	500	COA	C2A-N3A	3.81	1.38	1.32
2	A	500	COA	C3P-N4P	3.75	1.54	1.46
2	B	501	COA	O4B-C4B	3.46	1.52	1.45
2	A	500	COA	O4B-C4B	3.40	1.52	1.45
2	A	500	COA	P3B-O7A	3.36	1.60	1.50
2	B	501	COA	P3B-O7A	3.33	1.60	1.50
2	B	501	COA	C5B-C4B	2.65	1.59	1.51
2	A	500	COA	C5B-C4B	2.62	1.59	1.51
2	A	500	COA	C3B-C4B	2.52	1.59	1.52
2	B	501	COA	C3B-C4B	2.42	1.59	1.52
2	A	500	COA	OAP-CAP	2.32	1.46	1.42
2	B	501	COA	O9P-C9P	2.31	1.27	1.23
2	B	501	COA	O2B-C2B	-2.29	1.37	1.43
2	A	500	COA	O3B-C3B	2.27	1.51	1.44
2	B	501	COA	OAP-CAP	2.25	1.46	1.42
2	A	500	COA	O2B-C2B	-2.25	1.37	1.43
2	A	500	COA	O9P-C9P	2.25	1.27	1.23
2	B	501	COA	O3B-C3B	2.23	1.51	1.44
2	A	500	COA	O5B-C5B	2.15	1.52	1.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	COA	O5B-C5B	2.14	1.52	1.44
2	B	501	COA	P1A-O3A	-2.00	1.57	1.59

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	COA	O5P-C5P-C6P	-8.99	105.73	122.02
2	A	500	COA	O5P-C5P-C6P	-8.74	106.18	122.02
2	B	501	COA	O4B-C1B-N9A	-7.04	99.41	108.75
2	A	500	COA	C2B-C3B-C4B	6.94	115.40	103.24
2	A	500	COA	O4B-C1B-N9A	-6.88	99.62	108.75
2	B	501	COA	C2B-C3B-C4B	6.72	115.01	103.24
2	A	500	COA	O2B-C2B-C3B	5.84	127.54	111.19
2	B	501	COA	O2B-C2B-C3B	5.84	127.54	111.19
2	B	501	COA	O5P-C5P-N4P	5.43	133.68	123.03
2	A	500	COA	O5P-C5P-N4P	5.33	133.49	123.03
2	A	500	COA	N3A-C2A-N1A	-4.74	122.24	128.67
2	B	501	COA	N3A-C2A-N1A	-4.72	122.27	128.67
2	A	500	COA	P3B-O3B-C3B	4.64	135.84	123.43
2	B	501	COA	P3B-O3B-C3B	4.58	135.67	123.43
2	A	500	COA	C6P-C7P-N8P	-3.96	103.57	112.00
2	B	501	COA	O4B-C4B-C3B	-3.90	96.69	104.92
2	B	501	COA	C6P-C7P-N8P	-3.84	103.83	112.00
2	A	500	COA	O4B-C4B-C3B	-3.83	96.84	104.92
2	A	500	COA	P1A-O5B-C5B	3.66	142.35	121.35
2	B	501	COA	P1A-O5B-C5B	3.64	142.22	121.35
2	B	501	COA	C7P-N8P-C9P	-3.63	116.02	122.55
2	A	500	COA	C7P-N8P-C9P	-3.49	116.28	122.55
2	A	500	COA	C3P-N4P-C5P	-3.39	116.51	122.82
2	B	501	COA	C3P-N4P-C5P	-3.22	116.82	122.82
2	A	500	COA	CDP-CBP-CAP	3.13	114.11	108.77
2	B	501	COA	CDP-CBP-CAP	3.00	113.88	108.77
2	B	501	COA	C4A-C5A-N7A	2.97	112.48	109.34
2	B	501	COA	C2P-C3P-N4P	2.97	119.05	112.31
2	A	500	COA	C4A-C5A-N7A	2.91	112.42	109.34
2	A	500	COA	O3B-P3B-O7A	-2.89	99.02	109.33
2	B	501	COA	O3B-P3B-O7A	-2.86	99.14	109.33
2	A	500	COA	C2P-C3P-N4P	2.68	118.39	112.31
2	B	501	COA	O9P-C9P-CAP	-2.36	114.30	120.89
2	A	500	COA	O9P-C9P-CAP	-2.33	114.39	120.89
2	B	501	COA	C6P-C5P-N4P	2.33	120.59	116.34
2	B	501	COA	C3B-C2B-C1B	-2.30	94.83	99.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	COA	C6P-C5P-N4P	2.19	120.33	116.34
2	A	500	COA	C3B-C2B-C1B	-2.15	95.15	99.89
2	A	500	COA	C5A-C6A-N1A	-2.12	115.17	120.23
2	A	500	COA	N6A-C6A-N1A	2.12	122.86	118.33
2	B	501	COA	N6A-C6A-N1A	2.11	122.84	118.33
2	B	501	COA	C5A-C6A-N1A	-2.10	115.23	120.23
2	A	500	COA	O3B-C3B-C4B	-2.09	102.65	110.03
2	B	501	COA	O3B-C3B-C4B	-2.06	102.78	110.03

There are no chirality outliers.


All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	COA	C2B-C3B-O3B-P3B
2	A	500	COA	S1P-C2P-C3P-N4P
2	B	501	COA	S1P-C2P-C3P-N4P
2	B	501	COA	C3B-C4B-C5B-O5B
2	B	501	COA	C2B-C3B-O3B-P3B
2	A	500	COA	C4B-C3B-O3B-P3B
2	B	501	COA	O4B-C4B-C5B-O5B
2	B	501	COA	P2A-O3A-P1A-O1A
2	A	500	COA	C3B-C4B-C5B-O5B
2	B	501	COA	P2A-O3A-P1A-O2A
2	A	500	COA	P2A-O3A-P1A-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/406 (99%)	0.27	28 (6%) 24 23	4, 11, 25, 34	8 (1%)
1	B	394/406 (97%)	0.27	32 (8%) 19 18	5, 10, 27, 37	4 (1%)
All	All	798/812 (98%)	0.27	60 (7%) 22 21	4, 10, 26, 37	12 (1%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	405	LEU	5.5
1	B	217	THR	5.4
1	B	13	LEU	5.1
1	B	12	PRO	5.0
1	A	217	THR	4.7
1	B	14	MET	4.5
1	B	218	HIS	4.4
1	B	405	LEU	4.0
1	B	11	LEU	4.0
1	A	218	HIS	3.9
1	B	64	GLU	3.8
1	B	99	TYR	3.8
1	B	219	LEU	3.7
1	B	268	HIS	3.7
1	A	216	GLU	3.6
1	B	63	VAL	3.6
1	B	280	LEU	3.6
1	A	268	HIS	3.5
1	A	219	LEU	3.5
1	A	67	LYS	3.5
1	A	282	LYS	3.3
1	B	282	LYS	3.3
1	B	216	GLU	3.3
1	B	265	ASP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	266	VAL	3.2
1	A	215	ASN	3.2
1	A	286	MET	3.2
1	A	265	ASP	3.1
1	B	279	TYR	3.1
1	B	100	ASP	3.1
1	A	243	ILE	3.0
1	A	68	LYS	2.9
1	A	280	LEU	2.9
1	B	215	ASN	2.9
1	B	89	PHE	2.8
1	B	243	ILE	2.8
1	A	41	PRO	2.8
1	B	223	ILE	2.7
1	B	67	LYS	2.7
1	A	281	SER	2.6
1	A	92	LYS	2.6
1	B	71	HIS	2.5
1	A	283	GLY	2.5
1	B	15	GLU	2.5
1	A	89	PHE	2.4
1	B	375	GLU	2.4
1	A	131	GLY	2.4
1	B	68	LYS	2.3
1	A	309	GLU	2.3
1	B	22	LYS	2.3
1	B	60	GLU	2.3
1	A	63	VAL	2.3
1	B	93	TYR	2.3
1	A	66	LYS	2.2
1	A	279	TYR	2.2
1	B	65	LEU	2.1
1	A	93	TYR	2.1
1	A	1	GLY	2.0
1	A	70	ASP	2.0
1	A	43	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	CSD	A	177	8/9	0.89	0.11	8,9,16,19	0
1	CSD	B	177	8/9	0.92	0.10	8,9,15,15	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	COA	A	500	48/48	0.74	0.16	33,40,47,48	0
2	COA	B	501	48/48	0.76	0.16	31,37,42,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.