



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2023 – 11:12 AM EDT

PDB ID : 3D4C  
Title : ZP-N domain of mammalian sperm receptor ZP3 (crystal form I)  
Authors : Jovine, L.; Monne, M.  
Deposited on : 2008-05-14  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	
2	B	2	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, LINKER, Zona pellucida protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	470	3629	2322	604	692	11	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P0AEX9
A	3	THR	ILE	engineered mutation	UNP P0AEX9
A	360	ALA	GLU	engineered mutation	UNP P0AEX9
A	363	ALA	LYS	engineered mutation	UNP P0AEX9
A	364	ALA	ASP	engineered mutation	UNP P0AEX9
A	368	ASN	ARG	engineered mutation	UNP P0AEX9
A	474	LEU	-	expression tag	UNP P10761
A	475	GLU	-	expression tag	UNP P10761
A	476	HIS	-	expression tag	UNP P10761
A	477	HIS	-	expression tag	UNP P10761
A	478	HIS	-	expression tag	UNP P10761
A	479	HIS	-	expression tag	UNP P10761
A	480	HIS	-	expression tag	UNP P10761
A	481	HIS	-	expression tag	UNP P10761

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0


- Molecule 3 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	Cd	0	0
			7	7		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein, LINKER, Zona pellucida protein 3



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.04Å 95.12Å 141.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	16.50 – 2.90 16.50 – 2.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (16.50-2.90) 100.0 (16.50-2.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.98 (at 2.92Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.190 , 0.227 0.194 , 0.231	Depositor DCC
$R_{free}$ test set	844 reflections (5.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	92.5	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 69.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.013 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3659	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	113.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/3711	0.43	0/5044

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3629	0	3609	72	0
2	B	23	0	21	0	0
3	A	7	0	0	0	0
All	All	3659	0	3630	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:ILE:CD1	1:A:436:LYS:HE2	1.58	1.32

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:ILE:HD11	1:A:436:LYS:HE2	1.12	1.04
1:A:213:ILE:HD11	1:A:436:LYS:CE	1.87	1.02
1:A:381:LEU:HD13	1:A:469:CYS:SG	2.25	0.76
1:A:213:ILE:HD12	1:A:436:LYS:HE2	1.68	0.76
1:A:48:PHE:N	1:A:49:PRO:HD2	2.00	0.75
1:A:69:GLY:HA3	1:A:333:ASN:O	1.87	0.74
1:A:120:LYS:HG3	1:A:124:PRO:HA	1.69	0.74
1:A:213:ILE:CD1	1:A:436:LYS:CE	2.51	0.70
1:A:11:TRP:CD2	1:A:58:PRO:HG3	2.31	0.66
1:A:152:LEU:HD11	1:A:205:MET:HE3	1.79	0.64
1:A:131:GLU:O	1:A:134:PRO:HD2	1.99	0.62
1:A:52:ALA:HA	1:A:56:ASP:O	2.01	0.61
1:A:213:ILE:HD11	1:A:436:LYS:CD	2.30	0.61
1:A:129:THR:OG1	1:A:132:GLU:HG3	2.01	0.60
1:A:168:TYR:CE1	1:A:183:GLY:HA3	2.37	0.59
1:A:120:LYS:CG	1:A:124:PRO:HA	2.32	0.59
1:A:423:ALA:HB1	1:A:428:CYS:SG	2.43	0.58
1:A:193:LEU:O	1:A:197:VAL:HG23	2.03	0.58
1:A:118:TYR:OH	1:A:120:LYS:CE	2.53	0.56
1:A:169:ALA:O	1:A:182:VAL:HG13	2.05	0.56
1:A:11:TRP:CG	1:A:58:PRO:HG3	2.39	0.56
1:A:90:LEU:HD12	1:A:95:TRP:CZ2	2.42	0.55
1:A:118:TYR:OH	1:A:120:LYS:HE2	2.07	0.54
1:A:62:PHE:CE2	1:A:265:ALA:HB2	2.43	0.54
1:A:431:ARG:HH12	1:A:433:GLN:CD	2.12	0.53
1:A:47:LYS:C	1:A:49:PRO:HD2	2.29	0.53
1:A:48:PHE:N	1:A:49:PRO:CD	2.71	0.53
1:A:116:LEU:HD22	1:A:249:PRO:HD3	1.90	0.52
1:A:7:LYS:HA	1:A:34:ILE:HG23	1.92	0.52
1:A:138:LYS:NZ	1:A:204:HIS:CE1	2.79	0.51
1:A:448:ASP:OD1	1:A:450:ARG:HG2	2.10	0.51
1:A:443:THR:CG2	1:A:467:ILE:HB	2.41	0.51
1:A:138:LYS:NZ	1:A:204:HIS:HE1	2.09	0.51
1:A:94:THR:HB	1:A:108:PRO:HB3	1.94	0.50
1:A:170:PHE:CE1	1:A:182:VAL:HG22	2.47	0.50
1:A:233:TRP:HB2	1:A:299:PRO:HG2	1.94	0.50
1:A:138:LYS:HZ2	1:A:204:HIS:CE1	2.30	0.50
1:A:48:PHE:CG	1:A:61:ILE:HD12	2.47	0.49
1:A:364:ALA:HB1	1:A:466:PRO:HD3	1.94	0.49
1:A:415:ASP:HB2	1:A:418:VAL:HG22	1.95	0.48
1:A:443:THR:HG22	1:A:467:ILE:HB	1.95	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:HIS:N	1:A:41:PRO:HD3	2.29	0.48
1:A:159:TRP:N	1:A:160:PRO:CD	2.77	0.47
1:A:259:PHE:CD1	1:A:331:MET:HG2	2.50	0.47
1:A:260:VAL:HB	1:A:330:ILE:HA	1.96	0.47
1:A:19:ASN:O	1:A:23:GLU:HG3	2.15	0.47
1:A:349:ILE:HG21	1:A:470:ARG:HH21	1.78	0.46
1:A:256:SER:O	1:A:258:PRO:HD3	2.14	0.46
1:A:349:ILE:HG21	1:A:470:ARG:NH2	2.31	0.46
1:A:248:LEU:HD12	1:A:258:PRO:HG3	1.97	0.45
1:A:123:LEU:HD21	1:A:127:PRO:HD3	1.99	0.45
1:A:450:ARG:HA	1:A:451:PRO:C	2.37	0.44
1:A:368:ASN:HA	1:A:372:VAL:HB	1.99	0.44
1:A:259:PHE:CG	1:A:331:MET:HG2	2.53	0.43
1:A:308:TYR:O	1:A:312:LEU:HD13	2.18	0.43
1:A:90:LEU:HB2	1:A:95:TRP:NE1	2.34	0.43
1:A:168:TYR:CD1	1:A:183:GLY:HA3	2.54	0.43
1:A:184:VAL:O	1:A:362:LEU:HB3	2.19	0.43
1:A:247:VAL:HG21	1:A:327:LYS:NZ	2.34	0.42
1:A:46:GLU:O	1:A:49:PRO:HG2	2.19	0.42
1:A:381:LEU:CD1	1:A:469:CYS:SG	3.03	0.42
1:A:315:ASP:HA	1:A:316:PRO:HD2	1.80	0.42
1:A:425:LEU:HD12	1:A:425:LEU:HA	1.83	0.42
1:A:280:PHE:O	1:A:284:TYR:HB2	2.20	0.42
1:A:450:ARG:HE	1:A:450:ARG:HB3	1.69	0.41
1:A:248:LEU:CD1	1:A:258:PRO:HG3	2.50	0.41
1:A:49:PRO:HA	1:A:76:LEU:HD13	2.02	0.41
1:A:81:THR:N	1:A:82:PRO:HD3	2.35	0.41
1:A:234:SER:HB3	1:A:298:LYS:HD3	2.02	0.41
1:A:130:TRP:CE2	1:A:161:LEU:HD13	2.57	0.40
1:A:452:VAL:HG21	1:A:456:SER:OG	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/481 (97%)	458 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/393 (97%)	381 (100%)	1 (0%)	92	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	437	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	B	1	2	12,12,12	0.59	0	17,17,17	0.51	0
2	GLC	B	2	2	11,11,12	0.75	0	15,15,17	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	470/481 (97%)	0.05	30 (6%) <b>19</b> <b>15</b>	51, 111, 169, 210	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	454	GLY	6.5
1	A	456	SER	4.9
1	A	451	PRO	4.8
1	A	242	ASN	4.5
1	A	174	ASN	4.2
1	A	173	GLU	4.0
1	A	453	SER	3.9
1	A	474	LEU	3.7
1	A	392	THR	3.4
1	A	165	ASP	3.3
1	A	176	LYS	3.1
1	A	452	VAL	2.7
1	A	225	MET	2.7
1	A	296	LYS	2.6
1	A	455	LEU	2.5
1	A	289	GLU	2.5
1	A	147	ALA	2.5
1	A	7	LYS	2.5
1	A	240	LYS	2.4
1	A	450	ARG	2.4
1	A	407	GLY	2.4
1	A	142	ALA	2.4
1	A	459	ARG	2.4
1	A	145	LYS	2.3
1	A	436	LYS	2.3
1	A	125	ASN	2.3
1	A	323	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	118	TYR	2.1
1	A	314	LYS	2.0
1	A	30	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	B	1	12/12	0.97	0.14	82,89,99,100	0
2	GLC	B	2	11/12	0.98	0.14	62,67,78,82	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CD	A	484	1/1	0.83	0.05	160,160,160,160	0
3	CD	A	483	1/1	0.92	0.09	139,139,139,139	1
3	CD	A	486	1/1	0.94	0.04	176,176,176,176	0
3	CD	A	489	1/1	0.94	0.04	166,166,166,166	1
3	CD	A	487	1/1	0.97	0.05	153,153,153,153	0
3	CD	A	488	1/1	0.97	0.06	149,149,149,149	0
3	CD	A	485	1/1	0.97	0.39	153,153,153,153	0



## 6.5 Other polymers [i](#)

There are no such residues in this entry.