



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 28, 2023 – 02:15 PM JST

PDB ID : 5D4I  
Title : Intact nitrite complex of a copper nitrite reductase determined by serial femtosecond crystallography  
Authors : Fukuda, Y.; Tse, K.M.; Nakane, T.; Nakatsu, T.; Suzuki, M.; Sugahara, M.; Inoue, S.; Masuda, T.; Yumoto, F.; Matsugaki, N.; Nango, E.; Tono, K.; Joti, Y.; Kameshima, T.; Song, C.; Hatsui, T.; Yabashi, M.; Nureki, O.; Murphy, M.E.P.; Inoue, T.; Iwata, S.; Mizohata, E.  
Deposited on : 2015-08-07  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	 6% 88% 9% .
1	B	342	 9% 94% . .
1	C	342	 4% 90% 7% . .

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8260 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper-containing nitrite reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	2612	1668	444	488	12	0	7	0
1	B	341	2666	1703	456	496	11	0	7	0
1	C	334	2637	1685	446	494	12	0	10	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MET	-	initiating methionine	UNP P38501
A	341	LEU	-	expression tag	UNP P38501
A	342	VAL	-	expression tag	UNP P38501
A	343	PRO	-	expression tag	UNP P38501
A	344	ARG	-	expression tag	UNP P38501
B	3	MET	-	initiating methionine	UNP P38501
B	341	LEU	-	expression tag	UNP P38501
B	342	VAL	-	expression tag	UNP P38501
B	343	PRO	-	expression tag	UNP P38501
B	344	ARG	-	expression tag	UNP P38501
C	3	MET	-	initiating methionine	UNP P38501
C	341	LEU	-	expression tag	UNP P38501
C	342	VAL	-	expression tag	UNP P38501
C	343	PRO	-	expression tag	UNP P38501
C	344	ARG	-	expression tag	UNP P38501

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	0
			2	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Cu 2 2	0	0
2	C	2	Total Cu 2 2	0	0

- Molecule 3 is NITRITE ION (three-letter code: NO2) (formula: NO<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total N O 3 1 2	0	0
3	A	1	Total N O 3 1 2	0	0
3	C	1	Total N O 3 1 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	93	Total O 95 95	0	4
4	B	113	Total O 114 114	0	2
4	C	117	Total O 121 121	0	5


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Copper-containing nitrite reductase



- Molecule 1: Copper-containing nitrite reductase



- Molecule 1: Copper-containing nitrite reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.00Å 103.00Å 147.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.97 – 1.60 28.97 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.97-1.60) 100.0 (28.97-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.165 , 0.196 0.164 , 0.195	Depositor DCC
$R_{free}$ test set	6263 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.7	Xtrriage
Anisotropy	0.044	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	8260	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NO2, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.05	1/2683 (0.0%)	1.05	11/3654 (0.3%)
1	B	1.06	0/2738	1.05	3/3730 (0.1%)
1	C	1.10	1/2708 (0.0%)	1.11	5/3688 (0.1%)
All	All	1.07	2/8129 (0.0%)	1.07	19/11072 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	3
1	C	0	1
All	All	0	7

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	246	SER	CA-CB	5.36	1.60	1.52
1	C	313	GLU	CD-OE2	-5.34	1.19	1.25

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	250	ARG	NE-CZ-NH2	-9.96	115.32	120.30
1	B	253	ARG	NE-CZ-NH1	8.47	124.53	120.30
1	A	253	ARG	NE-CZ-NH2	7.10	123.85	120.30
1	A	231[A]	LYS	C-N-CA	6.99	139.18	121.70
1	A	231[B]	LYS	C-N-CA	6.99	139.18	121.70

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	250	ARG	NE-CZ-NH1	6.24	123.42	120.30
1	A	253	ARG	NE-CZ-NH1	-6.07	117.27	120.30
1	A	250	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	C	98	ASP	CB-CG-OD1	5.85	123.56	118.30
1	C	188	ASP	CB-CG-OD1	5.79	123.51	118.30
1	A	230	ASP	O-C-N	-5.66	113.65	122.70
1	B	256	LEU	CB-CG-CD2	-5.62	101.45	111.00
1	B	329	ASP	CB-CG-OD1	5.62	123.35	118.30
1	A	251	ASP	CB-CG-OD1	5.47	123.23	118.30
1	A	211	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	C	162	LEU	CB-CG-CD1	-5.26	102.06	111.00
1	A	54	ASP	CB-CG-OD1	5.16	122.94	118.30
1	A	53	ASP	CB-CG-OD1	5.10	122.89	118.30
1	A	54	ASP	CB-CG-OD2	-5.08	113.73	118.30

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	230	ASP	Mainchain
1	A	306	HIS	Peptide
1	B	306	HIS	Peptide
1	B	58	GLU	Mainchain
1	C	306	HIS	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2612	0	2532	9	0
1	B	2666	0	2601	7	0
1	C	2637	0	2557	10	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	6	0	0	0	0
3	C	3	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	95	0	0	0	0
4	B	114	0	0	0	0
4	C	121	0	0	2	0
All	All	8260	0	7690	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231[A]:LYS:HD2	1:A:231[A]:LYS:N	2.05	0.71
1:A:6:ALA:HB2	1:C:329:ASP:HB2	1.83	0.59
1:C:257:ILE:HD12	1:C:302:ALA:HB3	1.88	0.56
1:B:257:ILE:HD12	1:B:302:ALA:HB3	1.88	0.55
1:A:7:ALA:HA	1:A:8:GLU:HG2	1.88	0.55
1:A:187[B]:ARG:HD3	1:A:193:TYR:CE1	2.43	0.54
1:B:344:ARG:HD3	1:C:116:PRO:HG2	1.90	0.53
1:B:207:VAL:O	1:B:211[B]:ARG:HG3	2.10	0.50
1:A:69:PRO:HG3	1:A:179:GLY:HA3	1.95	0.48
1:A:132:PHE:CE2	1:A:269:LYS:HE3	2.48	0.48
1:C:50:ILE:O	1:C:58[A]:GLU:HA	2.17	0.45
1:C:88:PRO:HD2	1:C:91:ASN:ND2	2.32	0.45
1:A:257:ILE:HD12	1:A:302:ALA:HB3	1.98	0.44
1:A:235:ALA:O	1:A:322:VAL:HA	2.18	0.44
1:A:255:HIS:ND1	1:A:279:GLU:O	2.51	0.43
1:B:16[A]:LYS:HD2	1:B:17:VAL:N	2.34	0.43
1:C:235:ALA:O	1:C:322:VAL:HA	2.19	0.42
1:C:14[B]:ARG:HG3	1:C:38:VAL:HB	2.01	0.41
1:B:342:VAL:HB	1:B:343:PRO:HD2	2.03	0.41
1:C:37:LYS:HE2	4:C:622:HOH:O	2.20	0.41
1:C:267:THR:HG22	4:C:713:HOH:O	2.21	0.41
1:C:305:ASN:O	1:C:311:ALA:HB2	2.22	0.40
1:B:253:ARG:HA	1:B:281:TRP:O	2.21	0.40
1:B:58:GLU:CD	1:B:195:LYS:HE3	2.41	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/342 (99%)	333 (98%)	6 (2%)	0	100	100
1	B	346/342 (101%)	344 (99%)	2 (1%)	0	100	100
1	C	342/342 (100%)	338 (99%)	4 (1%)	0	100	100
All	All	1027/1026 (100%)	1015 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/270 (100%)	261 (97%)	9 (3%)	38	14
1	B	276/270 (102%)	269 (98%)	7 (2%)	47	22
1	C	273/270 (101%)	266 (97%)	7 (3%)	46	21
All	All	819/810 (101%)	796 (97%)	23 (3%)	49	18

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LYS
1	A	29	SER
1	A	33	GLU
1	A	89	GLU
1	A	160	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	192	LYS
1	A	231[A]	LYS
1	A	231[B]	LYS
1	A	325	GLU
1	B	16[A]	LYS
1	B	16[B]	LYS
1	B	159[A]	ARG
1	B	159[B]	ARG
1	B	168	LYS
1	B	341	LEU
1	B	344	ARG
1	C	37	LYS
1	C	58[A]	GLU
1	C	58[B]	GLU
1	C	166	LYS
1	C	168	LYS
1	C	197	GLU
1	C	253	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NO2	C	503	2	1,2,2	0.12	0	0,1,1	-	-
3	NO2	A	504	2	1,2,2	0.49	0	0,1,1	-	-
3	NO2	A	503	2	1,2,2	0.32	0	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	334/342 (97%)	0.11	22 (6%) 18 17	25, 37, 63, 83	0
1	B	341/342 (99%)	0.22	32 (9%) 8 7	23, 34, 63, 103	0
1	C	334/342 (97%)	-0.05	14 (4%) 36 33	22, 31, 54, 88	0
All	All	1009/1026 (98%)	0.09	68 (6%) 17 16	22, 34, 62, 103	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	ALA	9.6
1	B	7	ALA	8.5
1	C	7	ALA	5.8
1	C	165	GLY	5.6
1	B	5	THR	5.4
1	A	167	GLY	5.4
1	A	339	GLY	5.1
1	B	4	ALA	4.7
1	A	6	ALA	4.5
1	B	6	ALA	4.5
1	B	340	THR	4.3
1	C	339	GLY	3.9
1	B	33	GLU	3.9
1	A	166	LYS	3.8
1	A	190	ASN	3.8
1	B	154	ILE	3.7
1	B	167	GLY	3.4
1	B	81	LEU	3.4
1	A	189	GLU	3.3
1	B	99	PHE	3.3
1	A	325	GLU	3.2
1	B	169	ALA	3.2
1	C	189	GLU	3.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	229	GLY	3.1
1	B	190	ASN	3.1
1	B	189	GLU	3.0
1	B	133	VAL	3.0
1	C	190	ASN	2.9
1	B	132	PHE	2.8
1	C	166	LYS	2.8
1	B	230	ASP	2.8
1	C	168	LYS	2.8
1	B	197	GLU	2.7
1	B	10	ALA	2.7
1	B	97	ILE	2.7
1	B	11	ALA	2.6
1	B	341	LEU	2.6
1	A	163	HIS	2.6
1	A	338	SER	2.6
1	A	33	GLU	2.5
1	B	339	GLY	2.5
1	B	8	GLU	2.5
1	C	11	ALA	2.4
1	A	191	GLY	2.4
1	B	134	TYR	2.4
1	B	163	HIS	2.4
1	B	73	MET	2.3
1	B	34	GLY	2.3
1	A	34	GLY	2.3
1	A	11	ALA	2.3
1	B	72	LEU	2.3
1	B	83	LEU	2.3
1	B	168	LYS	2.2
1	A	337	PRO	2.2
1	A	168	LYS	2.2
1	A	165	GLY	2.2
1	C	326	TRP	2.2
1	A	12	LEU	2.1
1	A	15	GLN	2.1
1	B	342	VAL	2.1
1	A	18	GLU	2.1
1	A	192	LYS	2.1
1	C	9	ILE	2.1
1	C	192	LYS	2.0
1	C	338	SER	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	323	THR	2.0
1	C	133	VAL	2.0
1	B	124	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NO2	C	503	3/3	0.89	0.10	31,31,37,42	0
3	NO2	A	504	3/3	0.95	0.08	37,37,44,47	0
3	NO2	A	503	3/3	0.96	0.07	36,36,39,43	0
2	CU	B	502	1/1	1.00	0.05	32,32,32,32	0
2	CU	C	501	1/1	1.00	0.05	26,26,26,26	0
2	CU	C	502	1/1	1.00	0.04	25,25,25,25	0
2	CU	A	501	1/1	1.00	0.06	35,35,35,35	0
2	CU	A	502	1/1	1.00	0.05	31,31,31,31	0
2	CU	B	501	1/1	1.00	0.05	35,35,35,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.