



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 29, 2024 – 12:12 PM EDT

PDB ID : 9D5J
Title : Human Adenosine Deaminase Acting on dsRNA (ADAR2-RD) bound to dsRNA containing deoxyinosine at the -1 position of the guide strand
Authors : Fisher, A.J.; Cheng, J.; Manjunath, A.; Campbell, K.
Deposited on : 2024-08-13
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)
RNA backbone	3690	1037 (3.00-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	 48% 29% 21% 2%
1	B	487	 58% 30% 5% 7% 2%
2	C	32	 31% 56% 12%
3	D	32	 59% 31% 9%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 8019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform 4 of Double-stranded RNA-specific editase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	384	3019	1904	550	554	11	0	0	0
1	B	454	3540	2235	639	653	13	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	488	GLN	GLU	engineered mutation	UNP P78563
B	488	GLN	GLU	engineered mutation	UNP P78563


- Molecule 2 is a RNA chain called RNA Top Strand Containing 8-azanebularine (8AZ).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	32	685	305	126	223	31	0	0	0

- Molecule 3 is DNA/RNA hybrid called RNA Bottom Strand containing deoxyinosine complementary to a guanosine adjacent to the target site.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	32	674	303	121	219	31	0	0	0

- Molecule 4 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula: $C_6H_{18}O_{24}P_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O P 36 6 24 6	0	0
4	B	1	Total C O P 36 6 24 6	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Zn 1 1	0	0
5	B	1	Total Zn 1 1	0	0


- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	9	Total O 9 9	0	0
7	B	2	Total O 2 2	0	0
7	C	3	Total O 3 3	0	0
7	D	1	Total O 1 1	0	0



- Molecule 2: RNA Top Strand Containing 8-azanebularine (8AZ)



- Molecule 3: RNA Bottom Strand containing deoxyinosine complementary to a guanosine adjacent to the target site



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	174.69Å 63.13Å 141.78Å 90.00° 118.23° 90.00°	Depositor
Resolution (Å)	49.60 – 2.80 49.60 – 2.81	Depositor EDS
% Data completeness (in resolution range)	65.0 (49.60-2.80) 64.9 (49.60-2.81)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, R_{free}	0.188 , 0.235 0.190 , 0.237	Depositor DCC
R_{free} test set	1675 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	69.8	Xtrriage
Anisotropy	0.101	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 61.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8019	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 8AZ, EDO, IHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/3081	0.71	0/4164
1	B	0.41	0/3610	0.64	0/4876
2	C	0.70	0/740	1.28	4/1151 (0.3%)
3	D	0.65	0/727	1.25	2/1128 (0.2%)
All	All	0.51	0/8158	0.83	6/11319 (0.1%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	18	G	N3-C4-C5	7.71	132.46	128.60
2	C	18	G	N3-C4-N9	-6.39	122.17	126.00
2	C	16	G	N3-C4-N9	-6.06	122.36	126.00
3	D	10	C	C6-N1-C2	-5.83	117.97	120.30
2	C	18	G	C2-N3-C4	-5.67	109.07	111.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3019	0	3051	84	1
1	B	3540	0	3567	98	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	685	0	337	17	1
3	D	674	0	349	7	0
4	A	36	0	6	4	0
4	B	36	0	6	3	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	4	0	6	0	0
6	C	4	0	6	2	0
6	D	4	0	6	0	0
7	A	9	0	0	0	0
7	B	2	0	0	0	0
7	C	3	0	0	0	0
7	D	1	0	0	0	0
All	All	8019	0	7334	198	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 198 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:569:ASP:O	1:B:674:ARG:NH1	2.12	0.81
1:A:410:GLN:HG2	1:A:425:ILE:HG12	1.62	0.80
1:A:514:MET:HE1	1:A:688:VAL:H	1.47	0.79
1:A:461:GLU:HG3	1:A:462:PRO:HD2	1.64	0.76
1:B:558:ARG:HA	1:B:562:GLN:HB2	1.69	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:410:GLN:NE2	2:C:32:G:O2'[4_445]	2.17	0.03

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/487 (78%)	348 (91%)	32 (8%)	2 (0%)	25	56
1	B	450/487 (92%)	403 (90%)	45 (10%)	2 (0%)	30	61
All	All	832/974 (85%)	751 (90%)	77 (9%)	4 (0%)	25	56

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	322	VAL
1	B	497	ALA
1	B	318	HIS
1	A	467	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/415 (79%)	292 (89%)	37 (11%)	5	16
1	B	384/415 (92%)	326 (85%)	58 (15%)	2	8
All	All	713/830 (86%)	618 (87%)	95 (13%)	3	10

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	386	ARG
1	B	513	THR
1	B	390	LEU
1	B	448	THR
1	B	566	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	410	GLN
1	B	496	ASN
1	B	694	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	30/32 (93%)	5 (16%)	0
3	D	30/32 (93%)	7 (23%)	0
All	All	60/64 (93%)	12 (20%)	0

5 of 12 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2	C
2	C	6	C
2	C	14	G
2	C	26	A
2	C	29	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	8AZ	C	13	5,2	16,24,25	2.74	4 (25%)	12,35,38	1.22	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8AZ	C	13	5,2	-	2/3/35/36	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	13	8AZ	C2-N1	8.23	1.48	1.34
2	C	13	8AZ	N7-N8	4.74	1.42	1.34
2	C	13	8AZ	N8-N9	3.78	1.41	1.34
2	C	13	8AZ	C2-N3	2.94	1.34	1.29

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	13	8AZ	C1'-N9-N8	-2.28	113.46	118.20

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	13	8AZ	O4'-C4'-C5'-O5'
2	C	13	8AZ	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	13	8AZ	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	A	803	-	3,3,3	0.43	0	2,2,2	0.18	0
6	EDO	C	101	-	3,3,3	0.32	0	2,2,2	0.33	0
4	IHP	B	801	-	36,36,36	0.95	1 (2%)	60,60,60	0.74	1 (1%)
6	EDO	D	101	-	3,3,3	0.32	0	2,2,2	0.08	0
4	IHP	A	801	-	36,36,36	0.97	3 (8%)	60,60,60	0.89	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	803	-	-	0/1/1/1	-
6	EDO	C	101	-	-	1/1/1/1	-
4	IHP	B	801	-	-	7/30/54/54	0/1/1/1
6	EDO	D	101	-	-	0/1/1/1	-
4	IHP	A	801	-	-	4/30/54/54	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	801	IHP	P5-O15	2.76	1.64	1.59
4	B	801	IHP	P4-O14	2.73	1.64	1.59
4	A	801	IHP	P3-O13	2.56	1.64	1.59
4	A	801	IHP	P1-O11	2.53	1.64	1.59

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	801	IHP	P4-O14-C4	2.54	130.22	123.43
4	A	801	IHP	O16-P6-O26	-2.09	101.88	109.33

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	801	IHP	C4-C5-O15-P5
6	C	101	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	801	IHP	C6-C5-O15-P5
4	B	801	IHP	C1-O11-P1-O41
4	B	801	IHP	C3-O13-P3-O33


There are no ring outliers.


3 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	101	EDO	2	0
4	B	801	IHP	3	0
4	A	801	IHP	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	384/487 (78%)	-0.55	0 100 100	28, 47, 92, 136	0
1	B	454/487 (93%)	-0.15	10 (2%) 62 53	45, 76, 152, 204	0
2	C	31/32 (96%)	-0.56	0 100 100	42, 78, 134, 139	0
3	D	31/32 (96%)	-0.60	0 100 100	38, 81, 127, 133	0
All	All	900/1038 (86%)	-0.35	10 (1%) 77 71	28, 62, 141, 204	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	HIS	3.4
1	B	499	ILE	2.6
1	B	415	LEU	2.4
1	B	501	THR	2.4
1	B	700	THR	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	8AZ	C	13	22/23	0.96	0.07	35,38,42,45	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	D	101	4/4	0.83	0.12	47,52,58,61	0
6	EDO	C	101	4/4	0.84	0.15	66,70,77,82	0
6	EDO	A	803	4/4	0.89	0.10	46,49,54,58	0
4	IHP	B	801	36/36	0.95	0.06	47,70,82,84	0
4	IHP	A	801	36/36	0.98	0.05	28,36,42,47	0
5	ZN	B	802	1/1	0.99	0.02	55,55,55,55	0
5	ZN	A	802	1/1	0.99	0.05	42,42,42,42	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around IHP B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IHP A 801:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.