



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 18, 2024 – 06:10 PM EST

PDB ID : 9D8F
Title : Crystal structure of the ACVR1 (ALK2) Kinase Domain in complex with inhibitor CDD-2281
Authors : Ta, H.M.; Kim, C.; Jimmidi, R.; Matzuk, M.M.
Deposited on : 2024-08-19
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.21
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	 5% 85% 11% 5%
1	B	317	 5% 85% 9% 5%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10374 atoms, of which 5036 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Activin receptor type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	307	Total 4979	C 1580	H 2490	N 438	O 456	S 15	0	5	0
1	B	300	Total 4892	C 1550	H 2448	N 432	O 445	S 17	0	8	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 5	O 4	P 1	0	0
2	A	1	Total 5	O 4	P 1	0	0
2	A	1	Total 5	O 4	P 1	0	0
2	A	1	Total 5	O 4	P 1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	1
			28	6	16	6		
3	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is N-[3-(dimethylamino)propyl]-1-[(1r,3r)-3-(methylcarbamoyl)cyclobutyl]-2-(3,4,5-trimethoxyphenyl)-1H-1,3-benzimidazole-6-carboxamide (three-letter code: A1A3C) (formula: C₂₈H₃₇N₅O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
4	A	1	75	28	37	5	5	0	0
4	B	1	75	28	37	5	5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	148	149	149	0	1
5	B	109	112	112	0	3


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Activin receptor type-1



- Molecule 1: Activin receptor type-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.66Å 71.04Å 158.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.85 – 1.86 43.85 – 1.86	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.85-1.86) 100.0 (43.85-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 1.86Å)	Xtrriage
Refinement program	PHENIX (1.21rc1_5049: ???)	Depositor
R, R_{free}	0.186 , 0.223 0.186 , 0.223	Depositor DCC
R_{free} test set	48834 reflections (3.93%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtrriage
Anisotropy	0.245	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10374	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.92 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.4976e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, A1A3C, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2555	0.55	0/3464
1	B	0.29	0/2525	0.54	0/3421
All	All	0.30	0/5080	0.55	0/6885

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2489	2490	2482	23	1
1	B	2444	2448	2425	20	1
2	A	30	0	0	2	0
2	B	20	0	0	0	0
3	A	12	16	16	0	0
3	B	6	8	8	0	0
4	A	38	37	0	0	0
4	B	38	37	0	0	0
5	A	149	0	0	6	0
5	B	112	0	0	0	0
All	All	5338	5036	4931	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:GLU:OE2	5:A:701:HOH:O	1.66	1.10
2:A:602:PO4:O2	5:A:701:HOH:O	1.87	0.90
1:A:428:LYS:NZ	5:A:703:HOH:O	2.07	0.85
1:A:212:GLU:OE2	5:A:702:HOH:O	2.03	0.76
1:A:373:ASN:OD1	1:A:375:ARG:N	2.19	0.74
1:A:492:LYS:O	1:A:496:THR:HG23	1.93	0.69
1:B:204:VAL:HG12	1:B:207:GLN:H	1.57	0.69
1:A:240:ARG:NH2	5:A:705:HOH:O	2.26	0.68
1:A:250:GLU:O	1:A:254:THR:HG22	1.95	0.66
1:B:434:VAL:HG23	1:B:435:VAL:HG13	1.79	0.62
1:B:204:VAL:HG22	1:B:206:ARG:NH2	2.18	0.59
1:A:211:LEU:HD11	1:A:226[A]:SER:OG	2.06	0.56
1:B:472:LYS:O	1:B:476:GLU:HG3	2.08	0.54
1:A:200:VAL:O	1:A:204:VAL:HG23	2.09	0.53
1:B:204:VAL:HG11	1:B:207:GLN:OE1	2.08	0.53
1:A:407:PHE:CE2	1:A:491:ILE:HG21	2.44	0.53
1:B:251:LEU:C	1:B:251:LEU:HD23	2.32	0.50
1:A:197:PRO:O	1:A:198:PHE:CB	2.61	0.48
1:B:489:LEU:HD11	1:B:493:LYS:HZ3	1.79	0.48
1:B:306:LEU:HD13	1:B:499:ASP:HA	1.96	0.47
1:A:373:ASN:OD1	1:A:375:ARG:CA	2.61	0.47
1:A:497:LYS:HD2	1:A:497:LYS:O	2.14	0.47
1:A:251:LEU:C	1:A:251:LEU:HD23	2.35	0.47
1:B:287:GLU:OE1	1:B:346:LYS:HE3	2.15	0.47
1:A:196:LEU:CD1	1:A:204:VAL:HG21	2.45	0.46
1:A:260[B]:GLU:OE2	5:A:704:HOH:O	2.21	0.46
1:B:251:LEU:HD21	1:B:257:LEU:HD22	1.98	0.46
1:B:204:VAL:HG11	1:B:207:GLN:CG	2.46	0.45
1:B:287:GLU:OE1	1:B:346:LYS:CE	2.65	0.44
1:A:335:ARG:HD3	1:A:357:LEU:O	2.17	0.44
1:B:271:THR:HG21	1:B:273[B]:ARG:NE	2.33	0.44
1:A:392:GLN:HG2	1:A:395:CYS:CB	2.48	0.44
1:B:498:ILE:O	1:B:499:ASP:C	2.57	0.43
1:B:288[A]:MET:CE	1:B:344:VAL:HG12	2.50	0.42
1:B:323:ILE:HD12	1:B:329:LYS:HG2	2.01	0.42
1:A:407:PHE:CE2	1:A:411:LEU:HD11	2.53	0.42
1:A:338:LYS:HD2	2:A:601:PO4:O4	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:271:THR:HG21	1:B:273[B]:ARG:HE	1.83	0.42
1:A:287:GLU:OE1	1:A:346:LYS:HE3	2.21	0.41
1:A:433:ASP:OD2	1:A:433:ASP:N	2.44	0.41
1:B:407:PHE:CE2	1:B:491:ILE:HG21	2.56	0.41
1:B:457:ILE:HG23	1:B:461:TRP:CE3	2.56	0.40
1:B:211:LEU:HD11	1:B:226:SER:HB2	2.04	0.40
1:A:194:SER:OG	1:A:257:LEU:N	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:367:GLN:OE1	1:B:392:GLN:NE2[3_554]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/317 (98%)	301 (97%)	6 (2%)	3 (1%)	13	4
1	B	306/317 (96%)	297 (97%)	8 (3%)	1 (0%)	37	25
All	All	616/634 (97%)	598 (97%)	14 (2%)	4 (1%)	19	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	202	ARG
1	A	198	PHE
1	A	367	GLN
1	A	370	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/284 (98%)	274 (99%)	3 (1%)	70	62
1	B	272/284 (96%)	272 (100%)	0	100	100
All	All	549/568 (97%)	546 (100%)	3 (0%)	84	84

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256	MET
1	A	366	ASN
1	A	397	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	392	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	601	-	4,4,4	1.57	1 (25%)	6,6,6	0.45	0
2	PO4	A	603	-	4,4,4	1.39	1 (25%)	6,6,6	0.53	0
2	PO4	B	602	-	4,4,4	1.40	1 (25%)	6,6,6	0.51	0
3	GOL	A	607[A]	-	5,5,5	0.34	0	5,5,5	0.34	0
3	GOL	B	605	-	5,5,5	0.39	0	5,5,5	0.64	0
2	PO4	A	602	-	4,4,4	1.62	1 (25%)	6,6,6	0.84	0
2	PO4	A	606	-	4,4,4	1.35	1 (25%)	6,6,6	0.55	0
4	A1A3C	A	608	-	38,41,41	2.84	16 (42%)	48,58,58	2.86	19 (39%)
2	PO4	A	605	-	4,4,4	1.60	1 (25%)	6,6,6	0.54	0
2	PO4	B	601	-	4,4,4	1.47	1 (25%)	6,6,6	0.55	0
2	PO4	B	603	-	4,4,4	1.28	1 (25%)	6,6,6	0.64	0
3	GOL	A	607[B]	-	5,5,5	0.31	0	5,5,5	0.34	0
4	A1A3C	B	606	-	38,41,41	2.65	11 (28%)	48,58,58	2.00	12 (25%)
2	PO4	B	604	-	4,4,4	1.50	1 (25%)	6,6,6	0.51	0
2	PO4	A	604	-	4,4,4	1.60	1 (25%)	6,6,6	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	607[A]	-	-	0/4/4/4	-
3	GOL	B	605	-	-	2/4/4/4	-
4	A1A3C	A	608	-	-	12/27/39/39	0/4/4/4
3	GOL	A	607[B]	-	-	2/4/4/4	-
4	A1A3C	B	606	-	-	3/27/39/39	0/4/4/4

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	606	A1A3C	C19-N20	9.26	1.46	1.33
4	A	608	A1A3C	C19-N20	9.10	1.46	1.33
4	A	608	A1A3C	O34-C33	6.14	1.47	1.37
4	B	606	A1A3C	C02-N03	5.70	1.46	1.33
4	B	606	A1A3C	O28-C27	5.33	1.45	1.37
4	A	608	A1A3C	C02-N03	5.29	1.45	1.33
4	A	608	A1A3C	C12-C11	4.78	1.47	1.36
4	B	606	A1A3C	O34-C33	4.75	1.44	1.37
4	A	608	A1A3C	O28-C27	4.66	1.44	1.37
4	A	608	A1A3C	O31-C30	4.51	1.46	1.38
4	B	606	A1A3C	O31-C30	3.93	1.45	1.38
4	B	606	A1A3C	C12-C11	3.85	1.45	1.36
4	A	608	A1A3C	C10-C02	3.46	1.57	1.50
4	B	606	A1A3C	C38-C14	3.28	1.47	1.40
4	A	608	A1A3C	C38-C14	3.11	1.46	1.40
4	A	608	A1A3C	C36-C25	3.04	1.45	1.39
2	A	602	PO4	P-O1	2.98	1.57	1.50
2	A	604	PO4	P-O1	2.79	1.57	1.50
2	A	605	PO4	P-O1	2.73	1.57	1.50
2	A	601	PO4	P-O1	2.71	1.56	1.50
2	B	604	PO4	P-O1	2.60	1.56	1.50
2	B	601	PO4	P-O1	2.52	1.56	1.50
4	B	606	A1A3C	C38-C10	2.51	1.42	1.37
2	A	603	PO4	P-O1	2.38	1.56	1.50
2	B	602	PO4	P-O1	2.32	1.56	1.50
4	B	606	A1A3C	C27-C30	-2.28	1.36	1.41
4	A	608	A1A3C	C08-N07	2.24	1.53	1.46
2	A	606	PO4	P-O1	2.23	1.55	1.50
4	A	608	A1A3C	C38-C10	2.21	1.41	1.37
4	B	606	A1A3C	C33-C30	-2.13	1.36	1.41
4	A	608	A1A3C	C25-C24	2.11	1.53	1.47
4	B	606	A1A3C	O22-C19	2.09	1.27	1.23
4	A	608	A1A3C	C13-N37	2.09	1.45	1.38
2	B	603	PO4	P-O1	2.07	1.55	1.50
4	A	608	A1A3C	O22-C19	2.04	1.27	1.23
4	A	608	A1A3C	C05-C04	2.04	1.59	1.51
4	A	608	A1A3C	C33-C30	-2.04	1.36	1.41

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	608	A1A3C	C18-C19-N20	9.79	128.56	116.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	608	A1A3C	C10-C02-N03	8.11	133.98	117.12
4	A	608	A1A3C	O01-C02-N03	-6.47	109.99	122.59
4	A	608	A1A3C	O22-C19-N20	-6.21	112.78	123.13
4	B	606	A1A3C	C18-C19-N20	5.99	123.76	116.18
4	B	606	A1A3C	C11-C12-C13	-4.77	115.10	120.80
4	A	608	A1A3C	O28-C27-C26	-4.12	116.98	124.08
4	A	608	A1A3C	C11-C12-C13	-4.10	115.90	120.80
4	B	606	A1A3C	O22-C19-N20	-3.89	116.65	123.13
4	A	608	A1A3C	C04-N03-C02	3.81	130.72	122.11
4	B	606	A1A3C	O28-C27-C26	-3.67	117.75	124.08
4	B	606	A1A3C	C10-C02-N03	3.53	124.46	117.12
4	A	608	A1A3C	O28-C27-C30	3.37	120.91	115.14
4	B	606	A1A3C	O01-C02-N03	-3.30	116.16	122.59
4	B	606	A1A3C	O28-C27-C30	3.25	120.71	115.14
4	B	606	A1A3C	O34-C33-C36	-3.23	118.52	124.08
4	A	608	A1A3C	N37-C24-N15	-2.94	107.69	115.06
4	A	608	A1A3C	C38-C10-C02	-2.88	112.12	120.87
4	B	606	A1A3C	O34-C33-C30	2.81	119.95	115.14
4	A	608	A1A3C	O34-C33-C36	-2.80	119.26	124.08
4	A	608	A1A3C	O34-C33-C30	2.60	119.59	115.14
4	B	606	A1A3C	C17-C16-N15	-2.51	113.00	120.43
4	A	608	A1A3C	O01-C02-C10	-2.48	116.00	120.90
4	B	606	A1A3C	N37-C24-N15	-2.44	108.95	115.06
4	A	608	A1A3C	C17-C16-N15	-2.42	113.25	120.43
4	B	606	A1A3C	C11-C10-C38	2.37	122.08	119.21
4	A	608	A1A3C	C25-C24-N37	2.32	126.21	122.68
4	A	608	A1A3C	C38-C14-C13	2.31	122.83	120.54
4	A	608	A1A3C	O22-C19-C18	-2.29	118.66	122.19
4	A	608	A1A3C	C21-N20-C19	2.10	125.87	122.21
4	A	608	A1A3C	C11-C10-C02	2.06	127.29	120.60

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	605	GOL	O1-C1-C2-C3
4	A	608	A1A3C	C17-C18-C19-N20
4	A	608	A1A3C	C17-C18-C19-O22
4	A	608	A1A3C	C23-C18-C19-N20
4	A	608	A1A3C	C23-C18-C19-O22
4	A	608	A1A3C	C18-C19-N20-C21
4	A	608	A1A3C	O22-C19-N20-C21

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	608	A1A3C	O01-C02-N03-C04
4	A	608	A1A3C	C10-C02-N03-C04
3	A	607[B]	GOL	C1-C2-C3-O3
4	A	608	A1A3C	C05-C06-N07-C09
3	B	605	GOL	O1-C1-C2-O2
3	A	607[B]	GOL	O2-C2-C3-O3
4	A	608	A1A3C	C04-C05-C06-N07
4	A	608	A1A3C	C05-C06-N07-C08
4	B	606	A1A3C	C23-C18-C19-O22
4	B	606	A1A3C	C30-C27-O28-C29
4	B	606	A1A3C	C23-C18-C19-N20
4	A	608	A1A3C	N03-C04-C05-C06


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	PO4	1	0
2	A	602	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	307/317 (96%)	0.05	16 (5%) 34 35	17, 41, 84, 150	4 (1%)
1	B	300/317 (94%)	0.23	17 (5%) 30 32	19, 48, 97, 159	5 (1%)
All	All	607/634 (95%)	0.14	33 (5%) 32 34	17, 44, 97, 159	9 (1%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	432[A]	TYR	5.0
1	B	200	VAL	4.7
1	B	204	VAL	3.8
1	A	196	LEU	3.8
1	B	205	ALA	3.6
1	A	370	VAL	3.5
1	B	273[A]	ARG	3.1
1	A	368	LEU	3.1
1	B	498	ILE	3.0
1	B	201	GLN	3.0
1	A	371	GLY	3.0
1	A	365	THR	2.8
1	B	463	SER	2.8
1	B	227	TRP	2.7
1	A	256	MET	2.7
1	A	252	TYR	2.7
1	B	202	ARG	2.6
1	B	203	THR	2.5
1	A	284[A]	HIS	2.5
1	A	200	VAL	2.5
1	A	255	VAL	2.5
1	B	257	LEU	2.5
1	A	197	PRO	2.4
1	B	497	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	370	VAL	2.4
1	B	365	THR	2.4
1	A	202	ARG	2.4
1	B	499	ASP	2.4
1	A	198	PHE	2.3
1	B	271	THR	2.2
1	A	194	SER	2.2
1	A	199	LEU	2.0
1	B	470	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	607[A]	6/6	0.73	0.14	44,53,59,59	14
3	GOL	A	607[B]	6/6	0.73	0.14	44,52,57,59	14
3	GOL	B	605	6/6	0.87	0.10	38,58,74,75	0
2	PO4	A	606	5/5	0.91	0.11	40,41,52,52	0
2	PO4	A	605	5/5	0.92	0.09	37,52,62,65	0
4	A1A3C	A	608	38/38	0.93	0.10	22,38,68,69	0
4	A1A3C	B	606	38/38	0.94	0.08	23,36,63,76	0
2	PO4	B	604	5/5	0.96	0.08	36,36,41,41	0
2	PO4	B	601	5/5	0.97	0.06	28,37,42,44	0
2	PO4	B	603	5/5	0.97	0.07	33,36,39,40	0
2	PO4	A	602	5/5	0.97	0.06	33,35,37,42	0
2	PO4	A	604	5/5	0.97	0.06	37,39,45,48	0
2	PO4	A	603	5/5	0.98	0.06	27,32,34,36	0
2	PO4	B	602	5/5	0.98	0.04	35,36,39,42	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PO4	A	601	5/5	0.98	0.05	33,33,37,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.